



## Supplementary Material

# Prevalence, Haematological Alterations and Chemotherapy of Bovine Anaplasmosis in Sahiwal and Crossbred Cattle of District Faisalabad, Punjab, Pakistan

Aftab Shaukat<sup>1\*</sup>, Khalid Medmood<sup>3</sup>, Irfan Shaukat<sup>2</sup>, Tauseef ur Rehman<sup>3</sup>, Muhammad Ahsan Naem<sup>1,5</sup>, Ashar Mehfooz<sup>1</sup>, Muhammad Ijaz Saleem<sup>1</sup>, Zia-ud-Din Sindhu<sup>1</sup>, Shahid Ali Rajput<sup>4</sup>, Mubashar Hassan<sup>1,4</sup>, Saqib Umar<sup>1</sup>, Muhammad Ali Jamil<sup>1</sup>, Rao Zahid Abbas<sup>1</sup> and Anas Sarwar Qureshi<sup>1</sup>

<sup>1</sup>Faculty of Veterinary Sciences, University of Agriculture, Faisalabad

<sup>2</sup>Faculty of medicine, University of Lorraine, Nancy, France

<sup>3</sup>University College of Veterinary and Animal Sciences, The Islamia University of Bahawalpur

<sup>4</sup>College of Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China

<sup>5</sup>College of Veterinary and Animal Sciences (Narowal), University of Veterinary and Animal Sciences, Lahore, Pakistan

\* Corresponding author: aftabshaukat40@gmail.com

0030-9923/2019/0006-2023 \$ 9.00/0

Copyright 2019 Zoological Society of Pakistan

### Annexure

#### Title of project:

Date of survey: \_\_\_\_\_ Performa #: \_\_\_\_\_ Name of Owner: \_\_\_\_\_

Location /Address: \_\_\_\_\_

#### Breed

Sahiwal	Cross bred

#### Gender

Male	Female

#### Farming system

Separate	Mixed

#### Hygienic measure

Poor	Good	Excellent

#### Age group

<1 Year	1-2Year	>2-4 Years	>4 Years

#### Feeding System

Grazing	Stall Feeding	both

#### Floor Pattern

Un-cemented	Cemented	Partially-cemented

#### Herd size

Medium	large

#### Frequency of Acaricidal treatment

Once/year	Twice/year	Thrice/year	None

#### Number of cleaning time

Once a day	Twice a week	Once a week

Supplementary Fig. 1. Questionnaire designed for collection of information about *A. marginale* in cattle and associated determinants.