



## Supplementary Material

# Prevalence, Haematological Alterations and Chemotherapy of Bovine Anaplasmosis in Sahiwal and Crossbred Cattle of District Faisalabad, Punjab, Pakistan

Aftab Shaukat<sup>1\*</sup>, Khalid Mehmood<sup>3</sup>, Irfan Shaukat<sup>2</sup>, Tauseef ur Rehman<sup>3</sup>, Muhammad Ahsan Naem<sup>1,5</sup>, Ashar Mehfooz<sup>1</sup>, Muhammad Ijaz Saleem<sup>1</sup>, Zia-ud-Din Sindhu<sup>1</sup>, Shahid Ali Rajput<sup>4</sup>, Mubashar Hassan<sup>1,4</sup>, Saqib Umar<sup>1</sup>, Muhammad Ali Jamil<sup>1</sup>, Rao Zahid Abbas<sup>1</sup> and Anas Sarwar Qureshi<sup>1</sup>

<sup>1</sup>Faculty of Veterinary Sciences, University of Agriculture, Faisalabad

<sup>2</sup>Faculty of medicine, University of Lorraine, Nancy, France

<sup>3</sup>University College of Veterinary and Animal Sciences, The Islamia University of Bahawalpur

<sup>4</sup>College of Veterinary Medicine, Huazhong Agricultural University, Wuhan 430070, China

<sup>5</sup>College of Veterinary and Animal Sciences (Narowal), University of Veterinary and Animal Sciences, Lahore, Pakistan

\* Corresponding author: aftabshaukat40@gmail.com

0030-9923/2019/0006-2023 \$ 9.00/0

Copyright 2019 Zoological Society of Pakistan

### Annexure

#### Title of project:

Date of survey: \_\_\_\_\_ Performa #: \_\_\_\_\_ Name of Owner: \_\_\_\_\_

Location /Address: \_\_\_\_\_

Breed		Gender		Farming system	
Sahiwal	Cross bred	Male	Female	Separate	Mixed

  

Hygienic measure			Age group			
Poor	Good	Excellent	<1 Year	1-2Year	>2-4 Years	>4 Years

  

Feeding System			Floor Pattern		
Grazing	Stall Feeding	both	Un-cemented	Cemented	Partially-cemented

  

Herd size		Frequency of Acaricidal treatment			
Medium	large	Once/year	Twice/year	Thrice/year	None

  

Number of cleaning time		
Once a day	Twice a week	Once a week

Supplementary Fig. 1. Questionnaire designed for collection of information about *A. marginale* in cattle and associated determinants.