



Supplementary Material

Optimum Conditions for Enhanced Production of Podophyllotoxin from *Penicillium* sp. Isolated from Khanspur, Pakistan

Umar Farooq Gohar, Attia Majeed, Bushra Muneer* and Hamid Mukhtar


Institute of Industrial Biotechnology, Government College University, Lahore, Pakistan 54000.

Supplementary Table I. Isolation and screening of the fungal isolate for the production of podophyllotoxin through surface culture fermentation.


No. of samples	Fungal isolates	Amount of Podophyllotoxin produced ($\mu\text{g/ml}$)
MR-1	P*-15	10.799
	P-16	7.64
	P-18	88.14
	P-8	54.0
	P-1	3.8
	P-2	7.31
	P-3	5.68
MR-2	P-13	1.67
	P-12	2.63
MR-4	P-5	49.77
	P-7	14.71
KH-2	P-4	0.043
	P-6	0.65
	P-17	3.64
KH-3	P-19	7.9
	P-9	6.14
	P-10	12.8
MK-1	P-20	22.05
	P-14	3.64
MK-3	P-11	21.92

*MR, Murree; KH, Khanspur; MK, Mushkpuri. *P stands for podophyllotoxin.

* Corresponding author: bushramuneer11@yahoo.co.uk, dr.bushramuneer@gcu.edu.pk
0030-9923/2022/0006-2775 \$ 9.00/0



Supplementary Fig. 1. HPLC chromatogram of podophyllotoxin (A) Standard (B) Sample.



Supplementary Fig. 2. Colony of isolated fungal strain P-18.



Copyright 2022 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).



Supplementary Fig. 3. The sporangia and spore of isolated fungal strain P-18.