



# Bioaccumulation of Cu and Zn in *Schizothorax plagiostomus* and *Mastacembelus armatus* from River Swat, River Panjkora and River Barandu in Malakand Division, Pakistan

Hazrat Ali<sup>1\*</sup>, Wajid Ali<sup>1</sup>, Karim Ullah<sup>1</sup>, Fazle Akbar<sup>1</sup>, Sher Ahrar<sup>1</sup>, Irfan Ullah<sup>1</sup>, Irum Ahmad<sup>1</sup>, Ashfaq Ahmad<sup>1</sup>, Ikram Ilahi<sup>2</sup> and Muhammad Anwar Sajad<sup>3</sup>

<sup>1</sup>Green and Environmental Chemistry, Ecotoxicology and Ecology Laboratory, Department of Zoology, University of Malakand, Chakdara 18800, Dir Lower, Khyber Pakhtunkhwa, Pakistan

<sup>2</sup>Department of Zoology, University of Malakand, Chakdara 18800, Dir Lower, Khyber Pakhtunkhwa, Pakistan

<sup>3</sup>Department of Botany, Islamia College University Peshawar, Khyber Pakhtunkhwa, Pakistan

## ABSTRACT

Contamination of aquatic ecosystems with heavy metals leads to accumulation of these elements in aquatic biota including fish. Bioaccumulation of heavy metals in fish is of public health concern because fish are an important part of human diet. The present study was conducted to investigate the bioaccumulation of two heavy metals, Cu and Zn, in muscles of two economically important fish species, an herbivorous fish *Schizothorax plagiostomus* and a predatory fish *Mastacembelus armatus*, from three rivers of Malakand Division, Pakistan. Concentrations of Cu and Zn were determined in fish samples by atomic absorption spectrophotometer. Highest Cu concentration of  $4.52 \pm 0.24$  mg kg<sup>-1</sup> wet weight was observed in muscles of *M. armatus* at Chakdara on River Swat while highest Zn concentration of  $18.00 \pm 1.88$  mg kg<sup>-1</sup> wet weight was observed in muscles of *S. plagiostomus* at Daggar on River Barandu. A site wise comparison of both Cu and Zn concentrations in muscles of the study fish generally showed no significant differences among different sites on the rivers. However, in some cases, metal concentrations showed significant increase down the stream. Similarly, a comparison of metal concentrations in muscles of the two species generally did not show significant differences between the two species. Zn concentrations in fish muscles were higher than Cu concentrations in all cases. Bioaccumulation factor (BAF) values for Cu were in the order of kidneys > muscles > liver > gills > skin while for Zn the order was skin > kidneys > liver > gills > muscles.

## Article Information

Received 25 February 2017

Revised 22 March 2017

Accepted 12 May 2017

Available online 10 August 2017

## Authors' Contribution

HA and II conceived and designed the study. HA, WA, KU, FA, SA, IU, IA, AA and MAS collected, prepared and analyzed the samples. HA analyzed the data and wrote the manuscript. HA supervised the research work.

## Key words

Bioaccumulation, Heavy metal, *Mastacembelus armatus*, *Schizothorax plagiostomus*.

## INTRODUCTION

Environmental pollution is one of the major challenges faced by human society in the 21<sup>st</sup> century (Ali and Khan, 2017). Heavy metal pollution in aquatic ecosystems has become an issue of global concern due to industrialization (Javed and Usmani, 2016). Heavy metals are released to the environment from different natural and anthropogenic sources. Due to their non-biodegradable and persistent nature, they accumulate in the environment and subsequently accumulate in biota. Accumulation of

toxic heavy metals in biota leads to contamination of the food chains/webs with consequent risk to wildlife and human health. Aquatic animals are at more risk from toxic heavy metals in their habitats because they cannot escape from such contamination. Monitoring of heavy metal contamination in fish is important because fish constitute an important food source for humans. In routine monitoring of environmental contamination, fish muscle is the main tissue of interest due to their consumption by humans (Siraj *et al.*, 2014, 2016; Ambreen *et al.*, 2015). Fish are important bioindicators of water quality (Hussain *et al.*, 2016).

Fish consumption is recommended in human diet because they are a good source of proteins and unsaturated fatty acids. However, due to contamination of aquatic ecosystems with toxic heavy metals, it becomes necessary

\* Corresponding author: [hazratali@uom.edu.pk](mailto:hazratali@uom.edu.pk); [hazrataliuom@gmail.com](mailto:hazrataliuom@gmail.com)  
0030-9923/2017/0005-1555 \$ 9.00/0  
Copyright 2017 Zoological Society of Pakistan

to monitor contents of these toxic elements in fish. Scientific interest from environmental point of view in essential heavy metals such as Mn, Fe, Cu and Zn is due to the narrow window between their essentiality and toxicity (Yousafzai and Shakoori, 2008). Essential heavy metals are required for proper functioning of life processes but either deficiency or excess of these metals in the body creates health problems (Tabinda *et al.*, 2010, 2013). Although, Cu and Zn are essential elements but their excessive levels in fish can pose health risk to the consumer human population and to the exposed fish themselves. Excess of Cu in the body is associated with damage of the liver (Javed and Usmani, 2016). The Provisional Maximum Tolerable Daily Intake (PMTDI) of Cu and Zn given in Codex Alimentarius Commission are 0.05-0.5 mg/kg bw and 0.3-1 mg/kg bw, respectively (FAO/WHO, 2011). Some studies have investigated the effects of Cu on different biochemical parameters of fish (*e.g.*, Mutlu *et al.*, 2016). In fish, Cu can induce respiratory distress (Javed and Usmani, 2011) and interferes with osmoregulation; death is caused by tissue hypoxia associated with disrupted ATP synthesis (Eisler, 1998). Cu pollution in natural waters may severely damage the gills and cause histological changes in liver and kidneys (Javed and Usmani, 2013). It has been reported that exposure to Cu induced DNA damage in the gill cells of the freshwater mussels (*Anodonta anatina*) and the genotoxic effects were more at an exposure dose of 120  $\mu\text{g L}^{-1}$  (Sohail *et al.*, 2017). Excess Zn is teratogenic to embryos of frogs and fish (Eisler, 1993).

Riverine pollution has been reported as one of the possible reasons for the rapid population decline of the endangered freshwater fish *Tor puitora* in River Kabul, Pakistan (Yousafzai and Shakoori, 2009). Recently, marble industry effluents were associated with lower species abundance and diversity in River Barandu, Pakistan (Mulk *et al.*, 2016). In Pakistan, the major sources of heavy metals in the aquatic environment are agricultural runoff, domestic sewage and industrial effluents (Hussain *et al.*, 2014). The aim of the present research work was to investigate the bioaccumulation of two heavy metals Cu and Zn in muscles of two economically important fish species, *Schizothorax plagiostomus* and *Mastacembelus armatus* from three rivers of Malakand Division, Pakistan. We selected fish muscles for analysis of the heavy metals because muscles are the edible part of fish. Accumulation of toxic chemicals in edible tissues of fish is of public health concern. Fish skin is also mostly consumed with muscles and is therefore important on bioaccumulation point of view (Yousafzai *et al.*, 2010). However, fish muscles are usually analyzed for determining the levels of contaminants (Mortuza and Al-Misned, 2015).

## MATERIALS AND METHODS

### Study area

The study area of this research work included three rivers of Malakand Division *i.e.*, River Swat, River Panjkora and River Barandu. River Swat and River Panjkora meet at Bosaq Bridge, Sharbatti. Regarding ichthyofaunal diversity, River Swat has relatively more diversity compared to River Panjkora (Ullah *et al.*, 2014). Discharge of effluents causes heavy metal contamination in River Panjkora as observed from slight increase in heavy metal concentrations downstream (Ahmad *et al.*, 2014). River Barandu is also being polluted by effluents from marble industries in Buner (Khan *et al.*, 2012).

### Water samples

Water samples were collected according to Csuros and Csuros (2002) and transported to the laboratory in pre-cleaned polyethylene bottles. After filtration through a Whatman filter paper, a 50 mL sample was taken in a 100 mL beaker and acidified with 5 mL concentrated (65%)  $\text{HNO}_3$ . After acidification, the water samples were heated on a hot plate at 80°C until the volume was reduced to 50 mL (Malik and Maurya, 2014). These water samples were analyzed for Cu and Zn by Flame Atomic Absorption Spectrophotometer (Perkin-Elmer Model No. 2380).

### Fish samples

This study investigated bioaccumulation of Cu and Zn in a herbivorous snow carp, *Schizothorax plagiostomus* and a carnivorous *Mastacembelus armatus*. These fish species were selected because of their commercial importance in the study area. Furthermore, these species have different feeding habits. Dietary habits may affect metal bioaccumulation in different fish species (Papagiannis *et al.*, 2004). Therefore, one aim of the study was to compare metal bioaccumulation in two fish species with different feeding habits.

Three samples each of *S. plagiostomus* and *M. armatus* (where available) were collected at each pre-selected site on River Swat, River Panjkora and River Barandu. Fish muscles were collected according to Rosseland *et al.* (2017). Skin-free muscle samples were taken from dorsal side behind the dorsal fin with the help of clean knife and scissors. In case of tissue-specific metal bioaccumulation study of *M. armatus*, other tissues *i.e.*, gills, skin, liver and kidneys were also separated. Fish samples were digested according to Javed and Usmani (2016) with modifications. A sample of 1 g (wet weight basis) was taken in a clean beaker. To this sample, 7.5 mL  $\text{HNO}_3$  (65%) and 2.5 mL  $\text{HClO}_4$  (70%) were added and heated on a hot plate at 80°C. The sample was heated until the evolution of brown fumes

of NO<sub>2</sub> ceased and a clear (yellow) solution was obtained. The solution was cooled at room temperature and then filtered through a Whatman filter paper. The filtrate was diluted to 50 mL with analyte-free water. The heavy metals Cu and Zn were determined in these samples with Flame Atomic Absorption Spectrophotometer (Perkin-Elmer Model No. 2380).

For quality control check was performed in order to assure the accuracy of the metal analysis. The recoveries were obtained for the analyzed metals from reference materials (standards) (Table I).

**Table I.- Recoveries obtained for the analyzed metals from reference materials.**

Metal	Concentration (ppm)		Recovery (%)
	Measured	Standard	
Cu	4.998	5.000	99.96
Zn	0.997	1.000	99.70

#### Statistical analysis

Metal concentrations are shown as mean  $\pm$  standard deviation with  $n = 3$ . Experimental data were analyzed

with IBM SPSS Statistics 23. Mean metal concentrations at different river sites were compared using One-Way ANOVA (Tukey Test) while mean metal concentrations of the two fish species were compared using Independent-Samples T Test.  $P$  value of 0.05 was considered for statistical significance.

## RESULTS AND DISCUSSION

Concentrations of Cu and Zn in muscles of *S. plagiostomus* and *M. armatus* collected from different sites of River Swat, River Panjkora and River Barandu are given in Table II. Cu and Zn concentrations in muscles of *S. plagiostomus* at different sites of River Swat did not show significant differences (Table II). However, Cu concentration in muscles of *M. armatus* ( $4.52 \pm 0.24 \text{ mg kg}^{-1} \text{ ww}$ ) was significantly higher than that of *S. plagiostomus* ( $1.47 \pm 0.64 \text{ mg kg}^{-1} \text{ ww}$ ) at Chakdara site. In case of River Panjkora, Cu and Zn concentrations in muscles of *S. plagiostomus* increased down the stream (Table II). Furthermore, Zn concentration in muscles of *M. armatus* ( $7.28 \pm 1.71 \text{ mg kg}^{-1} \text{ ww}$ ) was significantly higher than that of *S. plagiostomus* ( $3.30 \pm 1.26 \text{ mg kg}^{-1} \text{ ww}$ ) at Khazana site.

**Table II.- Cu and Zn concentrations in muscles of *Schizothorax plagiostomus* and *Mastacembelus armatus* collected from River Swat, River Panjkora and River Barandu.**

Site	Metal concentration (mg kg <sup>-1</sup> wet weight)			
	Cu		Zn	
	<i>S. plagiostomus</i>	<i>M. armatus</i>	<i>S. plagiostomus</i>	<i>M. armatus</i>
<b>River Swat</b>				
Khwazakhela	$1.35 \pm 0.52^a$	NA	$12.12 \pm 2.19^a$	NA
Kabal	$1.07 \pm 0.36^a$	$0.62 \pm 0.10$	$9.45 \pm 1.15^a$	$8.95 \pm 3.54$
Chakdara	$1.47 \pm 0.64^a$	$4.52 \pm 0.24^{**}$	$7.53 \pm 3.48^a$	$12.35 \pm 1.48$
Batkhehla	$2.05 \pm 1.00^a$	NA	$12.80 \pm 1.08^a$	NA
<b>River Panjkora</b>				
Khall	$0.93 \pm 0.50$	NA	$2.72 \pm 2.36^b$	NA
Khazana	$2.18 \pm 0.46$	$3.92 \pm 2.77$	$3.30 \pm 1.26^b$	$7.28 \pm 1.71^*$
Shagukas	$2.72 \pm 1.15$	NA	$10.32 \pm 2.99^a$	NA
<b>River Barandu</b>				
Toor Warsak	$2.20 \pm 0.74^b$	$1.67 \pm 1.50$	$16.38 \pm 4.23$	$11.98 \pm 2.69$
Elai	$2.48 \pm 0.39^{ab}$	$1.13 \pm 0.80$	$10.17 \pm 4.79$	$11.58 \pm 2.50$
Daggar	$2.02 \pm 0.33^b$	$1.83 \pm 1.24$	$18.00 \pm 1.88$	$13.12 \pm 3.39$
Dewana Baba	$4.05 \pm 0.91^{a*}$	$2.38 \pm 0.41$	$12.08 \pm 3.83$	$12.27 \pm 4.20$

NA, not available (species sample); Results are shown as mean  $\pm$  standard deviation ( $n = 3$ ). Mean values in a column with same superscript letter are not significantly different at  $p \leq 0.05$  (One Way ANOVA, Tukey Test). \*\* Means of the two species significantly different at  $p \leq 0.01$  (Independent-Samples T Test).

The trend of increase down the stream in concentrations of both metals in muscles of *S. plagiostomus* is shown in Figure 1. Ullah *et al.* (2016) have also reported higher concentrations of heavy metals in three fish species from River Panjkora at Timergara compared to those at Khall. In case of River Barandu, Cu and Zn concentrations in muscles of the two fish species did not show significant differences at different river sites except Cu concentrations in muscles of *S. plagiostomus*. However, Zn concentrations were higher than Cu at all sites. Furthermore, at Dewana Baba site, Cu concentration in muscles of *S. plagiostomus* ( $4.05 \pm 0.91 \text{ mg kg}^{-1} \text{ ww}$ ) was significantly higher than that of *M. armatus* ( $2.38 \pm 0.41 \text{ mg kg}^{-1} \text{ ww}$ ). At other river sites, Cu as well as Zn concentrations in muscles of the two fish species did not differ significantly. Zn concentrations in field collections of biota are extremely variable and difficult to interpret (Eisler, 1993).




Fig. 1. Cu and Zn concentrations ( $\text{mg kg}^{-1}$  wet weight) in muscles of *Schizothorax plagiostomus* at different sites of River Panjkora.

#### Copper (Cu)

In this study, Cu concentrations in muscles of *S. plagiostomus* and *M. armatus* from River Swat ranged from  $1.07 \pm 0.36$  to  $2.05 \pm 1.00 \text{ mg kg}^{-1} \text{ ww}$  and  $0.62 \pm 0.10$  to  $4.52 \pm 0.24 \text{ mg kg}^{-1} \text{ ww}$ , respectively. Corresponding Cu concentrations from River Panjkora ranged from  $0.93 \pm 0.50$  to  $2.72 \pm 1.15 \text{ mg kg}^{-1} \text{ ww}$  and  $3.92 \pm 2.77 \text{ mg kg}^{-1} \text{ ww}$  respectively while corresponding Cu concentrations from River Barandu ranged from  $2.02 \pm 0.33$  to  $4.05 \pm 0.91 \text{ mg kg}^{-1} \text{ ww}$  and  $1.13 \pm 0.80$  to  $2.38 \pm 0.41 \text{ mg kg}^{-1} \text{ ww}$ , respectively. Ngelinkoto *et al.* (2014) have reported highest Cu concentration of  $15.7 \text{ mg kg}^{-1} \text{ ww}$  in muscles of some fish species from Kwilu-Ngongo River, Congo. In a recent study, Javed and Usmani (2016) have reported Cu concentration of  $41.36 \pm 0.38 \text{ mg kg}^{-1} \text{ dw}$  in muscles of *M. armatus* collected from an effluent-loaded canal in India. A comparison of our results with those of the above two studies shows that our reported Cu concentrations are less than those of the above mentioned studies.

#### Zinc (Zn)

In this study, Zn concentrations in muscles of *S. plagiostomus* and *M. armatus* from River Swat ranged from  $7.53 \pm 3.48$  to  $12.80 \pm 1.08 \text{ mg kg}^{-1} \text{ ww}$  and  $8.95 \pm 3.54$  to  $12.35 \pm 1.48 \text{ mg kg}^{-1} \text{ ww}$ , respectively. Corresponding Zn concentrations from River Panjkora ranged from  $2.72 \pm 2.36$  to  $10.32 \pm 2.99 \text{ mg kg}^{-1} \text{ ww}$  and  $7.28 \pm 1.71 \text{ mg kg}^{-1} \text{ ww}$ , respectively while corresponding Zn concentrations from River Barandu ranged from  $10.17 \pm 4.79$  to  $18.00 \pm 1.88 \text{ mg kg}^{-1} \text{ ww}$  and  $11.58 \pm 2.50$  to  $13.12 \pm 3.39 \text{ mg kg}^{-1} \text{ ww}$ , respectively. Ngelinkoto *et al.* (2014) have reported highest Zn concentration of  $45.3 \text{ mg kg}^{-1} \text{ ww}$  in muscles of some fish species from Kwilu-Ngongo River, Congo. In a recent study, Javed and Usmani (2016) have reported Zn concentration of  $186.19 \pm 0.12 \text{ mg kg}^{-1} \text{ dw}$  in muscles of *M. armatus* collected from an effluent-loaded canal in India. A comparison of our results with those of the above two studies shows that our reported Zn concentrations are less than those of the above mentioned studies.




Fig. 2. Cu and Zn concentrations ( $\text{mg kg}^{-1}$  wet weight) in different tissues of *Mastacembelus armatus* collected at Khazana on River Panjkora.

#### Tissue-specific bioaccumulation of Cu and Zn in *M. armatus*

We also determined Cu and Zn concentrations in five different tissues of *M. armatus* collected at Khazana site on River Panjkora. Results of such tissue-specific bioaccumulation of the two heavy metals in *M. armatus* are shown in Figure 2. From Figure 2, it can be seen that the order of accumulation of Cu was: kidneys > muscles > liver > gills > skin while for Zn the order was: skin > kidneys > liver > gills > muscles. Among fish tissues, liver accumulates relatively more quantities of heavy metals (Vinodhini and Narayanan, 2008).

In our study, liver was at intermediate position in concentrations of Cu and Zn. Recently, Ullah *et al.* (2016) have reported the order of bioaccumulation of heavy metals in different tissues of three fish species from River

**Table III.- Bioaccumulation factor (BAF) values of Cu and Zn in different tissues of *Mastacembelus armatus* at Khazana on River Panjkora.**

Heavy metal	Conc. in water (mg L <sup>-1</sup> )	BAF				
		Muscles	Gills	Skin	Liver	Kidneys
Cu	0.022	178.18	78.64	50.91	80.91	535.91
Zn	0.022	330.91	474.09	1017.27	603.64	987.73

Panjkora, Pakistan as: liver > kidney > muscles > gills. Kidney is the principal tissue for the storage of metals and the main excretion route for toxicants (Palaniappan and Karthikeyan, 2009). Malik *et al.* (2014) have also reported relatively higher concentrations of heavy metals in liver, kidneys and gills compared to those in skin and muscles of four edible fish species *i.e.*, *Tor putitora*, *Cirrhinus mrigala*, *Labeo calbasu* and *Channa punctatus* from Rawal Lake Reservoir, Pakistan.

In a recent study, Rosseland *et al.* (2017) have reported Cu concentrations of 2.5-5.2 and 7-1697  $\mu\text{g g}^{-1}$  dw in gills and liver, respectively of *M. armatus* from Lake Phewa, Nepal. The corresponding Zn concentrations in gills and liver were 93-189 and 38-213  $\mu\text{g g}^{-1}$  dw, respectively. Our reported Cu and Zn concentrations in gills and liver of *M. armatus* are less than those reported by the above mentioned study. Yousafzai and Shakoori (2008) have reported Cu concentrations of  $44.70 \pm 3.39$ ,  $67.70 \pm 2.38$  and  $76.70 \pm 4.82$   $\mu\text{g g}^{-1}$  ww in gills of *Tor putitora* collected from control and two polluted sites (site 1 and site 2), respectively of River Kabul, Pakistan. Their reported corresponding Zn concentrations were  $1993.90 \pm 126.70$ ,  $2124.90 \pm 43.81$  and  $2414.00 \pm 70.08$   $\mu\text{g g}^{-1}$  ww. Our reported Cu and Zn concentrations in gills are far less than those reported by the above mentioned study. The reason for low concentrations of the two heavy metals in different tissues of *M. armatus* in our study may be the fact that River Panjkora is less polluted than River Kabul.

#### Bioaccumulation factor

Bioaccumulation factor (BAF) is an index of the degree of accumulation of a contaminant in an organism relative to its environment. It is calculated by the following equation (Mortuza and Al-Misned, 2015):

$$BAF = \frac{\text{Conc. of metal in organism tissue}}{\text{Conc. of metal in abiotic medium}}$$

The BAF values of Cu and Zn in different tissues of *M. armatus* are given in Table III. From Table III, it is seen that BAF values of Zn are greater than those of Cu. Javed and Usmani (2013) have studied bioaccumulation of heavy metals in *M. armatus* collected from a rivulet

at Kasimpur, Aligarh, India. Their reported BAF values of Cu in muscles, gills, skin, liver and kidneys of *M. armatus* were 48.09, 232.41, 42.17, 315.89, and 204.52, respectively while corresponding BAF values of Zn were 620.63, 1831.10, 560.36, 5806.50, and 1170.93. These results are comparable to our reported BAF values of Cu and Zn in different tissues of *M. armatus*.

## CONCLUSIONS

The present study investigated the bioaccumulation of two heavy metals, Cu and Zn, in muscles of two economically important fish species *S. plagiostomus* and *M. armatus* from three rivers of Malakand Division, Pakistan. Generally, the concentrations of the two heavy metals in muscles of the study fish did not differ significantly at different sites of the rivers. However, in some cases Cu and Zn concentrations showed significant differences at different sites of the rivers. For example, in case of River Panjkora, Zn concentrations and in case of River Barandu Cu concentrations (except at Daggar) in muscles of *S. plagiostomus* increased down the stream. Regarding feeding habits of the two fish species, generally Cu and Zn concentrations did not show significant differences between the two species. However, in two cases, *M. armatus* accumulated more metals while in one case, *S. plagiostomus* showed more bioaccumulation of Cu. Thus, no clear trend could be noted in bioaccumulation of Cu and Zn regarding feeding habits of the study fish.

## ACKNOWLEDGEMENTS

The authors are thankful to Prof. Dr. Rup Lal, Professor at Department of Zoology, University of Delhi, India, for gifting a copy of standard literature keys for fish identification and to Mr. Abdur Rahman, Lecturer in Zoology, and Fisheries Specialist at Department of Zoology, University of Malakand, for his help in fish identification.

#### Statement of conflict of interest

The authors declare that there is no conflict of interests regarding the publication of this article.

## REFERENCES

- Ahmad, K., Azizullah, A., Shama, S. and Khattak, M.N.K., 2014. Determination of heavy metal contents in water, sediments, and fish tissues of *Shizothorax plagiostomus* in river Panjkora at Lower Dir, Khyber Pakhtunkhwa, Pakistan. *Environ. Monit. Assess.*, **186**: 7357-7366. <https://doi.org/10.1007/s10661-014-3932-1>
- Ali, H. and Khan, E., 2017. Environmental chemistry in the twenty-first century. *Environ. Chem. Lett.*, **15**: 329-346. <https://doi.org/10.1007/s10311-016-0601-3>
- Ambreen, F., Javed, M. and Batool, U., 2015. Tissue specific heavy metals uptake in economically important fish, *Cyprinus carpio* at acute exposure of metals mixtures. *Pakistan J. Zool.*, **47**: 399-407.
- Csuros, M. and Csuros, C., 2002. *Environmental sampling and analysis for metals*. Lewis Publishers, Boca Raton. <https://doi.org/10.1201/9781420032345>
- Eisler, R., 1993. *Zinc hazard to fish, wildlife, and invertebrates: A synoptic review*. U.S. Fish and Wildlife Service Biological Report 10, pp. 106.
- Eisler, R., 1998. *Copper hazards to fish, wildlife, and invertebrates: A synoptic review*. U.S. Geological Survey, Biological Resources Division, Biological Science Report USGS/BRD/BSR--1997-0002, pp. 98.
- FAO/WHO, 2011. *Joint FAO/WHO food standards programme Codex Committee on Contaminants in Food*. Codex Alimentarius Commission.
- Hussain, M., Muhammad, S., Malik, R.N., Khan, M.U. and Farooq, U., 2014. Status of heavy metal residues in fish species of Pakistan. In: *Reviews of environmental contamination and toxicology* (ed. D.M. Whitacre). Springer, pp. 111-132. [https://doi.org/10.1007/978-3-319-04411-8\\_5](https://doi.org/10.1007/978-3-319-04411-8_5)
- Hussain, M.Z., Latif, A., Ahmed, S.W., Hussain, S., Iqbal, R. and Ali, M., 2016. Diversity, abundance and seasonal variations of fish community in lentic water bodies of Indus River at Ghazi Ghat, Pakistan. *Pakistan J. Zool.*, **48**: 59-65.
- Jan, A., Rab, A., Ullah, R. and Haroon, U.I., 2017. On the breeding season of *Schizothorax plagiostomus*. *Pakistan J. Zool.*, **49**: 1-3. <https://doi.org/10.17582/journal.pjz/2017.49.2.sc8>
- Jan, M. and Ahmed, I., 2016. Length weight relationship and condition factor of snow trout, *Schizothorax plagiostomus* (Heckel, 1838) from Lidder River, Kashmir. *Int. J. Fish. Aquat. Stud.*, **4**: 131-136.
- Javed, M. and Usmani, N., 2011. Accumulation of heavy metals in fishes: A human health concern. *Int. J. Environ. Sci.*, **2**: 671-682.
- Javed, M. and Usmani, N., 2013. Assessment of heavy metal (Cu, Ni, Fe, Co, Mn, Cr, Zn) pollution in effluent dominated rivulet water and their effect on glycogen metabolism and histology of *Mastacembelus armatus*. *SpringerPlus*, **2**: Article No. 390. <https://doi.org/10.1186/2193-1801-2-390>
- Javed, M. and Usmani, N., 2016. Accumulation of heavy metals and human health risk assessment via the consumption of freshwater fish *Mastacembelus armatus* inhabiting, thermal power plant effluent loaded canal. *SpringerPlus*, **5**: Article No. 776. <https://doi.org/10.1186/s40064-016-2471-3>
- Khan, Z.R., Haq, F., Ullah, R. and Ullah, M.I., 2012. The analysis of water samples in different industrialization units of District Buner, Khyber Pakhtunkhwa, Pakistan. *Int. J. Biosci.*, **2**: 82-85.
- Malik, D.S. and Maurya, P.K., 2014. Heavy metal concentration in water, sediment, and tissues of fish species (*Heteropneustis fossilis* and *Puntius ticto*) from Kali River, India. *Toxicol. Environ. Chem.*, **96**: 1195-1206.
- Malik, R.N., Hashmi, M.Z. and Huma, Y., 2014. Heavy metal accumulation in edible fish species from Rawal Lake Reservoir, Pakistan. *Environ. Sci. Pollut. Res.*, **21**: 1188-1196. <https://doi.org/10.1007/s11356-013-1992-3>
- Mirza, M.R., 2006. A note on the fishes of Allai Khoar, NWFP, Pakistan. *Punjab Univ. J. Zool.*, **21**: 73-75.
- Mortuza, M.G. and Al-Misned, F.A., 2015. Heavy metal concentration in two freshwater fishes from Wadi Hanifah (Riyadh, Saudi Arabia) and evaluation of possible health hazard to consumers. *Pakistan J. Zool.*, **47**: 839-845.
- Mulk, S., Korai, A.L., Azizullah, A. and Khattak, M.N.K., 2016. Decreased fish diversity found near marble industry effluents in River Barandu, Pakistan. *Ecotoxicology*, **25**: 132-140. <https://doi.org/10.1007/s10646-015-1575-9>
- Mutlu, E., Aydın, S., Demir, T. and Yanık, T., 2016. Effect of zeolite and copper sulfate, administered alone and in combination on the biochemical components of blood serum of common carp, *Cyprinus carpio*. *Pakistan J. Zool.*, **48**: 1857-1863.
- Narejo, N.T., Rahmatullah, S.M. and Rashid, M.M., 2002. Studies on the reproductive biology of freshwater spiny eel, *Mastacembelus armatus* (Lacepede) reared in the cemented cisterns of BAU, Mymensingh, Bangladesh. *Pak. J. Biol. Sci.*, **5**: 809-811. <https://doi.org/10.3923/pjbs.2002.809.811>
- Ngelinkoto, P., Thevenon, F., Devarajan, N., Birane, N., Maliani, J., Buluku, A., Musibono, D., Mubedi, J.I.

- and Poté, J., 2014. Trace metal pollution in aquatic sediments and some fish species from the Kwilu-Ngongo River, Democratic Republic of Congo (Bas-Congo). *Toxicol. environ. Chem.*, **96**: 48-57. <https://doi.org/10.1080/02772248.2014.910211>
- Pal, M. and Ghosh, M., 2013. Assay of biochemical compositions of two Indian fresh water eel with special emphasis on accumulation of toxic heavy metals. *J. aquat. Fd. Prod. Technol.*, **22**: 27-35. <https://doi.org/10.1080/10498850.2011.622070>
- Palaniappan, P.R. and Karthikeyan, S., 2009. Bioaccumulation and depuration of chromium in the selected organs and whole body tissues of freshwater fish *Cirrhinus mrigala* individually and in binary solutions with nickel. *J. environ. Sci.*, **21**: 229-236. [https://doi.org/10.1016/S1001-0742\(08\)62256-1](https://doi.org/10.1016/S1001-0742(08)62256-1)
- Papagiannis, I., Kagalou, I., Leonardos, J., Petridis, D. and Kalfakakou, V., 2004. Copper and zinc in four freshwater fish species from Lake Pamvotis (Greece). *Environ. Int.*, **30**: 357-362. <https://doi.org/10.1016/j.envint.2003.08.002>
- Rosseland, B.O., Teien, H.C., Basnet, S., Borgstrøm, R. and Sharma, C.M., 2017. Trace elements and organochlorine pollutants in selected fish species from Lake Phewa, Nepal. *Toxicol. environ. Chem.*, **99**: 390-401. <https://doi.org/10.1080/02772248.2016.1189915>
- Serajuddin, M. and Mustafa, S., 1994. Feeding specializations in adult spiny eel, *Mastacembelus armatus*. *Asian Fish. Sci.*, **7**: 63-65.
- Singh, Y., Dewan, S. and Bhatt, J.P., 2016. Ascertaining spawning behavior of a Himalayan trout, *Schizothorax plagiostomus* using combinatorial approach. *Int. J. Res. Fish. Aquacul.*, **6**: 39-46.
- Siraj, M., Khisroon, M. and Khan, A., 2016. Bioaccumulation of heavy metals in different organs of *Wallago attu* from River Kabul Khyber Pakhtunkhwa, Pakistan. *Biol. Trace Elem. Res.*, **172**: 242-250. <https://doi.org/10.1007/s12011-015-0572-4>
- Siraj, M., Shaheen, M., Sthanadar, A.A., Khan, A., Chivers, D.P. and Yousafzai, A.M., 2014. A comparative study of bioaccumulation of heavy metals in two fresh water species, *Aorichthys seenghala* and *Ompok bimaculatus* at River Kabul, Khyber Pakhtunkhwa, Pakistan. *J. Biodiver. environ. Sci.*, **4**: 40-54.
- Sohail, M., Khan, M.N., Qureshi, N.A. and Chaudhry, A.S., 2017. Monitoring DNA damage in gills of freshwater mussels (*Anodonta anatina*) exposed to heavy metals. *Pakistan J. Zool.*, **49**: 305-311. <https://doi.org/10.17582/journal.pjz/2017.49.1.305.311>
- Tabinda, A.B., Hussain, M., Ahmed, I. and Yasar, A., 2010. Accumulation of toxic and essential trace metals in fish and prawns from Keti Bunder Thatta District, Sindh. *Pakistan J. Zool.*, **42**: 631-638.
- Tabinda, A.B., Zafar, S., Yasar, A. and Munir, S., 2013. Metals concentration in water, fodder, milk, meat, blood, kidney and liver of livestock and associated health impacts by intake of contaminated milk and meat. *Pakistan J. Zool.*, **45**: 1156-1160.
- Ullah, S., Hasan, Z., Ahmad, S., Rauf, M. and Khan, B., 2014. Ichthyofaunal diversity of Rhound Stream at District Lower Dir, Khyber Pakhtunkhwa, Pakistan. *Int. J. Biosci.*, **4**: 241-247.
- Ullah, S., Hussain, Z., Mahboob, S. and Al-Ghanim, K., 2016. Heavy metals in *Garra gotyla*, *Cyprinus carpio* and *Cyprinion watsoni* from the River Panjkora, District Lower Dir, Khyber Pakhtunkhwa, Pakistan. *Braz. Arch. Biol. Technol.*, **59**: e16160321. <https://doi.org/10.1590/1678-4324-2016160321>
- Vinodhini, R. and Narayanan, M., 2008. Bioaccumulation of heavy metals in organs of fresh water fish *Cyprinus carpio* (Common carp). *Int. J. environ. Sci. Technol.*, **5**: 179-182. <https://doi.org/10.1007/BF03326011>
- Yousafzai, A.M., Chivers, D.P., Khan, A.R., Ahmad, I. and Siraj, M., 2010. Comparison of heavy metals burden in two freshwater fishes *Wallago attu* and *Labeo dyocheilus* with regard to their feeding habits in natural ecosystem. *Pakistan J. Zool.*, **42**: 537-544.
- Yousafzai, A.M. and Shakoory, A.R., 2008. Heavy metal accumulation in the gills of an endangered South Asian fresh water fish as an indicator of aquatic pollution. *Pakistan J. Zool.*, **40**: 423-430.
- Yousafzai, A.M. and Shakoory, A.R., 2009. Fish white muscle as biomarker for riverine pollution. *Pakistan J. Zool.*, **41**: 179-188.