



Comparative Study on BPI Gene Expression in Various Tissues between Rongchang Pig and Landrace

Jing Xin Mao^{1,2}, Guo Wei Wang², Yuan She Huang³, Rui Wang², Guo Ze Wang⁴, Bing Zeng¹ and Fu Yuan Zuo^{1,*}

¹Department of Animal Science, Southwest University, Rongchang Campus, Chongqing, China

²College of Pharmaceutical Sciences, Southwest University, Chongqing, China

³College of Agronomy, AnShun University, Guizhou, China

⁴College of Pharmacy and Biological Engineering, Chengdu University, Chengdu, China

ABSTRACT

BPI is a pluripotent protein located in neutrophils and tissue that likely plays a vital role in host defense against GNB (gram-negative bacteria) and their endotoxin by means of its antibiotic and endotoxin neutralizing and disposing functions. BPI has several biological functions in mammals such as human, bovine, pig and rabbit, it also has major influence in the natural defense of the animal body. In this paper, by comparative study on BPI gene expression in various tissues between Rongchang pig and Landrace, we confirmed the expression of pig BPI gene in various tissues to determine the difference expression of antibacterial related gene among indigenous breed (Rongchang pig) and artificially cultivated breed (Landrace). The Rongchang pig (0 day old, 28 day old, 120 day old) and Landrace BPI (0 day old, 28 day old, 120 day old) gene were expressed in normal liver, heart, lung, kidney, spleen and small intestine. The results showed that: The expression level of BPI in lung of Rongchang pig was relatively higher level compared to that in heart, liver, kidney, spleen and small intestine. The expression level of BPI in liver of Landrace was relatively higher level compared to that in heart, lung, kidney, spleen and small intestine. Real-time fluorescence quantitative PCR analysis revealed that the native breed (Rongchang pig) has a better resistance to disease than Landrace. When Rongchang pig and Landrace were 0 day old, the expression level of BPI of Rongchang pig was higher than Landrace in heart, liver, spleen, kidney, small intestine, lower than Landrace in lung. When Rongchang pig and Landrace were 28 day old, the expression level of BPI of Rongchang pig was higher than Landrace in spleen ($p < 0.05$), kidney ($p < 0.05$), small intestine ($p < 0.05$); lower than Landrace in lung, heart, liver. When Rongchang pig and Landrace were 120 day old, the expression level of BPI of Rongchang pig was higher than Landrace in spleen, lung ($p < 0.05$), kidney ($p < 0.05$), small intestine ($p < 0.05$); lower than Landrace in heart, liver. The research trials showed that the expression of BPI gene increased with its increasing age in different growth stages. Ages and tissues are the main factors that influence the gene expression significantly.

Article Information

Received 01 December 2016

Revised 28 March 2017

Accepted 12 May 2017

Available online 18 December 2017

Authors' Contribution

FYZ conceived and designed the research. RW analyzed the data. YSH, JXM and GZW performed the experiments. JXM wrote the manuscript, FYZ revised it while BZ linguistically revised it.

Key words

Rongchang pig, Landrace, Tissue expression, BPI gene, Real-time fluorescence quantitative

INTRODUCTION

Disease has brought a great deal of breeding risk to swine production and increased the cost of epidemic prevention treatment, and the prevention and treatment of a large number of drugs, resulting in the potential threat of pork food safety (Zhong *et al.*, 2016). Utilize the genetic method, it can be a good solution to reduce the disease susceptibility and improve disease resistance (Henryon *et al.*, 2014). BPI gene has been studied in many other species, and it is expected to be a candidate gene for disease

resistance breeding. As a local pig variety, the Rongchang pig is one of the Chinese indigenous and famous breeds which produces high quality meat, it also has been characterized with high disease resistance (Liu *et al.*, 2009). The landrace is artificially cultivated breed with steady genetic feature and higher production level. We considered that it is necessary to compare those two typical indigenous and traditional swines in different genetic expressions research in order to enabling further investigation into the expression profiles and functional classifications of genes in Rongchang pig and landrace (Zou *et al.*, 2014).

Bactericidal/permeability-increasing protein (BPI) is a highly cationic protein which is located mainly in the primary granules of polymorphonuclear leucocyte (PMN). BPI has a high affinity for LPS of gram-negative bacteria

* Corresponding author: zfuyuan@163.com
0030-9923/2018/0001-0007 \$ 9.00/0
Copyright 2018 Zoological Society of Pakistan

(Elsbach and Weiss, 1993; Weiss, 2003). BPI plays an important role in the natural defense of the animal body (Schultz and Weiss, 2007; Balakrishnan *et al.*, 2013). It was reported that high expression of BPI may contribute to host immune defense against gram-negative bacterial infections in ark shell *Scapharca broughtonii* (Mao *et al.*, 2013). There are few domestic and international reports regarding porcine BPI gene polymorphisms and their associated impacts on disease resistance. In recent years, BPI was started to be identified as a candidate gene for disease-resistance breeding in pig (Christopher *et al.*, 2004). A researcher found that the porcine BPI gene may be the direct factor that caused resistance to ETEC F18 in weaning piglets (Keuzer *et al.*, 2013). Currently, although various kind of swine tissues transcriptomic data have been obtained, more studies focused on muscle to gain insights into improving meat quality and enhancing the ability of disease resistance from the perspective of agricultural economy in pig remains limited (Zhu *et al.*, 2013).

In this study, we used real-time fluorescence quantitative PCR technology (Tobias *et al.*, 2012) and took β -actin as an internal reference gene (Tesson *et al.*, 2002) to compare the differences of expression of BPI gene in tissues between Rongchang pig and Landrace. This study formed a foundation for the correlation of disease resistance and selective breeding on Rongchang pig and Landrace. Using molecular biology technology to provide an important theoretical significance and applicable value for livestock production and breeding.

Table I.- Ingredient composition of the diet.

Ingredient	% of diet	Ingredient	% of diet
Ground corn	47.60	Calcium carbonate	1.50
Soybean meal	37.00	Phosphoric acid	0.80
Whey	10.00	Vitamin premix	0.05
Poultry fat	3.00	Mineral mix	0.05
Antibiotic	1.00		

MATERIALS AND METHODS

Animal care and dietary treatments

Animal care, handling, sampling and administration procedures were approved by the Southwest University (Chongqing, China) Animal Care and Use Committee prior to initiation of the experiment. The basal diet was formulated based on NRC (1998) recommendations, the ingredient composition of the diet has shown in Table I (Zou *et al.*, 2014). All the pigs were housed and maintained according to the standard protocol and in an environmentally controlled nursery and provided ad libitum access to both feed and water.

Tissue sample preparation

All pigs (Rongchang pig and Landrace, 0 day old, 28 day old, 120 day old, 9 per age group) included in the study were healthy and raised in the same conditions with similar birth weights, weaning weights, and body sizes. Heart, liver, spleen, lung, kidney, small intestine were extracted immediately, and quick-frozen in liquid nitrogen tanks after butchering, then kept at -80 °C in the refrigerator for the extraction of total tissue RNA.

Primer design

The real-time fluorescence quantitative PCR primers used in the study were based on the published sequences of the porcine BPI and GAPDH genes. Referenced to mRNA sequence of pig's BPI gene and EST sequences, and BPI mRNA sequences of other animals provided by NCBI database (<http://www.ncbi.nlm.nih.gov/>), the specific primers were designed in highly conserved position. The primers were synthesized by Invitrogen Corporation (Carlsbad, CA, USA) and the sequence of the primers was presented in Table II.

Table II.- Sequence of the primers.

Primer	Sequence
Oligodt-Aaptor	TTTTTTTTTTTTTTTTTTT
PCR BPI	F: AGTCTGCGGCTGGGTTAT R: GGAGGACACGGTATTGGTC
PCR β -actin	F: TCTGGCACCACACCTTCT R: TGATCTGGGTCATCTTCTCAC

Total RNA extraction and detection

Total RNA was extracted from the tissues of the Rongchang pig and Landrace using a Trizol reagent (Invitrogen, Carlsbad, CA, USA), according to the manufacturer's instructions. In order to detect the integrity of RNA extracted, the RNA samples were electrophoresis at 260 V in 1% agarose gel for 7-8 min. Then 1 μ L of the total RNA was mixed with 99 μ L of DEPC water, the concentrations and OD values at 260 nm and 280nm were determined in the protein nucleic acid analyzer. An OD260/OD280 value of 1.8-2.0 indicated that the total RNA achieved the quality for the reverse transcription experiment (Cury and Koo, 2008).

Reverse transcription

Single-stranded cDNA was generated using the primescript RT-PCR Kit (TaKaRa) following the manufacturer's instructions. The 10 μ L of reagents containing 1 μ L of RT buffer (10 \times), 3.75 μ L of RNase free dH₂O, 0.5 μ L of AMV reverse transcriptase, 0.25

μL of RNase inhibitor, $0.5 \mu\text{L}$ of oligo dT, $1 \mu\text{L}$ of dNTP mixture (10 mM) and $1 \mu\text{L}$ of positive control RNA were added successively into 0.2 mL of PCR tube without RNA enzyme for reverse transcription reaction. The PCR reverse transcription process was conducted as following: $30 \text{ }^\circ\text{C}$ for 10 min , $42 \text{ }^\circ\text{C}$ for 30 min , $99 \text{ }^\circ\text{C}$ for 5 min , $5 \text{ }^\circ\text{C}$ for 5 min , then preserve at $-20 \text{ }^\circ\text{C}$. Then, the PCR amplification program was conducted according to the reference (Zhu *et al.*, 2013).

Real-time fluorescence quantitative PCR

The $25 \mu\text{L}$ reaction system consisted of $2 \mu\text{L}$ of cDNA ($10 \mu\text{M/L}$), $1 \mu\text{L}$ of BPI-F, $1 \mu\text{L}$ of BPI-R, $12.5 \mu\text{L}$ of SYBR Green Real-time PCR Master Mix ($2\times$), and $8.5 \mu\text{L}$ of dH_2O . The PCR reaction conditions included 40 cycles at $95 \text{ }^\circ\text{C}$ for 330 s , $56 \text{ }^\circ\text{C}$ for 30 s and $72 \text{ }^\circ\text{C}$ for 30 s . Each sample was tested using real-time PCR three times, and the average of three measurements was used. Real-time fluorescence quantitative PCR was performed using a Bio-Rad IQ5 sequence-detection system (Bio-Rad, CA, USA) with SYBR Green PCR Master Mix (TaKaRa), according to the manufacturer's instruction. The BPI fragment was amplified using the primers listed in Table II.

Statistical analysis

Triplicate PCR amplifications were performed for each sample. Samples without cDNA template were used as negative controls and pig's β -actin gene download from NCBI database (<http://www.ncbi.nlm.nih.gov/>) was used as an internal reference. The real-time fluorescence quantitative PCR results were processed using the $2^{-\Delta\Delta\text{Ct}}$ method (ΔCt =the mean expression level of BPI-the mean expression level of β -actin). The expression levels of this gene in other groups could be quantified since the group with the least ΔCt was defined as 1 ($\Delta\Delta\text{Ct}$ = ΔCt of the other group-group with the least ΔCt in each experiments). Based on the mathematical derivation of the Livak sehmittgen, when the amplification efficiency of target gene and reference gene are consistent, the expression of the target gene in the experimental group was compared with that of the control group. The standard curve for BPI was $Y = -3.353X - 1.839$ with a correlation coefficient of 0.998 , and the amplification efficiency was 92.3% . The standard curve for the β -actin gene was $Y = -3.341X - 1.827$ with a correlation coefficient of 0.997 , and the amplification efficiency was 91.2% (Livak and Sehmittgen, 2001). The consistent amplification efficiencies indicated that the $2^{-\Delta\Delta\text{Ct}}$ method can be used for quantitative calculations. Analysis of data were performed by ANOVA for a completely randomized design using procedure of SPSS 18.0 (SPSS Inc., Chicago, IL, USA). The LSD was used to analyze the significance of differences in BPI expression.

RESULTS

Real-time fluorescence quantitative PCR amplification results

The standard curve chart (Fig. 1) shows that the dilution gradient can be in a straight line, the correlation coefficient of the standard curve is between 0.99 and 1 , indicate that less error and higher reliability. Melt curve analysis of BPI and β -actin PCR product was shown in Figure 2. It can be seen that only single peak curve was presented, indicating that the amplified products was unitary and no primer-dimers and other non-specific amplification was produced.




Fig. 1. Real-time fluorescence quantitative PCR standard curve chart of BPI gene and β -actin.




Fig. 2. Melt curve analysis of BPI and β -actin PCR product.

Expression of BPI gene in different tissues of Rongchang pig and Landrace

Figure 3 was the relationship between expression and day old in tissues of Rongchang pig. From the Figure

3, it revealed that the BPI gene increased with the age of the pig and the increase trend of BPI gene in lung was more significant than that in other tissues. In addition, the expression of BPI gene in spleen was higher than that in other tissues in the embryonic (0 day old), the initial growth (28 day old) and growth (120 day old) stages. Figure 4 is the relationship between expression and days in tissues of Landrace. It revealed that the expression of BPI gene increased with the age of the pig and the increase trend of BPI gene in liver was more significant than that in other tissues. Figure 5 is the BPI expression of varies tissues between Rongchang pig and Landrace at different ages. It revealed that the expression of BPI gene increased with its increasing age in different growth stages. Figure 5A shows that the expression levels of BPI gene when Rongchang pig and Landrace were 0 day old; Figure 5B shows that the




Fig. 3. The relationship between BPI expression and days in tissues of Rongchang pig.




Fig. 4. The relationship between BPI expression and days in tissues of Landrace.




Fig. 5. The BPI expression of varies tissues between Rongchang pig and Landrace at different ages. A, 0 day old Rongchang pig and Landrace of varies tissues; B, 28 day old Rongchang pig and Landrace of varies tissues; C, 120 day old Rongchang pig and Landrace of varies tissues.

expression levels of BPI gene when Rongchang pig and Landrace were 28 day old; Figure 5C shows that the expression levels of BPI gene when Rongchang pig and Landrace were 120 day old.

DISCUSSION

In our study, cross-intron primer design method is used to eliminate the influence of DNA. The intron length of BPI and β -actin crossed are more than 500 bp and the genomic DNA cannot amplify at such a length of intron (Liang and Pardee, 1992). The specificity of RT-PCR depends on the specificity of the banding of primers and template DNA. Sequence alignment and structure function prediction indicate that the structure and function BPI gene of pig is similar as that of human and cow (Eckert *et al.*, 2006). The design of the primers refers the sequence of the cDNA of pig and BPI exon of human.

Given its ability to neutralize endotoxin and protect against gram-negative bacteria (Zhou *et al.*, 1999), the BPI has application prospects widely, it may be considered as a “super antibiotic”. Porcine BPI also has these functions (Tuggle *et al.*, 2006). In our study, we have confirmed the findings from studies done in Rongchang and Landrace piglets, that BPI gene expression levels are higher in spleen than in other tissues. We also have reconfirmed the findings from studies done in Rongchang and Landrace adult swines, that BPI gene expression levels are higher in lung and liver than in other tissues (Gan *et al.*, 2015). The increase of BPI gene in lung is more significant than that in other tissues with the increase of the age of Rongchang pig. The reason could be that the experiments are conducted in July and August, which is the hottest period in Chongqing, China. Thus the high incidence of respiratory disease leads to such a trend. On the other side, the BPI has the function of protection of lung. The researchers found that recombinant BPI administration protects pigs against endotoxemia-induced acute lung injury (Vandermeer *et al.*, 1994). While the reason for the increase trend of BPI gene in liver of Landrace is more significant could be attributed to the weak constitution and poor resistance of the Landrace. This is similar to the study of researcher (Gan *et al.*, 2015) in Rongchang pig. Since the Landrace is a species from Denmark and usually lives in relatively cold place, they are easy to have an attack of illness and then many drugs are fed. Therefore, the BPI gene increases more since liver is a detoxification and excretion organ (Gaur *et al.*, 2014). BPI is vital in the innate immune response, our findings may indicate that the high expression of BPI gene may be related to age in the lung and liver. In our trials, we found that the expression level of BPI gene in spleen of Rongchang pig was higher than other tissues (heart, liver, lung, kidney, small intestine). The spleen is one of the vital immune organs in body and is mainly responsible for making antibodies, regulating immune responses, filtering aging erythrocytes, storing blood. The normal structure and function of immune organs are connected with swine

immunity (Wu *et al.*, 2016). It has a high BPI expression level in spleen indicate that the Rongchang pig has a high resistance to disease.

We tested differential performances of BPI genes in 6 tissues, further molecular level exploration is required. In the embryonic stage of pigs, the BPI gene in lung of Rongchang pig is lower than Landrace, while in other tissues are higher than those of Landrace. In the initial growth stage, the BPI gene in spleen, kidney and small intestine of Rongchang pig is higher than that of Landrace, while others are lower. In the growth stage, the BPI gene in spleen, lung, kidney and small intestine of Rongchang pig is higher than that of Landrace, while others are lower. The studies show that the BPI gene also shows similar tissue-specific expression in a different breed of weaned piglets and adult swines. BPI can directly kill gram-negative bacteria such as E.coli F18, and that up-regulated expression is closely related to resistance to intestinal E.coli F18 (Keuzer *et al.*, 2013), also the BPI gene tissue specificity has vital implications in biological and engineering measures to utilize the activity of porcine endogenous BPI protein. Many researches indicated that BPI has the effect of neutralize porcine endogenous endotoxin and gram-negative bacteria (Bingle and Jeremy, 2004). We can see a difference in expression of antibody specific genes in two different pig breeds clearly, which indicate towards the underlying differences in genetic make-up of resistant and susceptible animals. The findings of this study provided a new insight for present research, the use of BPI as a disease resistance candidate gene for breeding still needs further study in mammals.

CONCLUSION

Real-time fluorescence quantitative PCR analysis revealed that Rongchang pig has a better resistance to disease and immunity than Landrace. The research trials showed that the expression of BPI gene increased with its increasing age in different growth stages, ages and tissues are the main factors that influence the gene expression significantly. BPI as a disease resistance candidate gene may play an important role for breeding, the resistance was related to the upregulation of BPI gene expression in the spleen, lung and liver.

ACKNOWLEDGMENTS

This work was supported by the Chinese National Science and Technology Ministry of state science and technology supporting plan. Collecting samples were supported by Chongqing Academy of Animal Sciences.

Statement of conflict of interest

Authors have declared no conflict of interest.

REFERENCES

- Balakrishnan, A., Marathe, S.A. and Joglekar, M., 2013. Bactericidal/permeability increasing protein: A multifaceted protein with functions beyond LPS neutralization. *J. Innate. Immun.* **19**: 339-347. <https://doi.org/10.1177/1753425912465098>
- Bingle, C.D. and Jeremy, C.C., 2004. Meet the relatives: A family of BPI- and LBP-related proteins. *J. Trends Immunol.*, **25**: 53-55. <https://doi.org/10.1016/j.it.2003.11.007>
- Christopher, K.T., Thomas, J.S. and Shi, X.W., 2004. Genetic markers for improved disease resistance in animals (BPI). *U.S. Patent.*, 20040234980.
- Cury, J.A. and Koo, H., 2008. Extraction and purification of total RNA from *Streptococcus mutans* biofilms. *J. Braz. Oral. Res.*, **22**: 216-222. <https://doi.org/10.1590/S1806-83242008000300005>
- Eckert, M., Wittmann, I. and Rollinghoff, M., 2006. Endotoxin induced expression of murine bactericidal permeability/increasing protein is mediated exclusively by toll/IL-1 receptor domain-containing adaptor inducing IFN-beta-dependent pathways. *J. Immunol.*, **176**: 522. <https://doi.org/10.4049/jimmunol.176.1.522>
- Elsbach, P. and Weiss, J., 1993. The bactericidal/permeability-increasing protein, a potent element in host-defense against gram-negative bacteria and lipopolysaccharide. *J. Immunobiol.*, **187**: 417-429. [https://doi.org/10.1016/S0171-2985\(11\)80354-2](https://doi.org/10.1016/S0171-2985(11)80354-2)
- Gan, L., Xie, L.W., Xiang, Z.H. and He, N.J., 2015. Transcriptomic analysis of Rongchang pig brains and livers. *J. Genet.*, **560**: 96-106. <https://doi.org/10.1016/j.gene.2015.01.051>
- Gaur, U., Wu, H.Y., Feng, Z., Yang, X.H., Guo, R., Wu, J.J., Li, K., Jin, M.L., Mei, S.Q. and Liu, G.S., 2014. Gene expressions in major organs between Enshi Black and Landrace pigs after *Streptococcus suis* infection. *Pak. Vet. J.*, **34**: 432-437.
- Henryon, M., Berg, P. and Sorensen, A.C., 2014. Animal-breeding schemes using genomic information need breeding plans designed to maximize long-term genetic gains. *Livest. Sci.*, **166**: 38-47. <https://doi.org/10.1016/j.livsci.2014.06.016>
- Keuzer, S., Reissmann, M. and Brockmann, G.A., 2013. New fast and cost-effective gene test to get the ETEC F18 receptor status in pigs. *J. Vet. Microbiol.*, **163**: 392-394. <https://doi.org/10.1016/j.vetmic.2012.12.040>
- Liang, P. and Pardee, A.C., 1992. Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science*, **257**: 967-971. <https://doi.org/10.1126/science.1354393>
- Liu, L., Yin, J.D., Li, W., Liu, K., Peng, Y., Tan, P.P. and Ma, R.L.Z., 2009. Construction of a bacterial artificial chromosome library for the Rongchang pig breed and its use for the identification of genes involved in intramuscular fat deposition. *Biochem. biophys. Res. Commun.*, **391**: 1280-1284. <https://doi.org/10.1016/j.bbrc.2009.12.060>
- Livak, K.J. and Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2^{-ΔΔCt} method. *Methods*, **25**: 402-408. <https://doi.org/10.1006/meth.2001.1262>
- Mao, Y.Z., Zhou, C.Y., Zhu, L., Huang, Y., Yan, T.R., Fang J.G. and Zhu. W., 2013. Identification and expression analysis on bactericidal permeability-increasing protein (BPI)/lipopolysaccharide-binding protein (LBP) of ark shell, *Scapharca broughtonii*. *J. Fish Shellfish Immunol.*, **35**: 642-652. <https://doi.org/10.1016/j.fsi.2013.05.025>
- NRC, 1998. *Nutrient requirements of swine*, 10th ed. National Academy Press, Washington, DC.
- Schultz, H. and Weiss. J.P., 2007. The bactericidal/permeability-increasing protein (BPI) in infection and inflammatory disease. *J. clin. Chim. Acta*, **384**: 12-23. <https://doi.org/10.1016/j.cca.2007.07.005>
- Tuggle, C.K., Stabel, T.J., Shi, X.W. and Mellencamp, M.A. 2006. *Genetic markers for improved disease resistance in animals (BPI)*. Iowa State University Patents, US 10/161, 968.
- Tesson, L., Heslan, J.M., Menoret, S. and Anegon, I., 2002. Rapid and accurate determination of zygosity in transgenic animals by real-time quantitative PCR. *Transgen. Res.*, **11**: 43-48. <https://doi.org/10.1023/A:1013928600442>
- Tobias, T.J., Bouma A., Klinkenberg, D.A., Daemen, J.J.M., Stegeman, J.A., Wagenaar, J.A. and Duimb, B., 2012. Detection of *Actinobacillus pleuropneumoniae* in pigs by real-time quantitative PCR for the *apxIVA* gene. *Vet. J.*, **193**: 557-560. <https://doi.org/10.1016/j.tvjl.2012.02.004>
- Vandermeer, T.J., Menconi, M.J., O'Sullivan, B.P., Larkin, V.A. and Wang, H., 1994. Bactericidal/permeability-increasing protein ameliorates acute lung injury in porcine endotoxemia. *J. appl. Physiol.*, **76**: 2006-2014.
- Weiss, J., 2003. Bactericidal/permeability-increasing protein (BPI) and lipopolysaccharide-binding protein (LBP): structure, function and regulation in host defence against Gram-negative bacteria.

- Biochem. Soc. Trans.*, **31**: 785-790. <https://doi.org/10.1042/bst0310785>
- Wu, X.J., Cao, W., Jia, G., Zhao, H., Chen, X.L., Wu, C.M., Tang, J.Y., Wang, J. and Liu, G.M., 2016. New insights into the role of spermine in enhancing the antioxidant capacity of rat spleen and liver under oxidative stress. *Anim. Nutr.*, **3**: 85-90. <https://doi.org/10.1016/j.aninu.2016.11.005>
- Zhu, J., Zi, C. and Wu, Z.C., 2013. Age-dependent expression of the BPI gene in Sutai piglets. *Genet. mol. Res.*, **12**: 2120-2126. <https://doi.org/10.4238/2012.April.12.1>
- Zou, T.D., Mao, X.B., Yu, B., He, J., Zheng, P., Yu, J. and Chen, D.W., 2014. Effects of dietary energy density and apparent ileal digestible lysine: digestible energy ratio on growth performance, meat quality, and peroxisome proliferator-activated receptor γ (PPAR γ) gene expression of muscle and adipose tissues in Landrace \times Rongchang crossbred pigs. *J. Livest. Sci.*, **167**: 219-226. <https://doi.org/10.1016/j.livsci.2014.04.032>
- Zhong, Y.Q., Huang, Z.H. and Wu, L.H., 2016. Identifying critical factors influencing the safety and quality related behaviors of pig farmers in China. *Fd. Contr.*, **71**: 264-272.
- Zhou, H., Yuan, J. and Zhou, L., 1999. The role of bactericidal / permeability-increasing protein of endotoxin neutralization *in vitro* and *in vivo*. *Nat. med. J. China*, **79**: 304-305.