



## Short Communication

# Evaluation of Lactation and Performance of Friesian Cow in Local Environment of Quetta, Balochistan, Pakistan

Madiha Khalil<sup>1</sup>, Amjad Farooq<sup>1</sup>, Asim Faraz<sup>2\*</sup>, Abdul Waheed<sup>2</sup>, Nasir Ali Tauqir<sup>3</sup>, Riaz Hussain Mirza<sup>2</sup> and Hafiz Muhammad Ishaq<sup>2</sup>

<sup>1</sup>Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan

<sup>2</sup>Department of Livestock and Poultry Production, Bahauddin Zakariya University, Multan, Pakistan

<sup>3</sup>Department of Animal Science, University of Sargodha, Sargodha, Pakistan

## ABSTRACT

The present study was conducted in order to analyze lactation performance of Friesian cattle in the local environment of Quetta, Pakistan and to evaluate non-genetic factors affecting the lactation traits. For this purpose, data on lactation performance of Holstein Friesian cows were collected from government livestock farms located at Quetta. Data obtained on calving from 608 cows during last sixteen years (2000-2016) were analyzed. After editing the data for consistency and errors it was subjected to statistical analysis using fixed effect models. Independent variables were year of birth, season of birth, year of calving, season of calving and parity. Age of animal (linear and quadratic effects) and lactation period were fitted as co-variables. Dependent variables were lactation performance traits (lactation milk yield, dry period, etc.). The results of all traits showed that the year of calving had a highly significant ( $p < 0.01$ ) effect on lactation length, month of calving and parity. Lactation milk yield was significantly affected by year of calving ( $p < 0.01$ ) and month of calving ( $p < 0.05$ ) while parity had a non-significant ( $p > 0.05$ ) effect. Milk yield over 305 days was significantly ( $p < 0.05$ ) affected by year and month of calving, while parity had a non-significant ( $p > 0.05$ ) effect. Similarly, year of calving and month of calving had a highly significant ( $p < 0.01$ ) effect on dry period which is non-significantly ( $p > 0.05$ ) affected by parity.

### Article Information

Received 28 March 2019

Revised 28 April 2020

Accepted 28 October 2020

Available online 07 December 2021  
(early access)

### Authors' Contribution

MK conducted research. AF supervised the work. AW analyzed the data. AF wrote the paper. NAT, RHM and HMI helped in write up.

### Key words

Lactation performance, Non-genetic factors, Holstein friesian, Quetta, Dairy production

Geographically Balochistan is the largest province of Pakistan; having social and cultural variations. Balochistan contains 44% of the national sheep population and smaller number of cows and buffalo due to which dairy production is low in this province (GOP, 2008-09). In 1977-78, the Livestock and Dairy Development Department Government of Balochistan decided to import the non-native cattle breeds to overcome the shortage of milk. For this purpose, Holstein Friesian cattle was imported from Denmark. About 175 pregnant heifers and 2 bulls were imported and stationed at Government Dairy Farm Quetta in order to breed non-native dairy cattle under different climatic conditions.

In spite of the import of non-native cattle breeds over thirty years, it is estimated that the desired production is not achieved up till now (Afzal and Naqvi, 2004). The main objective of this study is to evaluate the potentials, problems and solutions concerning the dairy industry in Pakistan

and to identify potential options to improve the lactation performance of Holstein Friesian cow kept under different conditions in different areas of country.

### Materials and methods

Several traits of interest were recorded from Holstein Friesian cattle ( $n=610$  out of which 2 died) reared at the Government Dairy Farm Pishin, Balochistan, Pakistan. These included various measures of lactation performance and general information on the cows: traits like lactation milk yield, lactation length, dry period, year of calving, month of calving and parity. Standard management conditions were offered (24-h *ad lib* water, 2 times green fodder) with concentrate (Anmol Vanda @ 3 kg per head). The animals were housed in semi-open system and vaccination against various diseases were done according to vaccination schedule given by Govt of Balochistan Livestock Department. Avg age at calving ranged from 24-30 months and 6 parities were recorded on average.

The data was fed into MS excel worksheet and edited. Any record beyond  $\pm 3$  S.D. (standard deviation) was omitted/removed. After editing, the data was analyzed

\* Corresponding author: drasimfaraz@bzu.edu.pk  
0030-9923/2021/0001-0001 \$ 9.00/0  
Copyright 2021 Zoological Society of Pakistan

using following mathematical models:

$$Y_{(ijklm)} = YOC_{(i)} + SOC_{(j)} + LL_{(k)} + Age_{(i)} + Age^2 + YOC \times SOC_{(ij)} + \varepsilon_{(ijklm)}$$

where  $Y_{ijklm}$  is lactation performance recorded,  $YOC_{(i)}$  is  $i^{\text{th}}$  year of calving,  $SOC_{(j)}$  is  $j^{\text{th}}$  season of calving,  $YOC \times SOC_{(ij)}$  is interaction of  $i^{\text{th}}$  and  $j^{\text{th}}$  year of calving,  $LL_{(k)}$  is  $k^{\text{th}}$  length of lactation related to any milk or lactation performance recorded,  $Age_{(i)}$  is age of the animal at time of calving recorded in months,  $Age^2$  is quadratic effects of age of animal (as both linear and quadratic effects of age are usually used in such models mostly as covariables),  $\varepsilon_{ijklm}$  is random error associated with each observation.

Preliminary descriptive analysis was conducted using SPSS software. Further analysis was conducted using ASREML software.

Different models were applied and finally one model was reached for optimum results for all traits. LL was both treated as dependent variable in milk traits and independent variable as well. Parity remained non-significant in all models so it was excluded ultimately.

**Table I.- Average mean values of LMY, LL, DP and MY-305 days of Holstein Friesian cattle**

Traits	Mean Values
LMY (liters)	2322.53±150.7
LL (days)	278.18±10.86
DP (days)	234.46±52.16
MY-305 days (liters)	2781.08±551.26

LL, lactation length; LMY, lactation milk yield; MY-305 days, milk yield 305 days; DP, dry period

**Table II. Analysis of variance of LL, LMY, MY-305 days and DP of Holstein Friesian cattle affected by YOC, MOC and Parity.**

Traits	YOC	MOC	Parity
LL	***	NS	NS
LMY	***	**	NS
MY-305 days	**	NS	**
DP	***	NS	NS

YOC has highly significant effect on LL, LMY and DP ( $p < 0.01$ ) while significant effect ( $p < 0.05$ ) on MY-305 days.

MOC non-significantly ( $p > 0.05$ ) effected on LL, MY-305 days and DP while this factor has significant effect ( $p < 0.05$ ) on LMY.

Parity has significant effect ( $p < 0.05$ ) on MY-305 days while all other traits were non-significantly affected ( $p > 0.05$ ) by this factor.

MOC, month of calving and YOC, year of calving. See Table I for remaining abbreviations.

### Results and discussion

The average lactation milk yield (LMY) of

2322.53±150.7 L as found in present study was greater than the values reported by different findings of many researchers. Nahar *et al.* (1992) reported LMY of Desi × HF as 1702.8±44.2 L, value of average LMY by Talbott (1994) was 1314±23.7 L in Desi cross. Mean lactation milk yield was found to be 1395±9.9 L in cross animals (Dahlin *et al.*, 1998). Average LMY was found to be 1432.8±51.2 L (Aslam *et al.*, 2002) and 429±11.4 L (Rehman *et al.*, 2008). LMY in the study of Habib *et al.* (2010) was found to be 500.7±19.3 L. LMY value reported by Hassan and Khan (2013) was 1633±49.03 L while in the findings of Fernando *et al.* (2016) stated LMY was 2235±77.6 L. All these given values of different research work were lower than present results. Similarly reported values of LMY of Afridi (1999), Sattar *et al.* (2005), Irshad *et al.* (2011), Sandhu *et al.* (2011), Usman *et al.* (2012) and Faid-Allah (2015) were 3002.66±45.49, 2772.76±65.0, 3992.41±16.20, 3977.75±37.20, 3438±887.19 and 7208.72±1753.6 L respectively, which are higher than the current findings.

The average lactation length (LL) in Holstein-Friesian cattle ( $n=608$ ) was found to be 278.18±10.86 days which is higher than those reported by other authors. Talbott (1994), Dahlin *et al.* (1998), Khatri *et al.* (2004) and Rehman *et al.* (2008) stated average lactation length as 238±2.6, 232±1.3, 226.98 and 235±2.2 days respectively in different cattle breeds and these values are lower than present results. Islam and Bhuiyan (1997) reported LL value of 247.23±3.51 days, Habib *et al.* (2010) as 259.6±6.2 days and Hassan and Khan (2013) as 240±5.5 days. All of these values are lower than current findings.

Some LL values were higher compared to present study as reported by Nahar *et al.* (1992) as 330.5±3.6 days, Afridi (1999) as 315.09±17.75 days, Sattar *et al.* (2005) as 291.86±6.55 days, Irshad *et al.* (2011) as 320.14±11.14 days, Sandhu *et al.* (2011) as 314.19±0.91 days, Usman *et al.* (2012) as 366.5±76.71 days, Fernando *et al.* (2016) as 323.78±91.02 days. Javed *et al.* (2004) reported this value as 278.40±90.17 days which was similar to the current results. The variations reported in the results from different researchers may be attributed to the variations in breed, housing, management and nutritional conditions.

Overall average dry period (DP) in present study was found to be 234.46±52.16 days that was considerably longer than reported by Nahar *et al.* (1992) as 148.1±8.8 days, Talbott (1994) as 228±11.7 days, Aslam *et al.* (2002) as 198.9±17.17 days, Sattar *et al.* (2005) as 224.99±10.00 days, Habib *et al.* (2010) as 162.4±7.7 days, Suhail *et al.* (2010) as 169.26±16.45 days, Sandhu *et al.* (2011) as 87.06±1.63 days, Usman *et al.* (2012) as 100.26±61.38 days, Faid-Allah (2015) as 72.33±20.03 days and Fernando *et al.* (2016) as 120±12.2 days. DP reported by

Khatri et al. (2004) and Rehman et al. (2008) was 455.1 and 244±2.8 days respectively, which were higher than the current findings.

Regarding average 305 days milk yield (days MY), results were 2781.08±551.26 L which are higher than reported values of different researchers. Average values of MY-305 days reported by Mohiuddin et al. (1991), Chaudary et al. (1994), Dahlin et al. (1998), Aslam et al. (2002), Rehman et al. (2008), Hassan and Khan (2013) were found to be 1603±12.3, 2729.0±669.3, 1363±30.1, 1864.5±77.1, 1393±11.7 and 1613±49.03 L. The results of Nawaz et al. (1993) and Faid-Allah (2015) were 2889±92 and 6384.95±1236.9 L, which were considerably higher than current values.

Non-genetic factors like year of calving (YOC), month of calving (MOC) and parity also have different effects on the lactation traits. LMY was significantly ( $p < 0.01$ ) affected by year of calving and month of calving while parity influenced non-significantly ( $p > 0.05$ ) on this lactation trait. Similarly, year of calving and month of calving also significantly ( $p < 0.01$ ) affected the lactation length. In this case parity showed non-significant ( $p > 0.05$ ) effect as in previous trait. Milk yield 305 days and dry period was significantly ( $p < 0.01$ ) affected by year of calving and month of calving. Parity has non-significant ( $p > 0.05$ ) effect on both traits.

The relationship between fixed factors and lactation performance could be used to increase production. For example, if year of calving had a positive effect on milk yield, we can use this relationship in practice to increase milk yield by adopting the management practices to inseminate the animal regarding our desired calving year. Month of calving and year of calving has significant effect on milk yield, hence the husbandry practices could be adopted for fruitful gains.

Supplementary Table 1 gives study conditions, parameters investigated and findings from literature data by different researchers to have a deep look on productive performance of crossbred and exotic cattle.

### Conclusion

Non-native cattle breeds imported and kept at Balochistan at different locations are performing well regarding production. The climate in Quetta is cool, humid and temperate so these climatic conditions are favoring the production potential exploitation. Better performance of dairy cattle could be achieved by modern husbandry practices, nutritional manipulations and proper disease control. By adopting these manipulations, we can attain the desired effects of YOC, MOC, SOC and age of the cow to increase milk production.

### Acknowledgements

Authors are grateful to the Government Livestock Farms in extending their cooperation for this research work.

### Supplementary material

There is supplementary material associated with this article. Access the material online at: <https://dx.doi.org/10.17582/journal.pjz/20190328080332>

### Statement of conflict of interest


The authors have declared no conflict of interest.

### References

- Afridi, R.J., 1999. *Pak. Vet. J.*, **19**: 192. <https://doi.org/10.1088/0952-4746/19/2/608>
- Afzal, M. and Naqvi, A.N., 2004. *Q. Sci. Vis.*, **9**: 1-14.
- Aslam, M., Nawaz, M. and Khan, M.S., 2002. *Int. J. Agric. Biol.*, **1560**: 565-567.
- Chaudary, M.Z., Tahir, M.J. and Rafique, M., 1994. *Asian Austral. J. Anim. Sci.*, **7**: 385-387. <https://doi.org/10.5713/ajas.1994.383>
- Dabdoub, S.A.M., 2009. *Al-Anbar. J. Vet. Sci.*, **2**: 48.
- Dahlin, A.K., Zafar, U.N., Saleem, A.H., Chaudary, M. and Philipsson, M.A.J., 1998. *Anim. Sci.*, **66**: 307-318. <https://doi.org/10.1017/S1357729800009437>
- Faid-Allah, E., 2015. *Indonesian J. Anim. Vet. Sci.*, **20**: 10-17. <https://doi.org/10.14334/jitv.v20i3.1182>
- Fernando, P.R.P., Sinniah, J. and Thatchaneshkanth, S., 2016. *Glob. Vet.*, **17**: 392-400.
- GOP, 2008-09. Economic Advisor's Wing. Ministry of Finance, Government of Pakistan Islamabad, Pakistan.
- Habib, M.A., Afroz, M.A. and Bhuniyan, A.K.F.H., 2010. *Bangladesh Vet. J.*, **27**: 18-25. <https://doi.org/10.3329/bvet.v27i1.5911>
- Hassan, F. and Khan, M.S., 2013. *J. Anim. Pl. Sci.*, **23**: 705-714.
- Irshad, A., Tariq, M.M., Bajwa, M.A., Abbas, F., Isani, G.B., Soomro, G.H., Waheed, A. and Khan, K., 2011. *Igdir Univ. J. Inst. Sci. Technol.*, **1**: 53-57.
- Islam, S.S. and Bhuiyan, A.K.F.H., 1997. *Asian Austral. J. Anim. Sci.*, **10**: 581-586. <https://doi.org/10.5713/ajas.1997.581>
- Javed, K., Afzal, M., Sattar, A. and Mirza, R.H., 2004. *Pak. Vet. J.*, **24**: 58.
- Khatri, P., Bakhsh, K., Mirbahar and Samo, U., 2004. *J. Anim. Vet. Adv.*, **3**: 353-355.
- Mohiuddin, G., Yaqoob, M., Nawaz, H. and Akhtar, P., 1991. *Pak. Vet. J.*, **14**: 238-241.
- Nahar, T.N., Aslam, M. and Hasanath, M.A., 1992. *Asian Austral. J. Anim. Sci.*, **5**: 435-438. <https://doi.org/10.17582/journal.pjz/20190328080332>

- [org/10.5713/ajas.1992.435](https://doi.org/10.5713/ajas.1992.435)
- Nawaz, M., Anwar, M., Mirza, M.A. and Nawaz, R., 1993. *Asian Austral. J. Anim. Sci.*, **6**: 549-553. <https://doi.org/10.5713/ajas.1993.549>
- Rehman, Z.U., Khan, M.S., Bhatti, S.A., Iqbal, J. and Iqbal, A., 2008. *Arch. Tierz., Dummerstorf*, **51**: 305-317. <https://doi.org/10.5194/aab-51-305-2008>
- Sandhu, Z.S., Tariq, M.M., Baloch, M.H. and Qaimkhani, M.A., 2011. *Pak. J. Life Soc. Sci.*, **9**: 128-133.
- Sattar, A., Mirza, R.H., Niazi, A.A.K. and Latif, M., 2005. *Pak. Vet. J.*, **25**: 75.
- Suhail, S.M., Ahmed, I., Hafeez, A., Ahmed, S., Jan, D., Khan, S. and Rehman, A., 2010. *Sarhad J. Agric.*, **26**: 87-91.
- Talbott, C.W., 1994. *Potential to increase milk yield efficiency in tropical countries*. Ph.D. thesis, North California State Univ., Raleigh, USA.
- Usman, T., Guo, G., Suhail, S.M., Ahmad, S., Qiaoxiang, L., Qureshi, M.S. and Wang, Y., 2012. *J. Pl. Sci.*, **22**: 92-95.

Online First Article



## Supplementary Material

# Evaluation of Lactation and Performance of Friesian Cow in Local Environment of Quetta, Balochistan, Pakistan

Madiha Khalil<sup>1</sup>, Amjad Farooq<sup>1</sup>, Asim Faraz<sup>2\*</sup>, Abdul Waheed<sup>2</sup>, Nasir Ali Tauqir<sup>3</sup>, Riaz Hussain Mirza<sup>2</sup> and Hafiz Muhammad Ishaq<sup>2</sup>

<sup>1</sup>Institute of Pure and Applied Biology, Bahauddin Zakariya University, Multan, Pakistan

<sup>2</sup>Department of Livestock and Poultry Production, Bahauddin Zakariya University, Multan, Pakistan

<sup>3</sup>Department of Animal Science, University of Sargodha, Sargodha, Pakistan

### Supplementary Table I. Meta analysis of productive and reproductive traits from literature data.

Research	Parameters	Results
Afridi, 1999	To analyze productive performance of Holstein-Friesian cattle in NWFP	The overall performance of imported Holstein-Friesian was better than local born Holstein-Friesian. The milk yield and lactation length were maximum in autumn.
Javed <i>et al.</i> , 2004	To study environmental factors affecting milk yield in Friesian cows in Punjab	The milk yield ↑ with ↑ in lactation length, yet it did not seem advantageous to have lactations exceeding one year. The daily milk yield in the later stages of lactation ↓ and hence affects the lifetime production. Moreover, longer lactations prolong the calving interval, thereby decreasing the number of calves that could be obtained during the life span of a cow.
Sattar <i>et al.</i> , 2005	To investigate productive and reproductive performance of Holstein-Friesian cows in Pakistan	The reproductive and productive performance of Holstein-Friesian herd at Bhunikey is declining gradually, indicating that the managerial practices at the farm need to be improved for better reproductive and productive performance of the herd.
Dabdoub, 2009	To check milk production and reproductive efficiency in Friesian and Friesian×Sharabi cows	For improving efficiency in Friesian and crossbreed cattle, it is necessary that functional traits interms of reproductive efficiency and age at first calving deserve primary consideration beside milk production into national breeding programs.
Irshad <i>et al.</i> , 2011	Performance analysis of Holstein-Friesian cattle herd under semi-intensive management at Pishin dairy farm Balochistan	The productive and reproductive traits of Holstein-Friesian cattle were low to moderate. Therefore, it is necessary to improve managerial practices at the farm for better reproductive and productive traits.
Sandhu <i>et al.</i> , 2011	Performance analysis of Holstein-Friesian cattle in intensive management at dairy farm Quetta, Balochistan, Pakistan	Productive and reproductive performance of Holstein-Friesian cattle were low to modest so it is required to improve managerial practices at the farm for better productive and reproductive performance.
Usman <i>et al.</i> , 2012	Performance traits study of Holstein Friesian cattle under subtropical conditions	Significant effect of calving season on the studied traits and low to moderate heritability and repeatability estimates depicted that these traits were highly influenced by non-genetic factors and efforts to improve subtropical environmental conditions will bring an improvement in these traits.

Continue on next page.....

\* Corresponding author: [drasimfaraz@bzu.edu.pk](mailto:drasimfaraz@bzu.edu.pk)  
0030-9923/2021/0001-0001 \$ 9.00/0  
Copyright 2021 Zoological Society of Pakistan

<b>Research</b>	<b>Parameters</b>	<b>Result</b>
Hassan and Khan, 2013	Performance of crossbred dairy cattle at military dairy farms in Pakistan	Performance of crossbred dairy cattle in first lactation may be lower than earlier reports on crossbred cattle in Pakistan. This poor performance may be due to existing culling procedures used at military farms. Every female offspring born at military farms are retained for raising as cow to produce milk. No clear criteria are followed for selection of female calves born at farms.
Faid-Allah, 2015	Genetic and non-genetic analysis for milk production and reproductive traits in Holstein cattle in Egypt	The Holstein breed in Egypt can show high milk production and good reproductive traits under adequate circumstances. Moderate heritability estimates and positive genetic correlation for most of traits studied suggested that genetic improvement of these traits would be achieved via multi-trait selection. The high positive genetic correlations between traits especially productive ones clarified that these traits could be improved simultaneously via multi-trait selection breeding program.
Fernando <i>et al.</i> , 2016	Productive and reproductive performance of Jersey cattle in the hill country of Sri Lanka	The values for total milk yield, birth weight, lactation length and number of services per conception falls within the average values observed in other tropical countries. However, values for calving interval, age at first calving, calving to service period and days open were higher than the optimum values recommended to maintain reproductive efficiency.

Online First Article