

Population Dynamics of Sucking Insect Pest and Predator, *Chrysoperla carnea* in Transgenic *Bt* Cotton and Non *Bt* Cotton Varieties

Qadeer Ahmed Soomro^{1*}, Mubasshir Sohail¹, Riffat Sultana², Niaz Hussain Khuhro¹ and Raza Muhammad Memon¹

¹Nuclear Institute of Agriculture (NIA), Tandojam-70060

²Department of Zoology, University of Sindh, Jamshoro-76080

ABSTRACT

Transgenic *Bt* cotton, initially introduced to combat bollworm infestations, inadvertently resulted in increased populations of sucking pests such as *Thrips tabaci*, *Bemisia tabaci*, and *Amerasia devastans* in *Bt* cotton fields. The heavy reliance on indiscriminate pesticide application to control these pests has detrimental effects on natural fauna. This study aimed to explore the efficacy of natural enemies as an alternative to pesticides for managing sucking pests in different varieties of *Bt* cotton. Field experiments were conducted on various *Bt* cotton varieties, including Super NIAB-602, Super NIAB-992, Super NIAB-3701, Super NIAB-886, Super NIAB-142, NIABt-102, IR-443, NIA-Bt-100, IR-1513, IR-NIBGE-1524, and the non-transgenic *Bt* variety Sadori. Population densities of *A. devastans*, *T. tabaci*, *B. tabaci*, and the predator *Chrysoperla carnea* were recorded on the upper, middle, and lower leaves of twenty plants in each group. Results revealed that Super NIAB-142 exhibited the highest population of *A. devastans* (10.00±0.57), followed by NIABt (8.00±0.57). Likewise, Super NIAB-602 demonstrated the highest population of *T. tabaci* (7.66±0.33), with IR-1513 ranking second (6.66±0.88). For *B. tabaci*, Super NIAB-142 displayed the highest population (10.00±0.57), closely followed by Super NIAB-602 (8.00±1.15). Conversely, Sadori exhibited the lowest populations of *A. devastans*, *T. tabaci*, and *B. tabaci*, with values of 0.33±0.33, 1.00±0.57, and 1.0±0.57, respectively. Additionally, Super NIAB-142 showed significantly higher populations of *C. carnea* eggs, larvae, and adults (0.54±0.0, 0.44±0.01, and 0.38±0.05), followed by Super NIAB-602. In contrast, Sadori displayed the lowest populations of *C. carnea* eggs, larvae, and adults (0.07±0.01, 0.16±0.01, and 0.06±0.06). These findings underscore the potential of utilizing natural enemies, particularly *C. carnea*, as a viable method for managing sucking pests in transgenic *Bt* cotton, thereby reducing the excessive dependence on pesticides.

Article Information

Received 31 December 2021

Revised 05 September 2023

Accepted 20 September 2023

Available online 30 December 2023 (early access)

Published 21 April 2025

Authors' Contribution

QAS planned the research project, wrote, revised and edited the manuscript. MS participated in the critical review and provide guidance during writing research manuscript. RS conceived and supervised the overall study and made critical revision. NH provide guidance during execution of experiment. RM made critical revisions to the manuscript and helped to analyze the data.

Key words

Transgenic, Population fluctuation, Natural enemies

INTRODUCTION

Cotton is a commercially important fiber and cash crop cultivated in tropical and warm temperate regions worldwide. It is grown annually and has various valuable products derived from its seeds, such as oil, lint, hulls, and animal feed (Ozyigit *et al.*, 2007). In Pakistan, cotton was grown on 3.009 million hectares in 2016-17, resulting in an average production of 14.101 million bales. In the Sindh province alone, cotton was cultivated on 0.660 million hectares, yielding an average of 4.500 million bales (Daily

Ali *et al.*, 2019). Cotton production is affected by various factors, and insect pests pose a major threat, causing significant yield reduction. Among the insect pests, the bollworm complex and sucking complex are the two major types damaging cotton crops. Economically damaging bollworms include spotted bollworm (*Earias vitella*), American bollworm (*Helicoverpa armigera*), and pink bollworm (*Pectinophora gossypiella*). To modernize agriculture and manage insect pests, the introduction of insecticidal crop cultivars with transgenic traits has played a significant role globally. Introducing new resistance genes into economically important crops can help develop insect resistance and integrate pest management strategies (Gatehouse and Gatehouse, 1998). *Bt* cotton, which is genetically modified to resist bollworm infestations, was introduced. However, gradually, sucking insects such as whiteflies (*Bemisia tabaci*), thrips (*Thrips tabaci*), and jassids (*Amerasia devastans*) became serious problems for transgenic cotton. Losses in *Bt* cotton ecosystems due to sucking pests have been reported to reach up to 8.37% (Banerjee, 2002). Cotton growers indiscriminately use

* Corresponding author: soomroqadeer77@yahoo.com
0030-9923/2025/0003-1067 \$ 9.00/00



Copyright 2025 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

pesticides to combat these sucking insect pests. Globally, there is a growing concern to implement alternative methods of crop protection that reduce the use of agrochemicals. Integrated pest management systems rely on continuous field monitoring, physical and biological control, and the performance of natural enemies in cotton ecosystems (Konradsen, 2007; Eddleston *et al.*, 2002). Natural enemies, including *Coccinella septempunctata* Linnaeus, *Chrysoperla carnea* Stephen, *Menochilus sexmaculatus* Fabricius, *Trichogramma brasiliensis* (Ashmead), *Trichogramma achaeae* Nagraja, spiders, and *Chelonus blackburni* Cameron, play a significant role in managing sucking insect pests in cotton crops (Dhaka and Pareek, 2007). *C. carnea*, in particular, has contributed greatly to the biological control of sucking insect pests, helping to keep their populations below economically damaging levels (Hoy and Nguyen, 2000). The population development of predators depends on the availability of prey, and beneficial insect populations thrive in field conditions where their food sources are abundant (Solangi *et al.*, 2005). Predators like coccinellids (Coleoptera: Coccinellidae) are highly efficient in managing small and soft-bodied insects, and more than 400 species of these predators have been documented worldwide (Michaud, 2001). The incidence of insect pests greatly reduces the yield and value of cotton crops. The upsurge in the occurrence of these insect pests is primarily dependent on relative humidity, rainfall, and temperature (Aheer *et al.*, 1994). Considering the losses caused by various sucking insect pests in cotton crops, this study focuses on the population dynamics of these pests and the predator *Chrysoperla carnea*.

MATERIALS AND METHODS

Studies on the seasonal occurrence of sucking pests and the predator *C. carnea* in transgenic cotton varieties, along with a non-transgenic variety, were conducted during the Kharif season of 2020 at the Nuclear Institute of Agriculture (NIA) Experimental Farm. The cotton varieties used in the experiment included Super NAIB-602, Super NIAB-992, Super NIAB-3701, IR-443, NIABt-100, IRNIBGE-1524, IR-1513, NIABt-102, Super NIAB-142, Super NIAB-886, and the non-*Bt* cotton variety Sadori. Agronomic practices recommended for cultivation were followed in the experimental plots, and no plant protection measures were applied. The experiment was set up using a randomized complete block design (RCBD), with three replications, and the selected plot size was 15x3 m. Observations on the population densities of sucking insect pests such as jassids (*A. devastans*), thrips (*T. tabaci*), Whiteflies (*B. tabaci*), and the field population of the predator *C. carnea* were recorded from three leaves of five

randomly selected plants at weekly intervals, starting 30 days after the cotton crop germinated. These three leaves were chosen from the top, middle, and bottom canopy of the plants.

Statistical analysis

Statistical analysis of the data was conducted using Statistix® version 8.1. The analysis involved the use of ANOVA (analysis of variance) to determine significant differences between the treatments. Additionally, the least significant difference (LSD) test, as described by Steel *et al.* (1997), was used to compare means and identify statistically significant variations among the treatment groups.

RESULTS

Studies on population dynamics of the predator *Chrysoperla carnea* and sucking insect pests in transgenic *Bt* cotton and non-*Bt* cotton concluded that lower numbers of sucking pests were recorded in the cotton variety Sadori compared to other *Bt* cotton varieties.

Sucking insect pests

Jassids (*Amerasia devastans*)

The population of *A. devastans* was higher throughout the cropping season, as shown in Figure 1A. The maximum population of *A. devastans* ($10.0 \pm 0.6a$ per 5 leaves) was observed in Super NIAB-142, followed by NIABt-100 ($8.0 \pm 0.6ab$ per 5 leaves). The population of *A. devastans* on other varieties decreased in the following order: Super NIAB-886 ($2.6 \pm 0.3g$ per 5 leaves), Super NIAB-602 ($3.0 \pm 0.6fg$ per 5 leaves), Super NIAB-992 ($3.3 \pm 0.8efg$ per 5 leaves), IR-443 ($3.6 \pm 0.3efg$ per 5 leaves), Super NIAB-3701 ($5.0 \pm 1.1def$ per 5 leaves), IR-NIBEG-1524 ($5.3 \pm 0.3cde$ per 5 leaves), NIABt-102 ($7.0 \pm 1.1bcd$ per 5 leaves), IR-1513 ($7.3 \pm 0.6bc$ per 5 leaves). The lowest population of *A. devastans* ($0.3 \pm 0.3h$ per 5 leaves) was observed in the non-*Bt* cotton variety Sadori.

Thrips (*Thrips tabaci*)

The results regarding the *T. tabaci* in transgenic *Bt* cotton and non-*Bt* cotton throughout the cropping season are shown in Figure 1B. The maximum population of *T. tabaci* ($7.6 \pm 0.3a$ per 5 leaves) was observed in the Super NIAB-602 variety, followed by IR-1513 ($6.6 \pm 0.8ab$ per 5 leaves). Similarly, the population of *T. tabaci* decreased in the following order in other varieties: Super NIAB-886 ($2.0 \pm 0.5gh$ per 5 leaves), Super NIAB-3701 ($3.0 \pm 0.5fg$ per 5 leaves), NIABt-102 ($3.3 \pm 0.6efg$ per 5 leaves), Super NIAB-142 ($4.0 \pm 0.6def$ per 5 leaves), Super NIAB-992 ($5.0 \pm 0.6bcde$ per 5 leaves), IR-443 ($5.6 \pm 0.3bcd$ per 5 leaves), NIABt-100 ($6.0 \pm 1.1abc$ per 5 leaves), and IRNIBGE-1524 ($4.6 \pm 0.3cdef$ per 5 leaves). However, the

lowest population of *T. tabaci* (1.0 ± 0.6 h per 5 leaves) was observed in the non-Bt cotton variety Sadori.




Fig. 1. Average population density of *Amrasca devastans* (A), *Thrips tabaci* (B) and *Bemisia tabaci* (C) on transgenic Bt and nonBt varieties.

White flies, Bemisia tabaci (Gennad)

The population of *Bemisia tabaci* was comparatively higher than that of jassids throughout the entire cropping season, as shown in Figure 1C. The maximum population of *B. tabaci* (10.0 ± 0.6 a per 5 leaves) was observed in the transgenic Bt cotton variety Super NIAB-142, followed by Super NIAB-602 (8.0 ± 1.1 ab per 5 leaves). However, the population of *B. tabaci* decreased in the following order in other transgenic Bt cotton varieties: Super NIAB-992 (2.0 ± 0.6 g per 5 leaves), Super NIAB-3701 (3.0 ± 0.6 fg per 5 leaves), Super NIAB-886 (3.3 ± 0.3 defg per 5 leaves), NIABt-102 (4.33 ± 0.88 cdef per 5 leaves), IR-443 (4.66 ± 1.20 de per 5 leaves), IR-1513 (5.6 ± 0.6 bcd per 5 leaves), IR-NIBGE-1524 (6.3 ± 0.8 bc per 5 leaves), and NIABt-100 (6.3 ± 0.8 bc per 5 leaves). The lowest population of *B. tabaci* (1.0 ± 0.6 g per 5 leaves) was recorded in the non-Bt cotton variety Sadori.

Chrysoperla carnea (Stephen)

C. carnea eggs

The population of *C. carnea* eggs was observed throughout the entire cropping season, as shown in Figure 2A. The highest population of *C. carnea* eggs (0.57 ± 0.03 a per 5 leaves) was noticed in the transgenic Bt cotton variety Super NIAB-142, followed by Super NIAB-602 (0.41 ± 0.01 b per 5 leaves). In descending order, the population of *C. carnea* eggs in other transgenic Bt cotton varieties was as follows: Super NIAB-992 (0.25 ± 0.02 d per 5 leaves), Super NIAB-886 (0.35 ± 0.01 c per 5 leaves), IR-443 (0.17 ± 0.1 ef per 5 leaves), NIABt-100 (0.19 ± 0.1 e per 5 leaves), IR-1513 (0.12 ± 0.03 fg per 5 leaves), IR-NIBGE-1524 (0.10 ± 0.03 fg per 5 leaves), NIABt-102 (0.13 ± 0.02 ef per 5 leaves), and Super NIAB-3701 (0.21 ± 0.01 de per 5 leaves). However, the population of *C. carnea* eggs was found to be the lowest (0.07 ± 0.01 h per 5 leaves) in the non Bt cotton variety Sadori.

C. carnea larvae

The population trend of *Chrysoperla carnea* larvae was observed throughout the entire cropping season in both transgenic Bt cotton varieties and non-Bt cotton variety, as documented in Figure 2B. The highest number of *C. carnea* larvae (0.44 ± 0.01 a per 5 leaves) was noticed in the transgenic Bt cotton variety Super NIAB-142, followed by Super NIAB-602 with a population of (0.38 ± 0.01 b per 5 leaves). On the other hand, the population of *C. carnea* larvae decreased in the following order in other transgenic Bt cotton varieties: Super NIAB-886 (0.29 ± 0.01 efg per 5 leaves), Super NIAB-992 (0.29 ± 0.02 def per 5 leaves), NIABt-100 (0.36 ± 0.1 bcd per 5 leaves), IR-443 (0.22 ± 0.01 hi per 5 leaves), IR-NIBGE-1524 (0.21 ± 0.03 gh per 5 leaves), Super NIAB-3701 (0.31 ± 0.01 cde per 5 leaves), NIABt-102 (0.36 ± 0.01 bc per 5 leaves), and IR-1513

(0.21 ± 0.01 ghi per 5 leaves). Likewise, the lowest number of *C. carnea* larvae (0.16 ± 0.01 i per 5 leaves) was noticed in the non-*Bt* cotton variety Sadori.




Fig. 2. Average population density of *Chrysoperla carnea* eggs (A), larvae (B) and adults (C) on transgenic *Bt* and non *Bt* varieties.

C. carnea adults

The adult population of the predator *C. carnea* was observed throughout the entire cropping season, as demonstrated in Figure 2C. The maximum population of *C. carnea* (0.38 ± 0.05 a per 5 leaves) was observed in the transgenic *Bt* cotton variety Super NIAB-142, followed by Super NIAB-602 with a recorded population of (0.32 ± 0.01 ab per 5 leaves). The population trend decreased in the following order in other transgenic *Bt* cotton varieties: Super NIAB-886 (0.26 ± 0.01 bc per 5 leaves), NIABt-102 (0.24 ± 0.01 bcd per 5 leaves), IR-NIBGE-1524 (0.13 ± 0.01 ef per 5 leaves), IR-1513 (0.10 ± 0.01 ef per 5 leaves), Super NIAB-992 (0.18 ± 0.01 de per 5 leaves), Super NIAB-3701 (0.12 ± 0.01 ef per 5 leaves), NIABt-100 (0.22 ± 0.01 cd per 5 leaves), and IR-443 (0.11 ± 0.01 ef per 5 leaves). However, the least population *C. carnea* adults (0.06 ± 0.00 f per 5 leaves) was noticed in the non-*Bt* cotton variety Sadori.

DISCUSSION

Sucking insect pests such as jassids, thrips, and whiteflies have become a serious problem in transgenic *Bt* cotton ecosystems compared to non-*Bt* cotton. The highest population of *A. devastans* was observed in the transgenic *Bt* variety Super NIAB-142, followed by NIABt-100. However, the jassid population reduced compared to other transgenic *Bt* cotton varieties, namely SuperNIAB-602, IR-443, SuperNIAB-992, IR-NIBGE-1524, SuperNIAB-886, SuperNIAB-3701, NIABt-102, and IR-1513. The lowest population of jassids was recorded in the non-*Bt* variety Sadori. According to Ashfaq *et al.* (2010) the maximum population *A. devastans* was 2.00 per leaf in the transgenic *Bt* genotype CP-1401, while the minimum population of 1.19 per leaf was observed in the non-transgenic genotype CIM-496. Similar findings on the population fluctuation of Jassids in transgenic *Bt* cotton were reported by Rekha *et al.* (2008). The highest population of thrips was observed in SuperNIAB-602, followed by IR-1513, while the population decreased on other varieties such as SuperNIAB-886, SuperNIAB-3701, NIABt-102, SuperNIAB-IR-NIBGE-1524, SuperNIAB-992, IR-443, and NIABt-100. The lowest population of thrips was found in the non-*Bt* cotton variety Sadori. The higher population of thrips on transgenic *Bt* cotton compared to traditional cultivars, as reported by Men *et al.* (2005) and Naveen *et al.* (2007), supports our findings. Similar results on the population fluctuation of *T. tabaci* on *Bt* cotton and non-*Bt* cotton were also examined by Godhani (2006), who reported a maximum thrips population of 23.66 per 3 leaves in the *Bt* cotton system. Regarding whiteflies (*B. tabaci*), their population in transgenic *Bt* cotton was comparatively higher than that of Jassids over the entire cropping season.

The highest populations of *B. tabaci* were observed in Super NIAB-142, followed by Super NIAB-602. The population of *B. tabaci* decreased in SuperNIAB-992, Super NIAB-3701, SuperNIAB-886, NIABt102, IR-443, IR-1513, IR-NIBGE-1524, and NIABt-100 ($6.33 \pm 0.88bc$ per 5 leaves). The least population of *B. tabaci* was found in the non-*Bt* cotton variety Sadori. Similar results on a higher incidence of *B. tabaci* in transgenic *Bt* cotton hybrids, Super NIAB-886, NIAB-992, NIAB-3701, NIABt-100, IR-443, NIABt 102, and IR- NIBGE-1524, were reported by Jeyakumar *et al.* (2008). *C. carnea* plays a vital role in reducing the population of these insect pests in the transgenic *Bt* cotton ecosystem. However, the results revealed that the population of *C. carnea* eggs, larvae, and adults were higher in the transgenic *Bt* cotton variety Super NIAB-142, followed by SuperNIAB-602. The population of *C. carnea* decreased in other transgenic *Bt* cotton varieties, including SuperNIAB-886, SuperNIAB-3701

CONCLUSION

The study concluded that transgenic *Bt* cotton varieties harboured a higher population of sucking insects, including *Thrips tabaci*, *Amrasica devastans*, and *Bemisia tabaci*, compared to the non-transgenic *Bt* cotton variety Sadori. Additionally, transgenic *Bt* cotton varieties exhibited increased activity of the generalist predator *Chrysoperla carnea* in comparison to the non-transgenic *Bt* cotton variety Sadori.

ACKNOWLEDGEMENT

Authors wish to acknowledge Biological Control Group, Plant Protection Division, Nuclear Institute of Agriculture, Tandojam for providing conducive environment for research.

Funding

The study received no external funding.

Ethical statement

We ensure that all research is conducted in accordance with ethical principles. Neither human was the subject in this research nor such kind of animal, which required any administrative approval.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Aheer, G.M., Ahmed, K.J. and Ali, A., 1994. Role of weather in fluctuating aphid density in wheat crop. *J. agric. Res.*, **32**: 295-301.
- Ali, M.A., Farooq, J., Batool, A., Zahoor, A., Azeem, F., Mahmood, A. and Jabran, K., 2019. Cotton production in Pakistan. In: *Cotton production* (eds. K. Jabran and B.S. Chauhan). pp. 249-276.
- Ashfaq, M., Noor-ul-Aen, M., Zia, K., Nasreen, A. and Mansoor-ul-Hasan, 2010. The correlation of abiotic factors and physico-morphic characteristics of (*Bacillus thuringiensis*) *Bt* transgenic cotton with whitefly, *Bemisia tabaci* (Homoptera: Aleyrodidae) and jassid, *Amrasca devastans* (Homoptera: Jassidae) populations. *Afr. J. agric. Res.*, **5**: 3102-3107.
- Badiger, H.K., Patil, S.B., Udikeri, S.S. and Chattannavar, S.N., 2012. Comparative efficacy of single (BG I) and stacked *Bt* cotton hybrids (BG II) under integrated pest management practices. *J. Cotton Res. Dev.* **26**: 117-123.
- Banerjee, S.K., 2002. *Estimation of losses due to major pests of cotton*. CICR, Annual Report 2001- 02, pp. 54-55.
- Dhaka, S.R. and Pareek, B.L., 2007. Seasonal incidence of natural enemies of key insect pests of cotton and their relationship with weather parameters. *J. Pl. Prot. Res.*, **47**: 418-419.
- Dhillon, M.K. and Sharma, H.C., 2013. Comparative studies on the effects of *Bt*-transgenic and non-transgenic cotton on arthropod diversity, seed cotton yield and bollworms control. *J. environ. Biol.* **34**: 67-73.
- Dhillon, M., Pampapathy, K.G., Wadaskar, R.M. and Sharma, H.C., 2012. Impact of *Bt* transgenic cottons and insecticides on target and non- target insect pests, natural enemies and seed cotton yield in India. *Indian J. agric. Res.*, **82**: 248-254.
- Eddleston, M., Karalliedde, L., Buckley, N., Fernando, R. and Hutchinson, G., 2002. Pesticide poisoning in the developing world a minimum pesticides list. *Lancet*, **360**: 1163-1167. [https://doi.org/10.1016/S0140-6736\(02\)11204-9](https://doi.org/10.1016/S0140-6736(02)11204-9)
- Gatehouse, A.M.R. and Gatehouse, J.A., 1998. Identifying proteins with insecticidal activity: Use of encoding genes to produce insect-resistant transgenic crops. *Pestic. Sci.*, **52**: 165-175. [https://doi.org/10.1002/\(SICI\)1096-9063\(199802\)52:2<165::AID-PS679>3.0.CO;2-7](https://doi.org/10.1002/(SICI)1096-9063(199802)52:2<165::AID-PS679>3.0.CO;2-7)
- Godhani, P.H., 2006. *Impact of intercropping on the insect pests suppression in Hybrid cotton-10*. Ph.D thesis, Anand Agricultural University, Anand, Gujarat (India).
- Hoy, M.A. and Nguyen, R., 2000. Classical biological

- control of brown citrus aphid: Release of *Lipolexis scutellaris*. *Citrus Indust.*, **81**: 24-26.
- Jeyakumar, P.R., Tanwar, C.K.M., Singh, A., Monga, D. and Bambawale, O.M., 2008. Performance of *Bt* cotton against sucking pests. *J. Biopest.*, **1**: 223-225.
- Konradsen, F., 2007. Acute pesticide poisoning a global public health problem. *Danish med. Bull.*, **54**: 58-59.
- Men, X.Y.F., Yardim, G.E. and Parajulee, M.N., 2005. Behavior response of *Helicoverpa armigera* (Lepidoptera: Noctuidae) to cotton with and without expression of the CryIAc delta-endotoxin protein of *Bacillus thuringiensis* Berliner. *J. Ins. Behav.*, **18**: 33-50. <https://doi.org/10.1007/s10905-005-9345-9>
- Men, X.Y., Ge, F., Liu, X.H. and Yardim, E.N., 2003. Diversity of arthropod communities in transgenic *Bt* cotton and non-transgenic cotton agro ecosystems. *Environ. Ent.*, **32**: 270-275. <https://doi.org/10.1603/0046-225X-32.2.270>
- Michaud, S.P., 2001. Population dynamics of bean aphid (*Aphis craccivora* Koch) and its predatory coccinellid complex in relation to crop type (Lentil, Lathyrus and Faba bean) and weather conditions. *J. entomol. Res.*, **18**: 25-36.
- Naveen, A., Brar, D.S. and Butter, G.S., 2007. Evaluation of *Bt* and non *Bt* version of two cotton hybrid under different spacing against sucking insect pest and natural enemies. *J. Cotton Res. Dev.*, **21**: 106-110.
- Ozyigit, I.I., Kahraman, M.V. and Ercan, O., 2007. Relation between explants age, total phenols and regeneration response in tissue cultured cotton (*Gossypium hirsutum* L.). *Afr. J. Biotech.*, **6**: 3-8.
- Rafee, C.M., 2010. *Insect pest management in desi cotton*. Ph.D. thesis submitted to Department of Agricultural Entomology, College of Agriculture, Dharwad University of Agricultural Sciences, Dharwad, and Karnataka.
- Rekha, S., Nageswararao, G. and Dhurera, S., 2008. Effect of legume intercrops on yield and profitability of rain M. fed cotton in vertisols. *J. Cotton Res. Dev.*, **22**: 256-260.
- Sagar, D., Patil, B. and Bheemanna, M., 2011. Pest status and cry protein content in transgenic cotton. *J. Cotton Res. Dev.*, **25**: 258-262.
- Sharma, H.C. and Agarwal, R.A., 1983. Role of some chemical components and leaf hairs in varietal resistance in cotton to jassid, *Amrasca biguttula biguttula* Ishida. *J. entomol. Res. Soc.*, **7**: 145-149.
- Solangi, B.K., Talpur, M.A. and Nizamani, I.A., 2005. Population of spotted *Bollworm earias* spp. and its predators (Natural enemies) on cotton. *J. appl. Sci.*, **5**: 1402-1404. <https://doi.org/10.3923/jas.2005.1402.1404>
- Steel, R.G.D., Torrie, J.H. and Dicky, D.A., 1997. *Principles and procedures of statistics. A biometrical approach 3rd ed.* McGraw-Hill, Singapore.
- Udikeri, S.S., Patil, B., Basavanagoud, V.K., Khadi, B.M., Kulkarni, K.A. and Vamadevaiah, H.M., 2012. Impact of *Bt* transgenic cotton on population dynamics of aphids and natural enemies. *Indian J. agric. Sci.*, **82**: 555-560.