



## Supplementary Material

# Short Communication: Specific PCR-based Marker for Identification of Common Carp ‘Hainan’ in China

Muhammad Younis Laghari<sup>1,2</sup>, Zixia Zhao<sup>1</sup>, Yiwen Gong<sup>3</sup>, Punhal Lashari<sup>1,2</sup>, Shangqi Li<sup>1</sup>, Jian Xu<sup>1</sup>, Ateeque Rahman Khooharo<sup>4</sup>, Jiongtang Li<sup>1,\*</sup> and Yan Zhang<sup>1\*</sup>

<sup>1</sup>Key Laboratory of Aquatic Genomics, Ministry of Agriculture, CAFS Key Laboratory of Aquatic Genomics and Beijing Key Laboratory of Fishery Biotechnology, Chinese Academy of Fishery Sciences, Beijing, 100141, China

<sup>2</sup>Department of Freshwater Biology and Fisheries, University of Sindh, Jamshoro, 76080, Pakistan


<sup>3</sup>National Demonstration Center for Experimental Fisheries Science Education (Shanghai Ocean University)

<sup>4</sup>Centre of Excellence in Marine Biology, Karachi University, Sindh-Pakistan

\* Corresponding author: [zhangy@cafs.ac.cn](mailto:zhangy@cafs.ac.cn); [lijt@cafs.ac.cn](mailto:lijt@cafs.ac.cn)

0030-9923/2020/0003-1177 \$ 9.00/0

Copyright 2020 Zoological Society of Pakistan



Supplementary Fig. 1. Specific PCR-based marker amplified in four populations of common carp. DL2000 Marker; A1-10, TL1 marker was amplified in HLJ (1-2), CJ (3-4), Hainan (5-6); 8-9, YLH (7-8) and repeat in Hainan (9-10). B1-10, SL1 marker was amplified in HLJ (1-2), CJ (3-4); Hainan (5-6); YLH (7-8) and TL1 marker was amplified in Hainan (9-10).