



Short Communication

Molecular Characterization and Pathogenesis of Local Isolates of *Clostridium perfringens* Type D from Sheep in District Peshawar

Saqib Nawaz¹, Sher Bahadar Khan^{1*}, Umar Sadique¹, Muhammad Israr², Mumtaz Ali Khan³, Hameed Ullah Khan⁴, Muhammad Irshad⁵, Ihsan Ali¹ and Haq Aman Ullah¹

¹College of Veterinary Sciences, The University of Agriculture, Peshawar

²Pakistan Science Foundation, Islamabad

³Civil Veterinary Hospital, Takht Bhai, Mardan

⁴Veterinary Research Institute, Peshawar

⁵Livestock Research and Developmental Station, Surezi, Peshawar

ABSTRACT

The main objective of this study was isolation and determination of changes caused by *Clostridium perfringens* type D from sheep. For this purpose, a total of 272 fecal samples were collected from sheep and lambs from four different zones of district Peshawar. Isolates were morphologically, biochemically and molecularly confirmed through Gram staining, gelatin liquefaction test (GLT) and polymerase chain reaction (PCR), respectively. Out of total, 85 (31.3%) were positive for *C. perfringens* type D on PCR. The results revealed 6-34% prevalence of *C. perfringens* type D in different regions. Similarly, significantly ($P < 0.001$) higher prevalence was observed in lambs (54%) as compared to sheep (23%). For gross and histopathological lesions a total of 50 animals were necropsied infected with *C. perfringens* type D. A total of 43 (86%) sheep were recorded having hemorrhages on serosa of intestine followed by gas filled small intestine in 41 sheep (82%), lesions in ileum and colon in 39 sheep (78%), soft kidney 27 (54%), dark congested liver 26 (52%), nephritis 23 (46%) and splenitis 13 (26%). Microscopic lesions revealed a score of 10 of kidney tissue out of 15 while the lesion score of intestine revealed 08 out of lesion score of 15.

Article Information

Received 03 February 2020

Revised 29 June 2020

Accepted 07 July 2020

Available online 08 June 2021

Authors' Contribution

SN, SBK and US designed the idea, performed experiments and wrote the article. M Israr, MAK and HUK did analysis and reviewed the article. M Irshad, IA and HAU collected and processed the samples.

Key words

Clostridium perfringens, Enterotoxaemia, Pathogenesis, Pulpy kidney disease, PCR

Pulpy kidney disease (PKD) is one of the major endemic diseases of small ruminant. This disease mainly occurs in sheep in region where there is abundant availability of green fodder and is caused by different types of toxins of *Clostridium perfringens* with high case fatality rates leading to considerable economic losses to the farmers. The disease mostly transmitted through feces to sheep and rarely to large ruminants (McClane *et al.*, 2006).

C. perfringens is non motile, spore forming, large rod shaped gram-positive, catalase-negative, oxidase-negative and anaerobic bacteria with an abundant ecological spreading in soil, sewage, diet, feces, and in the normal gut flora of humans and animals (McClane *et al.*, 2013). *C. perfringens* produces 4 major toxins which is categorized into 5 toxino-types (A, B, C, D and E), namely alpha (α) (CPA), beta (β) (CPB), Epsilon (ϵ) (ETX), and iota (ITX). α toxin is produced by all 5 toxino-types while β toxin is produced by *C. perfringens* type B and C. ϵ toxin is produced

by *C. perfringens* type B and D while iota toxin is produced by *C. perfringens* type E. *C. perfringens* also produces 15 other minor toxins including lethal toxins such as enterotoxin (CPE), perfringolysin O (PFO) and β 2 toxin (CPB2) (Garmory *et al.*, 2000). *C. perfringens* type A and B cause yellow lamb disease and lamb dysentery in sheep respectively, while *C. perfringens* type C causes haemorrhagic enteritis (struck in disease) and *C. perfringens* type D causes pulpy kidney disease.

In Pakistan, the prevalence of *C. perfringens* in small ruminants and sheep is 26% and 32.1%, respectively while some study reported 31% (Radad and Khalil, 2011). In Khartoum State, Iran and Al-Ahsa, Kingdom of Saudi Arabia the prevalence of *C. perfringens* in sheep was found to be 23.8% and 26.5%, respectively (Aschfalk *et al.*, 2002). Alteration in the gastro intestinal tract (GIT) is the predisposing factor for *C. perfringens* type D due to which it starts proliferation and produces large number of copies of ϵ toxin in the form of inactive prototoxin which get activated by proteolytic degradation of C-terminal 14 amino acid which produces both local and systemic effects (Cheung *et al.*, 2004). In kidney it lead to autolysis while in lungs it

* Corresponding author: sbvetdr@yahoo.com
0030-9923/2021/0004-1595 \$ 9.00/0
Copyright 2021 Zoological Society of Pakistan

cause pulmonary edema and in brain it lead to perivascular edema and cerebral necrosis (Garmory *et al.*, 2000).

Clinically in sheep pulpy kidney disease is mainly characterized by nervous signs like staggering and knuckling at fetlock in forelimbs, champing of jaws, teeth grinding, salivation, rapid shallow and irregular respiration along with sudden onset of death (Fernandez and Uzal, 2003). On postmortem examination, empty and gas filled small intestine, congested liver and dark, pulpy soft kidney, increased fluid in pericardial sac and petechial and ecchymotic hemorrhages in abdominal muscles and serosa of intestine in sheep are seen (Radad and Khalil, 2011). The main objectives of this study were isolation, histopathological and molecular identification of local isolates of *C. perfringens* type D from sheep and lambs to assess the prevailing situation of *C. perfringens* type D in district Peshawar, to reduce the economic losses and to make devise strategies for effective control of this fatal disease.

Materials and methods

Fecal samples were randomly collected from sheep and lambs suspected on clinical signs and symptoms for *C. perfringens* type D infection present in four regions of District Peshawar, Khyber Pakhtunkhwa province, Pakistan. A total of 272 fecal samples were used in the study including 200 from sheep (50 from each region) and 72 from lambs (18 from each region). Region I consist of Pelosi, university dairy farm and adjacent areas. Region II consists of Tajabad and Achenai bala. Region III consists of Nasir Pur and Charsadda Road while Region IV consists of GT Road and adjacent area. Samples were labeled, transported in ice box and stored at 4°C until use. A preformed questionnaire was developed to collect detail data of the animals.

Fecal sample were washed with phosphate buffer saline and culture in fluid thioglycollate broth (HI Media Laboratories Pvt. Ltd., India) for enrichment, followed by incubation at 37°C for 24h anaerobically using CO₂ incubator (Galaxy 48 S, New Brunswick an eppendorf Company) and observed for any growth. After obtaining mass turbidity in broth, the samples were subcultured on tryptose sulphite cycloserine (TSC) agar (HI Media Laboratories Pvt. Ltd., India) by taking 0.1ml of the inocula for obtaining suspected grayish black colonies having diameter of 3-5mm. The suspected isolates were initially confirmed through Gram staining and gelatin liquefaction test (Kalender *et al.*, 2005).

For PCR amplification of α and ϵ toxins, genomic DNA was extracted by using the commercially available DNA extraction kit (GeneAllBldg, 303-7 Dongnam-ro, Songpa-gu, Seoul, South Korea) according to the manufacturer's instructions.

PCR was conducted in 25 μ L reaction mixture

containing 10 μ l of MasterMix, 1 μ l each of Forward Primer and Reverse Primer, 8 μ l of PCR water and 5 μ l of DNA template using Bio-Rad thermal cycler. Amplification was done with initial denaturation at 94°C for 5 min followed by 30 cycles, each consisting of 60 sec of denaturation at 94°C, 60 sec of annealing at (α 55°C, ϵ 53.4°C), 60 sec of extension at 72°C and 10 min of final extension at 72°C. Two sets of primers used for the detection of *C. perfringens* type D α /cpa and ϵ /etx in fecal samples are:

α /cpa Primer F-5-TGC TAA TGTTAC TGC CGT TGA TAG-3
Primer R-5-TGC TAA TGTTAC TGC CGT TGA TAG-3
 ϵ /etx Primer F-5-ATT AAA ATC ACA ATC ATT CAC TTG-3
Primer R-5-CTT GTG AAG GGA CAT TAT GAG TAA-3
with products of base pair 247 and 206, respectively as described by Garmory *et al.* (2000).

PCR products were run on 1.5% agarose gel with gel red (3 μ l) added along DNA ladder of 1 Kb (Genesta™) as a Marker and optimized positive and negative control product at 120 V, charge of 500 mA current for 35min.

A total of 50 sheep were examined to study the gross and histopathological lesions. Similarly those sheep showing diarrhoea, staggering and knuckling at fetlock in forelimbs, champing of jaws and teeth grinding were suspected for Enterotoxaemia and were purchased and sacrificed for recording pathological lesions. On post mortem inspection, detailed lesions were recorded in small intestine, liver, kidney, pericardial sac and abdominal muscles. The detail lesions scoring were documented in different organs of each necropsied animal (Wesonga *et al.*, 2004). Tissues samples were collected from kidney and intestine in aseptic condition in sterile container under refrigeration for histopathology and preserved in neutral buffered formalin (10%) for histopathological examinations. Samples were properly labelled and transported to the Microbiology laboratory, Department of Animal Health, the University of Agriculture, Peshawar for further processing.

For gross and histopathological lesion scoring of different organs was made on the basis of severity of lesion. The severity level was 0, 1, 2, 3 and 4 that represent normal, mild, moderate, severe and highly severe, respectively (Sadique *et al.*, 2012). For histopathology a small cross-section were trimmed off from intestine and kidney. Then suitable sections were made by placing these samples in 10 % neutral buffered formalin for fixing of these samples. Then samples were further processed for histopathology and routine staining by following the method suggested by Bancroft *et al.* (2007). The data collected in the present study was analyzed through SPSS 8.1 using post Hoc and lesion scoring performed for evaluation of pathogenesis and severity of infection.

Results and discussion

C. perfringens type D was confirmed on PCR that

all 85 samples (46 from sheep and 39 from lambs) were positive for α and ϵ toxin gene (Table I). The maximum prevalence of *C. perfringens* type D in sheep and lambs was found in region III (34%) and (89%), respectively. On PCR analysis an amplicon size of 247 bp for α and 206 bp for ϵ was obtained that represents *C. perfringens* type D in sheep and lambs. Out of total 200 examined sheep, 123 (61.5%), 112 (56%), 105 (52.5%) and 81 (40.5%) showing diarrhoea, staggering and knuckling at fetlock in forelimbs, champing of jaws and teeth grinding, respectively. Along with these symptoms the frothing of mouth 75 (37.5%) and rapid shallow and irregular respiration 41 (20.5%) was also recorded. Most of the sheep's showing signs of mild fever 97 (48.5%) with anorexia, anemia and lethargy. Out of total 50 sheep died from pulpy kidney disease were examined for postmortem lesions, 43 (86%), 41(82%), 39(78%), 27(54%), 26(52%), 23(46%) and 13(26%) sheep showing hemorrhages on serosa of intestine, gas filled small intestine, lesions in ileum and colon, soft kidney, dark congested liver, nephritis and Spleenitis, respectively were recorded.

Table I. Incidence of *C. Perfringens* type D in sheep and lamb on PCR in Peshawar, KPK.

Regions	Sheep	Lamb
1	50/03	18/02
2	50/12	18/11
3	50/17	18/16
4	50/14	18/10
Total	200/46	72/39

Kidney and intestine were severely infected so the tissue sections were studied for histopathology. *C. perfringens* type D were confirmed that in kidney, it causes degenerative changes in the glomerulus, cast disposition, necrotic changes in tubular epithelium accompanied by congestion and leukocytic infiltration (Figs. 1A, B). The sections of intestine of sheep showed erosion of epithelium lining, sloughing of villi, leukocytic infiltration and aggregation of inflammatory cells (Figs. 1C, D). The

gross lesion score was carried out to record and evaluate the severity of *C. perfringens* type D infection in kidney and intestine of sheep. Out of total lesion score of 15 the distribution of lesion score was 08 in kidney while the lesion score of intestine is 07. However, the microscopic lesions score revealed that out of total lesion score of 15, the kidney present a score of 10 while the lesion score of intestine was 08 (Table II).

The isolates were identified on the basis of their culture, morphological and biochemical characteristics and its gross and histopathological lesion in its predirection sites (kidney and intestine). The isolates show grayish black colonies on TSC agar while Gram staining were also performed for morphological identification and show purple rods having spore which agree with the result of (Cheung *et al.*, 2004). During current study the prevalence rate in sheep




Fig. 1. Histological structure of sheep kidney (A, B) and intestine (C, D). Control kidney (A), sheep kidney infected with *C. perfringens* type D (B) showing leukocyte infiltration, cast deposition, tubular epithelial cells and severe degeneration in tubular epithelium. Control intestine (C), intestine of infected sheep (D) showing sloughing of villi and leukocytic infiltration (D). **Stain: H&E (Hematoxyllin and Eosin); Magnification: A, B, 100x; C, D, 400X.

Table II. Gross and microscopic lesion score in sheep suspected for *C. perfringens* type D infection.

Tissue	Gross lesions	Scoring	Microscopic lesions	Scoring
Intestine	Serosae hemorrhages	2	Hemorrhagic enteritis	2
	Enteritis	2	Sloughing of Villi	3
	Bloody ingesta	1	Necrosis of epithelial cells	3
	Edematous	2	Leukocytic infiltration	0
Kidney	Pyelonephritis	3	Loss of glomerulus	3
	Pus in hilus and pelvis	2	Distal tubular necrosis	2
	Focal necrosis	3	Nephrosis	2
			Congestion	3
			Leukocyte Infiltration	0

Gross lesions scoring on the bases of tissue severity; 0, Normal; 1, Mild; 2, Moderate; 3, Severe; 4, highly severe.

in region I, II, III and IV were (06%), (24%), (34%) and (28%) while the overall prevalence of enterotoxaemia in sheep is 23% which is in line with the finding of (Greco *et al.*, 2005) from Italy who reported 25% prevalence of enterotoxaemia in sheep. These results are in contrast with the result of (Bachhil and Jaiswal, 1989) who reported less prevalence when compared with our findings. This high occurrence rate may be due to different climatic condition, breed, and poor farm management, system of rearing and lack of information about disease to farmer community.

The prevalence rate in lamb in region I, II, III and IV was 11%, 61%, 88% and 55% while the overall prevalence of enterotoxaemia in lambs is 54% which is in close accordance with the result of (Kalender *et al.*, 2005) from turkey who reported 50% prevalence of enterotoxaemia in lambs. The results are in contrast with results of (Aschfalk *et al.*, 2002) from West Africa who reported 71% prevalence of enterotoxaemia in lambs. The high occurrence rate of (Aschfalk *et al.*, 2002) may be due to random selection of sheep, different climatic condition, breed and system of rearing.

PCR is used for molecular identification of toxinotyping of *C. perfringens* type D due to its high efficiency and sensitivity. α and ϵ primers with an amplicon size of 247bp and 206bp respectively were used. As per PCR findings, out of 272 suspected fecal samples (200 from sheep and 72 from lamb), 85 samples were positive for α and ϵ toxin which confirmed that *C. perfringens* type D is the most predominant type of *C. Perfringens* in sheep and lamb. These results are in line with (Kalender *et al.*, 2005).

Beside it the histopathological changes in intestine are dependent on the involvement of *C. perfringens* type D that causes lesions in the form of hyperplasia, granuloma, enteritis, congestion and fibrosis. There was sloughing of villi, desquamation of epithelial cells, hyperplasia of serosa and scattered hemorrhages in serosa layer of intestine infiltrated with leukocytes. The lesions noted in the current study are in line with the findings of (Kalender *et al.*, 2007). The section of kidney showed congestion, extensive degeneration in the glomeruli, and necrosis in tubular epithelium, urinary tubules filled with cast and loss of glomeruli.

Conclusion

The overall prevalence of enterotoxaemia in sheep and lamb population in study area was 23% and 54% respectively. The region wise prevalence of enterotoxaemia in sheep in region I, II, III and IV were 06%, 24%, 34% and 28% respectively while the region wise prevalence of enterotoxaemia in Lamb in region I, II, III and IV were 11%, 61%, 88% and 55% respectively. Region III showed the highest incidence rate of 34% and 88% in sheep and lamb respectively. PCR confirmed the α and ϵ toxin with an amplicon size of 247bp and 206bp for the presence of

C. perfringens type D respectively while the gross and microscopic lesion scoring revealed that kidney is more severely infected than intestine.

Statement of conflict of interest

The authors have declared no conflict of interest.

References

- Aschfalk, A., Valentin-Weig, P., Muller, W. and Goethe, R., 2002. *Vet. Rec.*, **151**: 210–213. <https://doi.org/10.1136/vr.151.7.210>
- Bachhil, N. and Jaiswal, T.N., 1989. *Indian Vet. Med. J.*, **13**: 229–233. <https://doi.org/10.17953/aicr.13.3-4.5613516316x12124>
- Bancroft, J.M. and Gamble, M., 2007. *Theory and practice of histopathological techniques*. Churchill Livingstone, London. pp. 125-138.
- Cheung, J.K., Awad, M.M., McGowan, S.J. and Rood, I., 2004. *J. Vet. Microbiol.*, **105**: 130-136.
- Fernandez Miyakawa, M.E. and Uzal, F.A., 2003. *Vet. Res. Commun.*, **27**: 231-241. <https://doi.org/10.1023/A:1023348708599>
- Garmory, H.S., Chanter, N. and French, N.P., 2000. *Epidemiol. Infect.*, **124**: 61–67. <https://doi.org/10.1017/S0950268899003295>
- Greco, G., Madio, A., Buonavoglia, D., Totaro, M., Corrente, M., Martella, V. and Buonavoglia, C., 2005. *Vet. J.*, **170**: 346-350. <https://doi.org/10.1016/j.tvjl.2004.08.001>
- Kalender, H., Ertas, H.B., Cetinkaya, B., Muz, A., Arslan, N. and Kilic, A., 2005. *Vet. Med. Czech.*, **50**: 141-148.
- Kalender, H., Kili, A. and Atil, E., 2007. *Turk. J. Vet. Anim. Sci.*, **31**: 83-84.
- McClane, B.A., Uzal, F.A., Miyakawa, M.F., Lyerly, D. and Wilkins, T., 2006. The Enterotoxic Clostridia. In: *The prokaryotes*, Vol. 4 (eds. M. Dworkin, S. Falkow and E. Stackebrandt). Springer-Verlag, NY, pp. 698-752. https://doi.org/10.1007/0-387-30744-3_22
- McClane, B.A., Robertson, S.L. and Chen, L.J., 2013. *Clostridium perfringens*. In: *Food microbiology: Fundamentals and frontiers*. (eds. R.L. Buchanan and M.P. Doyle). American Society for Microbiology, Washington, DC. pp. 465–489. <https://doi.org/10.1128/9781555818463.ch18>
- Radad, K. and Khalil, S., 2011. *Braz. J. Vet. Pathol.*, **4**: 219-224.
- Sadique, U., Zafer, R., Younas, Z.U., Hassan, M., Idrees, M., Mushtaq, A., Sajid, M. and Sabtain, M., 2012. *Pak. J. Anim. Pl. Sci.*, **22**: 33-37.
- Wesonga, H.O., Bolske, G., Thiaucourt, F., Wanjohi, C. and Lindberg, R., 2004. *Acta Vet. Scand.*, **45**: 167-179. <https://doi.org/10.1186/1751-0147-45-167>