

# Description of *Hemicycliophora pyri* n. sp., with Observations on *Psilenchus vincigurrae* Brzeski, 1991 and *Ditylenchus equilis* Heyns, 1964 (Nematoda: Tylenchida) from Gilgit-Baltistan, Pakistan

Sagir Hussain<sup>1</sup>, Erum Iqbal<sup>2\*</sup>, Nasira Kazi<sup>2</sup>, Sher Wali Khan<sup>1</sup>, Qamar Abbas<sup>1</sup> and Abdul Razaq<sup>1</sup>

<sup>1</sup>Department of Biological Sciences, Karakorum International University, Gilgit, Gilgit-Baltistan, Pakistan

<sup>2</sup>National Nematological Research Centre, University of Karachi, Karachi-75270, Pakistan

## ABSTRACT

The present study provides the morphological and morphometric characterization of some plant parasitic nematode populations, recovered from agricultural fields during surveys of the districts Gilgit and Nager, Gilgit-Baltistan, Pakistan. The analysis of samples yielded a new nematode species and two new reported species belonging to the order Tylenchida as new geographical records for Pakistan. *Hemicycliophora pyri* n. sp., is characterized by the broadly rounded lip region with two indistinct annuli, closely fitting sheath with 231-247 body annuli, stylet 95-100 µm long, non-raised and non-separated labial disc and gradually tapering long tail. *Psilenchus vincigurrae* Brzeski, 1991 and *Ditylenchus equilis* Heyns, 1964 have been reported as new record species from Pakistan. These species are described and briefly re-described along with their morphometric data, photomicrographs and illustrations.

## Article Information

Received 17 November 2022

Revised 10 April 2022

Accepted 11 May 2022

Available online 21 July 2022

(early access)

Published 21 July 2023

## Authors' Contribution

SH and QA performed survey, sampling and sample processing. EI and NK did identification and description. SWK reviewed the literature. DAR proofread the manuscript and support in the data sampling.

## Key words

*Hemicycliophora pyri* n. sp., New geographic records, *Psilenchus vincigurrae*, *Ditylenchus equilis* taxonomy, Gilgit-Baltistan

## INTRODUCTION

In the present study, systematic surveys were conducted of different areas of Gilgit, Baltistan to identify the economically important plant parasitic nematodes that damaged the crops. As a result, some plant parasitic nematode populations, recovered from the agricultural fields of the Districts Gilgit and Nager, Gilgit-Baltistan. The analysis of samples yielded a new nematode species *Hemicycliophora pyri* n. sp., and two new reported species *Psilenchus vinciguerrae* (Brzeski, 1991) and *Ditylenchus equilis* (Heyns, 1964) belonging to order Tylenchida as new records for Pakistan.

*Hemicycliophora pyri* n. sp. belongs to the superfamily Hemicycliophoroidea that received the common name

sheath nematodes due to the presence of an additional cuticle or sheath. The superfamily comprises two families, Hemicycliophoridae and Caloosiidae. The Hemicycliophoridae contain only a single genus *Hemicycliophora*, with 132 species while Caloosiidae comprises two genera *Caloosia* and *Hemicaloosia* with eight and nine species, respectively. They are obligate ectoparasites of plants and inhabit moist soil and aquatic environments. Several *Hemicycliophora* species are known to damage agricultural crops in different countries (Chitambar and Subbotin, 2014).

Members of Hemicycliophoroidea are vermiform, tapering at both extremities but more so posteriorly, rounded to truncate, slightly flattened at the anterior end with an attenuated or filiform tail. Body distinctly, slightly of not recessed immediately posterior to vulva. Two to three labial annuli not modified or separated. Vulva a transverse slit over 0.5 body diameter long, lips either modified and projecting, or not modified and rounded; vagina straight or curved. The male body is generally shorter and more slender than that of female, with a rounded anterior end, offset and tapering posterior end. Spicules arcuate, semi circular, or hook shaped, cloacal lip elongated to form a penial tube. Caudal alae covering less than one third of tail. Tail longer of female.

\* Corresponding author: erum\_i@yahoo.com  
0030-9923/2023/0004-1975 \$ 9.00/0



Copyright 2023 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Two species of the genus *Hemicycliophora* have been reported from Pakistan so far. Akhtar (1962) reported a known species *Hemicycliophora gracilis* Thorne, 1955 from *Agrostis* spp., for the first time from Lahore. A new species *Hemicycliophoraveechi* was described by Maqbool *et al.* (1986) as a new species from soil around the roots of herbaceous plants and grasses from slopes of hill around the lake of Saiful Muluk, Naran, Pakistan (Maqbool and Shahina, 2001). During the present study *Hemicycliophora pyri* n. sp., has been described from district Gilgit of Gilgit-Baltistan. Many species of the genus *Psilenchus* and *Ditylenchus* were reported earlier including new species viz., *P. khuzdarensis* (Khan *et al.*, 2004) and *D. bilqeesae* (Khan *et al.*, 2004; Zarina and Shahina, 2012; Shahina *et al.*, 2019). *Psilenchus vinciguerrae* (Brzeski, 1991) and *Ditylenchus equilis* (Heyns, 1964) have been reported as new records from district Nager for the first time from Gilgit, Baltistan.

## MATERIALS AND METHODS

Soil samples were collected from different agricultural fields of districts Gilgit and Nager, Gilgit-Baltistan. Nematodes were extracted from soil samples by using Cobb sieving and decanting technique (Cobb, 1918), followed by modified Baermann funnel method (Baermann, 1917). Nematodes were then gently heat killed, fixed in TAF (Tri-ethanol-amine Formaldehyde) solution (Courtney *et al.*, 1955). From TAF, nematodes were transferred to glycerine (Seinhorst, 1959) and subsequently mounted on microscopic slides for identification (Siddiqi, 2000). Measurements, including Demanian indices (de Man, 1880) and other ratios, were taken with the compound microscope Nikon Eclipse E400, equipped with a drawing tube (Camera Lucida) attached to it for illustrations. Photomicrographs were made with a Nikon DS, Film camera, attached to the same microscope.

### *Hemicycliophora pyri* n.sp. (Fig. 1, Table I)

#### Description

##### Female

Body straight or very slightly curved ventrad. Cuticular sheath closely fitting, sheath annules slightly flattened than body annules. Lip region rounded with two annuli. Labial area with open amphids, labial disc not raised and merged with the lip annules. Labial frame weakly sclerotized. Stylet slender, slightly curved dorsad. Metenchium 80-88µm long and telenchium 12-17µm long. Stylet knobs well-sloped anteriorly with a distinct cavity posteriorly. Oesophagus typical of the genus. Median

oesophageal bulb amalgamated with procorpus 14-18µm wide with a distinct valve. Isthmus short encircled by nerve ring. Nerve ring 144-150µm from anterior end. Hemizonid not clearly seen. Excretory pore situated from two to four annuli posterior to base of oesophagus. Vulval lips elongated. Vulval sheath one to two annuli long, vagina thick walled. Gonad prodelphic, spermatheca round to oval, mostly filled with small round sperm cells. Oocytes arranged in a single row. Anus indistinct. Body posterior to vulva gradually narrows to about last quarter. Tail narrows uniformly, ending in a narrow rounded or sometimes slightly wider rounded terminus, annuli on last distinct and smaller than on rest of body.




Fig. 1. *Hemicycliophora pyri* n. sp., A, Whole body of female; B, Vulva and reproductive part; C, Oesophageal region of female; D, Vulva and anus with tail terminus; E, F, Variations of female vulvae shape.

##### Male

Not found.

#### Type habitat and locality

The specimens were collected from soil round the roots of apple (*Pyrus malus* L.) and cherry (*Prunus avium*

L.) from district Gilgit (village Danyour), Gigit-Baltistan, Pakistan.

**Table I. Morphometrics of *Hemicycliophora pyri* n.sp. (Measurments in  $\mu\text{m}$  except L).**

Morphological characters	Holotype female	Paratype females (n=15) Mean $\pm$ SD (range)
L	0.99	1.02 $\pm$ 0.05 (0.90-1.12)
a	21.5	21.01 $\pm$ 0.94 (19.8-23.0)
b	5.8	5.9 $\pm$ 0.50 (5.1-6.8)
c	12.6	10.35 $\pm$ 2.12 (10.7-12.6)
c'	3.1	3.28 $\pm$ 0.29 (3.0-4.1)
V%	85	83.05 $\pm$ 2.00 (80.1-85.7)
O	7.5	7.5 $\pm$ 1.09 (6-9.3)
DGO	6.5	7.3 $\pm$ 1.10 (6-9)
Stylet length	98	97.83 $\pm$ 2.03 (95-100)
Metenchium length	84	83.58 $\pm$ 3.01 (80-88)
Telenchium length	15.5	15.58 $\pm$ 1.32 (12-17)
Stylet knob width	6.5	7.07 $\pm$ 0.82 (6-8)
Stylet knob height	4.5	4.54 $\pm$ 0.44 (4-5)
Excretory pore from head end	178	185.90 $\pm$ 5.93 (177-196)
Width at mid body	49.2	49.27 $\pm$ 4.15 (45-58)
Lip region width	16.6	16.61 $\pm$ 1.07 (15-18)
Lip region height	5.8	6.07 $\pm$ 0.82 (5-7)
Annules width	5.5	5.58 $\pm$ 0.49 (5-6)
Tail length	94.6	94.61 $\pm$ 7.61 (80-108)
Oesophagus length	176.8	176.81 $\pm$ 3.2 (172-182)
Vulva- anus distance	55	58.33 $\pm$ 3.59 (54-66)
R	232	238.54 $\pm$ 10.44 (231-247)
Rst	26	26.0 $\pm$ 1.23 (24-28)
Roes	44.5	44.58 $\pm$ 1.97 (41-47)
Rex	47	47.45 $\pm$ 1.37 (44-49)
RV	45	44.6 $\pm$ 5.37 (36-49)
R Van	18.1	18.1 $\pm$ 1.58 (15-20)
Ran	22	24.3 $\pm$ 2.3 (21-29)
VL/VB	3.5	3.52 $\pm$ 0.33 (3.1-4.0)
St % L	9.4	9.45 $\pm$ 0.66 (8.4-11.0)
Vulval width	44.5	46.2 $\pm$ 2.18 (44-50)

#### Type material

Holotype slide no NNRC 126/1 and paratypes slide Nos 126/2-6 (14 female) deposited in the National Nematode Collection of NNRC University of Karachi, Karachi 75270, Pakistan. Slide no 126/7 (one female) deposited in the British Nematode Collection at the Food and Environmental Research Agency, Sand Hutton York, England.

#### Diagnosis and relationship

*Hemicycliophora pyri* n. sp., is characterized by the presence of broadly rounded, continuous lip region with two indistinct annuli, closely fitting sheath with 231-247 body annuli, stylet 95-100 $\mu\text{m}$  long non raised and non-separated labial disc and long gradually tapering tails.

According to the dicotomus key proposed by Chitambar and Subbotin (2014), the new species comes close to *H. subaolice*. Jairajpuri and Bagri (1973) in body and stylet length and number of body annules (R), but differs from it in labial disc which is not elevated vs elevated; lateral field without anastomoses vs lateral field, marked with anastomoses and there are no delicate scratches outside the lateral field annuli vs outside later field without lateral field annuli may be marked with numerous delicate scratches. It also differs from it by slightly higher number of Rst and Rvan (24-28 vs 19-24 and 15-20 vs 10), respectively.

#### *Psilenchus vinciguerrae* Brzeski, 1991 (Fig. 2, Table II)




Fig. 2. *Psilenchus vinciguerrae*. A, Female whole body; B, Male whole body; C, Vulval region; D, Head region; E, Oesophageal region; F, Male tail; G, Female tail.

**Table II. Morphometric data of *Psilenchus vinciguerrae* and *Ditylenchus equilis*. Measurements (except where indicated) are in  $\mu\text{m}$  and in the form mean  $\pm$  SD (range).**

Morphological characters	<i>P. vinciguerrae</i>		<i>D. equilis</i>
	Female (n=5)	Male (n=5)	Female (n=8)
L (mm)	0.89 $\pm$ 0.10 (0.69-0.98)	0.79 $\pm$ 0.07 (0.69-0.88)	0.61 $\pm$ 0.09 (0.54-0.84)
a	49.23 $\pm$ 1.56 (46.6-51.63)	42.94 $\pm$ 2.67 (39.15-46.2)	32.38 $\pm$ 2.81 (28.77-37.6)
b	6.77 $\pm$ 0.49 (6.02-7.56)	6.11 $\pm$ 0.29 (5.72-6.50)	5.19 $\pm$ 0.67 (4.21-6.39)
c	8.04 $\pm$ 0.29 (7.94-8.76)	6.90 $\pm$ 0.65 (6.50-8.05)	12.19 $\pm$ 1.51 (10.37-14.06)
c'	8.81 $\pm$ 1.05 (7.46-10.6)	9.23 $\pm$ 1.09 (7.81-10.86)	4.0 $\pm$ 0.61 (2.85-4.72)
V %	49.18 $\pm$ 0.57 (48.72-50.16)	-	83.15 $\pm$ 2.82 (81.06-90)
Stylet	12.8 $\pm$ 1.16 (11-14)	12.75 $\pm$ 0.82 (12-14)	8.12 $\pm$ 1.53 (7.0-12.0)
Excretory pore	97.8 $\pm$ 6.46 (86-104)	96 $\pm$ 10.08 (79-109)	86.87 $\pm$ 12.38 (78.0-112.0)
Oesophagus	132.4 $\pm$ 11.05 (116-148)	-	119.37 $\pm$ 11.93 (101-132)
Tail length	130.4 $\pm$ 192 (122-188)	-	50.12 $\pm$ 5.10 (40.0-60.0)
Maximum body width	18.2 $\pm$ 1.72 (18-20)	18.5 $\pm$ 2.17 (15-21)	12.75 $\pm$ 2.22 (11.0-18.0)
Anal body width	12.4 $\pm$ 2.24 (10-15)	-	19.12 $\pm$ 2.36 (15.0-24.0)
Spicule	-	18.25 $\pm$ 1.78 (16-20)	-
Gubernaculum	-	6.75 $\pm$ 0.43 (6.0-7.0)	-

*Morphological characterization**Female*

After killing with the gentle heat, the body slightly to strongly ventrally curved and open c-shaped. The head continuous, 2-4  $\mu\text{m}$  high. Body has small annules less than 0.1  $\mu\text{m}$  wide but below the anus 0.2-2.5  $\mu\text{m}$ . Lateral field with four lines and 1/2  $\mu\text{m}$  of the total body. Stylet 11-14  $\mu\text{m}$  in length without knobs and lumen pass through median bulb towards basal bulb. Amphids aperture invisible. The length of procrpus 70-80  $\mu\text{m}$  from the head and the length from middle of the median bulb to base of basal bulb 54-60  $\mu\text{m}$ . Isthmus 30-32  $\mu\text{m}$ . The nerve ring 90-110  $\mu\text{m}$  from the anterior region. The distance of excretory pore 86-104  $\mu\text{m}$  from anterior end at the upper side of basal bulb. Basal bulb is flask like 20-22  $\mu\text{m}$  long and oval in shape.

Hemizonid anterior to excretory pore. Ovary didelphic with vulval flaps at position of 48-50%. Spermatheca filled and 12-14  $\mu\text{m}$  long filled with sperms. Phasmids below the anus. Tail elongate filiform and 122-188  $\mu\text{m}$  in length.

*Male*

Males are similar to females and slightly smaller in body. Spicules ventrally curved and 16-20  $\mu\text{m}$  in length. Gubernaculum below the specula with 6-7  $\mu\text{m}$  long.

*Remarks*

This species is reported for the first time from Pakistan. Specimens of *Psilenchus vinciguerrae* were collected from the soil around the roots of potato from district Nager (village Hoper and Jaffarabad). Measurement and morphology of this population fit with the original description given by Brzeski, 1991. But the only difference was in body width (a= 37-43 vs 46-51  $\mu\text{m}$ ).


*Ditylenchus equilis* Heyns, 1964  
(Fig. 3, Table II)

Fig. 3. *Ditylenchus equilis*. A, Female whole body; B, Reproductive region; C, Oesophageal region; D, Anterior region; E, Anus with tail terminus; F, Lateral lines.



*Morphological characterization**Female*

After killing by gentle heat body slightly arcuate or opens "C" shaped but tail portion carved. Head 2-3µm high and 5-6µm wide, continuous with body. Body with small annuli 0.5-1µm at the vulval position. Stylet 8-12µm long, shaft 5-6 and conus 4-5µm long. DGO near the base of stylet at 1-2µm. Nerve ring 70µm from the head end. The length of isthmus 65-70µm. The length of median bulb 20-24µm and width 4-7µm, oval elongate in shape. Hemizonid near the excretory pore. Excretory pore 78-112µm from anterior end.

Ovary monodelphic. Vulva posterior 81-90%. The distance from vulva to anus 60-62µm. Lateral field with four incisures, 1/3 of the total body length, the maximum body width 15-24µm. The length of post uterine sac 12-14µm. The length of rectum 1µm. Tail length 40-60µm.

*Male*

Not found.

*Remarks*

This species is reported for the first time from Pakistan. Specimens of *Ditylenchus equalis* were collected from soil around the roots of pear district Nager (village Hoper). Measurement and morphology of this population fit with the original description given by Heyns (1964).

*Statement of conflict of interest*

The authors have declared no conflict of interest.

## REFERENCES

- Akhtar, S.A., 1962. Free-living nematodes inhabiting Lahore soils. *Agric. Pak.*, **13**: 64-80.
- Baermann, G., 1917. Eine einfache Method zur Affindung von Ankylostomum- (Nematoden)-larven in Erdproben. *Geneesk. Tijdschrift. Nederl. Indië*, **57**: 131-137.
- Brzeski, M.W., 1991. *Psilenchus vinciguerrae* n. sp. (Nematoda: Tylenchidae). *Nematologica*, **37**: 1-7. <https://doi.org/10.1163/187529291X00015>
- Chitambar, J.J., and Subbotin, S.A., 2014. Systematics of the sheath nematodes of the superfamily Hemicycliophoroidea. In: *Nematology monographs and perspectives* (eds. D.J. Hunt and R.N. Perry), 10, (Series Editors), Brill, Leiden, Boston, pp. 732. <https://doi.org/10.1163/9789004187894>
- Cobb, N.A., 1918. Estimating the nema population of soil. *Agric. Tech. Circ. Dept. Agric.*, **1**: 48.
- Courtney, W.D., Polley, D. and Miller, V.L., 1955. TAF, an improved fixative in nematodes technique. *Pl. Dis. Rep.*, **39**: 570-571.
- de Man, J.G., 1880. Die einheimischen, frei in der reinen Erde und im süßen Wasser lebenden Nematoden. Vorläufiger Bericht und descriptive systematischer Theil. *Tijdsch. Nederl. Dierk. Vereenig.*, **5**: 1-104.
- Heyns, J., 1964. *Aphelenchoides helices* n. sp. and *Ditylenchus equalis* n. sp. two new soil inhabiting nematodes. *South Afr. J. agric. Sci.*, **7**: 147-150.
- Khan, A., Batool, S.K., and Khatoon, N., 2004. Two new species of Tylenchidae, *Psilenchus khuzdarnsis* n. sp. and *Ditylenchus bilqeesae* n. sp. (Nematoda: Tylenchidae). *Proc. Parasitol.*, **38**: 81-87.
- Jairajpuri, M.S. and Baqri, Q.H., 1973. Four new species of Tylenchida. Nematodes of high altitudes in India. *Nematologica*, 19-30.
- Maqbool, M.A. and Shahina, F., 2001. *Systematic and distribution: Biodiversity of nematode fauna in Pakistan*. National Nematological Research Centre, University of Karachi, Karachi-75270 Pakistan, pp. 179.
- Maqbool, M.A., Shahina, F. and Zarina, B., 1986. Two new species of Hemicycliophorinae Skarbilovich, 1959 (Nematoda: Cricematidae) from Pakistan. *Pak. J. Nematol.*, **4**: 43-49.
- Seinhorst, J.W., 1959. A rapid method for the transfer of nematodes from fixative to anhydrous glycerin. *Nematologica*, **4**: 67-69. <https://doi.org/10.1163/187529259X00381>
- Shahina, F., Nasira, K., Firoza, K. and Erum, Y.I., 2019. Overview of the nematode fauna of Pakistan. *Pak. J. Nematol.*, **37**: 171-243. <https://doi.org/10.18681/pjn.v37.i02.p171-243>
- Siddiqi, M.R., 2000. *Tylenchida: Parasites of plant and insect*, 2<sup>nd</sup> Edition. CABI Publication, Wallingford, UK, pp. 833.
- Thorne, G., 1955. Fifteen new species of the genus *Hemicycliophora* with an emended description of *H. typica* de Man (Tylenchida: Cricematidae). *Proc. helminthol. Soc. Washington*, **22**: 1-16.
- Zarina, B., and Shahina, F., 2012. *Annotated bibliography on nematology in Pakistan*. 2<sup>nd</sup> Edition. National Nematological Research Centre, University of Karachi, Karachi-75270, Pakistan, pp. 850.