A Checklist of family Salticidae (Arachnida: Araneae) from some localities of Pakistan

NUSRAT MAJEED* & ABIDA BUTT

Department of Zoology, University of the Punjab, Quaid-i-Azam Campus, Lahore, Pakistan

ABSTRACT

Salticid fauna of Pakistan had not been comprehensively explored. This study was an attempt to provide inventory of salticids, which would be the baseline documentation for future studies in Pakistan. This survey was conducted from June 2013 to May 2017, to collect jumping spiders from twelve different localities of Pakistan. Specimens (2556) were collected which belong to twenty-eight species and twenty-one genera. This study added twenty-three first records of species from ten genera. The female of *Phintella indica* is in this study for the first time.

Key words: Distribution, taxonomy, jumping spiders, diurnal hunters

INTRODUCTION

Salticidae is the largest family of spiders that comprises of 625 genera and 5979 species (World Spider Catalog, 2017). Jumping spiders have diverse body forms and behaviors (Maddison et al., 2014). Despite the highest diversity of jumping spiders in the world among the Araneae fauna (Bao & Peng, 2002), they have been poorly studied in Pakistan. In Global Species Database of Salticidae, thirty three salticid species have been reported from Pakistan (Proszynski, 2016). Among these eleven were reported from Punjab, only one from Kashmir and Balochistan each and nineteen are uncertain species. However, according to worldwide database of jumping spiders thirty nine species of salticidae were recorded from Pakistan (Metzner, 2017). Due to diverse range of habitats such as grasslands, forests. deserts. mountains and other ecosystems, there is a possibility of greater diversity of salticids in Pakistan

Perveen et al. (2012) reported onlytwo species, that belong to different genera from Peshawar. Parveen et al. (2007) recorded nine genera with twenty-seven salticid species from Punjab. Four species of salticids were reported from Sargodha by Mukhtar et al. (2012). The checklist of spider fauna from Cholistan and

neighbouring areas has not reported any salticid species (Sial et al., 2012). Another checklist of from Sheringal, spider fauna Khvber Pakhtunkhwa has not presented any record of salticids (Perveen & Khan, 2015). Only two species, Plexippoides flavescens (O. Pickard-Cambridge, 1872) and Menemerus marginatus (Kroneberg, 1875) were recorded Balochistan province (Bauer et al., 2015). Ghazanfar et al. (2016) reviewed all checklists of spiders and concluded thatonly seven species of salticidswere recorded from Punjab, twenty six from Sindh and one from Khyber Pakhtunkhawa. More extensive and serious efforts are needed to explore salticid diversity in Pakistan. This study aimed to develop a comprehensive biodiversity inventory of family salticidae from Pakistan.

MATERIALS AND METHODS

Salticids were collected from localities like Jallo park (Lahore), Jinnah garden (Lahore), Botanical garden of University of the Punjab (Lahore), Kallar Kahar lake (Chakwal), Hastal village (Chakwal), Hazara University (Mansehra), Pai Reserve Forest (Sakrand), Lal Suhanra National Park (Bhawalpur), Derawar Fort (Bhawalpur), Islamia University

(Bahawalpur), Bhakkar and Dera Bakha at Sutlej river (Bahawalpur) as shown in(Fig.1).

Hand picking and jerking were the most effective and useful method due to their wandering and predatory nature. Specimens were collected directly from the ground or by picking up substrate e.g. leaf litter, bunches of grass. For the jerking method, a cloth was laid on the ground underneath contiguous branches of trees/shrubs. The sampled branches were shaken by hand, beating the foliage and their trunks with mallets for 30 seconds to dislodge spiders. Then spiders dropped on the cloth were collected. The plastic bags and glass vials were

also used to capture the spiders because they were very active. Specimens were washed with 95% alcohol and preserved in absolute alcohol with the proper labeling of collection site, date of collection, collector name and other important notes. Identifications up to species level was done by using the historic keys to the salticid groups by Simon (1897-1903), literature cited within the "World Spider Catalog" (2017) and the "Global Species Database of Salticidae" (Proszynski, 2016). Specimens housed at room temperature in Arachnology Lab, Department of Zoology, University of the Punjab, Quaid-i-Azam Campus, Lahore, Pakistan.




Fig. 1: Map showing different collection sites of salticids from Pakistan

RESULTS

A total of 2556 salticids were collected from twelve localities and identified to species level (Table I). Collected specimens belong to twenty two genera and twenty eight species. Due to immature specimens in genera *Siler*, *Rhene*,and *Sitticus*, their species could not identified. The most specious genus was *Myrmarachne* with four species. The most

dominant species were *Thyene imperialis, Menemerus marginattus, Menemerus bivittatus* and *Carrhotus tristis.* However, the least number of specimens were recorded from *Sitticus sp., Titanattus cretatus, Siler sp., Rhene sp.* and *Chrysilla lauta.* This study added first record of twenty-three species and ten genera listed in (Table I). This suggests that Pakistan has reasonable high diversity of salticids that needs to be explored more extensively.

Table I: Checklist of salticids biodiversity from Pakistan. All species are new records to Pakistan except no. 9, 12 and 20.

Sr. #	Species	No of specimens	Sex	Distribution	Coordinates
1	Bianor maculatus (Keyserling, 1883)	180	7 0 94	Botanical garden, University of the Punjab, Lahore Hastal village, Chakwal	31.5004N, 74.3008E 28.6377N, 77.0490E
2	Carrhotus tristis Thorell, 1895	250	8	Pai Forest, Sakrand Islamia University Bhawalpur	26.0593N,68.1346E 29.3960N, 71.6626E
3	Chrysilla lauta Thorell,1887	12	8	Lal Suhanra National Park, Bhawalpur	29.3170N,71.9045E
4	Harmochirus brachiatus (Thorell, 1877)	44	7 0 9	Botanical garden, University of the Punjab, Lahore Hastal village, Chakwal	31.5004N,74.3008E 28.6377N, 77.0490E
5	Hasarius adansoni (Audouin, 1826)	59	7 0 O+	Botanical garden , University of the Punjab, Lahore	31.5004N,74.3008E
6	Iciusalboterminus (Caleb, 2014)	10	3	Hastal village, Chakwal Pai Forest, Sakrand	28.6377N, 77.0490E26.0593N, 68.1346E

7	Langona bristowei Berland & Millot, 1941	140	50 0 +	Botanical garden , University of the Punjab, Lahore Pai Forest, Sakrand	31.5004N,74.3008E 26.0593N, 68.1346E
8	Langona alfensis Heciak & Prószyński, 1983	100	9	Pai Forest, Sakrand	26.0593N, 68.1346E
9	Marpissa tigrina Tikader, 1965	36	7 0 P	Kallar Kahar, Lahore Jallo Park, Lahore Jinnah Garden, Lahore Botanical garden, University of Punjab Lahore	32.7760N, 72.7008E 31.5755N, 74.4943E 33.5723N, 73.1748E 31.5004N, 74.3008E
10	Menemerus marginatus (Kroneberg, 1875)	291	% 0+	Kallar Kahar, Chakwal Jallo Park, Lahore Jinnah Garden, Lahore Lal Suhanra National Park, Bhawalpur Derawar Fort, Bhawalpur Hastal village, Chakwal Botanical garden, University of the Punjab, Lahore	32.7760N, 72.7008E 31.5755N, 74.4943E 33.5723N,73.1748E 29.3170N, 71.9045E 28.4622N, 71.2019E 28.6377N, 77.0490E 31.5004N, 74.3008E
11	Menemerus bivittatus (Dufour, 1831)	276	7 0 Q+	Kallar Kahar, Lahore Jallo Park, Lahore Jinnah Garden, Lahore Botanical garden University of the Punjab, Lahore Lal Suhanra National Park, Bhawalpur Derawar Fort, Bhawalpur Hastal village, Chakwal	32.7760N, 72.7008E 31.5755N, 74.4943E 33.5723N, 73.1748E 31.5004N,74.3008E 29.3170N, 71.9045E 28.4622N, 71.2019E 28.6377N, 77.0490E
12	Myrmarachne melanotarsa Wesolowska & Salm, 2002	135	8	Jallo Park, Lahore Jinnah Garden, Lahore Hazara university, Mansehra Islamia University, Bhawalpur Lal Suhanra National Park, Bhawalpur Botanical garden, University of the Punjab, Lahore	31.5755N, 74.4943E 33.5723N,73.1748E 34.4207N, 73.2506E 29.3960N, 71.6626E 29.3170N, 71.9045E 31.5004N, 74.3008E

13	Myrmarachne melanocephala MacLeay, 1839	60	9	Lal Suhanra National Park, Bhawalpur Botanical garden, University of the Punjab, Lahore	29.3170N, 71.9045E 31.5004N, 74.3008E
14	Myrmarachne tristis (Simon, 1882)	60	3	Lal Suhanra National Park, Bhawalpur Botanical garden, University of the Punjab, Lahore	29.3170N, 71.9045E 31.5004N, 74.3008E
15	Myrmarachne plataleoides (O. Pickard-Cambridge, 1869)	53	ð	Lal Suhanra National Park, Bhawalpur Botanical garden, University of the Punjab, Lahore Pai Forest, Sakrand	29.3170N, 71.9045E 31.5004N, 74.3008E 26.0593N, 68.1346E
16	Napoca insignis (O. Pickard-Cambridge, 1872)	43	8	Lal Suhanra National Park, Bhawalpur Botanical garden, University of the Punjab,Lahore Pai Forest, Sakrand	29.3170N, 71.9045E 31.5004N,74.3008E 26.0593N, 68.1346E
17	Phintella indica (Simon, 1901)	20	7 0 9	Hastal village, Chakwal Pai Forest, Sakrand	28.6377N, 77.0490E 26.0593N, 68.1346E
18	Plexippus clemens (O. Pickard-Cambridge, 1872)	130	3 0 9	Lal Suhanra National Park, Bhawalpur Botanical garden, University of the Punjab,Lahore Pai Forest, Sakrand Bhakkar	29.3170N, 71.9045E 31.5004N, 74.3008E 26.0593N, 68.1346E 31.6266N, 71.0617E
19	Plexippus petersi (Karsch, 1878)	81	7 0	Botanical garden, University of the Punjab,Lahore Pai Forest, Sakrand Bhakkar	31.5004N, 74.3008E 26.0593N, 68.1346E 31.6266N, 71.0617E
20	Plexippoides flavescens (O. Pickard-Cambridge, 1872)	6	3	Pai Forest, Sakrand	26.0593N, 68.1346E

21	Pseudicius admirandus Logunov, 2007	89	₹ 0 0+	Botanical garden, University of the Punjab,Lahore Jallo Park, Lahore Jinnah Garden, Lahore Hazara university, Mansehra	31.5004N, 74.3008E 31.5755N, 74.4943E 33.5723N,73.1748E 34.4207N, 73.2506E
22	Rhene sp.	10	Im mat ure	Hazara university, Mansehra	34.4207N, 73.2506E
23	Siler sp.	5	Sub adul t	Hastal village, Chakwal	28.6377N, 77.0490E
24	Sitticus sp.	2	Im mat ure	Derawar Fort,Bhawalpur	28.4622N, 71.2019E
25	Stenaelurillus lesserti Reimoser, 1934	50	7 0 9	Botanical garden, University of the Punjab,Lahore	31.5004N, 74.3008E
26	Telamonia dimidiata (Simon, 1899)	120	7 0 P	Botanical garden, University of the Punjab,Lahore Lal Suhanra National Park, Bhawalpur Jallo Park, Lahore Jinnah Garden, Lahore	31.5004N, 74.3008E 29.3170N, 71.9045E 31.5755N, 74.4943E 33.5723N,73.1748E
27	Thyene imperialis (Rossi, 1846)	292	7 0 9+	Botanical garden, University of the Punjab,Lahore Jallo Park, Lahore Jinnah Garden, Lahore Pai Forest, Sakrand Lal Suhanra National Park, Bhawalpur Kallar Kahar, Chakwal	31.5004N, 74.3008E 31.5755N, 74.4943E 33.5723N,73.1748E 26.0593N, 68.1346E 29.3170N, 71.9045E 32.7760N, 72.7008E
28	Titanattus cretatus Chickering, 1946	2	9	Dera Bakha near Sutlej river, Bahawalpur	29.2641N, 71.5134E

DISCUSSION

The salticid fauna of Pakistan is rich like neighboring countries. According to Worldwide Database of Jumping Spiders (Metzner, 2017), 319 salticids from India, 498 from China, nine from Bangladesh and 106 from Sri Lanka have been recorded. However, in Pakistan serious efforts are required to explore the diversity of salticids. Few species of salticids are recorded from Pakistan which reflects the incomplete indigenous fauna. Majority of presently explored salticids were not reported earlier from Pakistan as previous work mentioned in the introduction. Most of the efforts done before this study focused on the spider fauna generally and not the salticids specifically. This is a preliminary survey on salticids and more extensive survey and collection of this diverse family in the Pakistan is required. It is expected that further studies will explore more and some interesting salticid species from Pakistan.

ACKNOWLEDGMENT

We are indebted especially to Dr. Riffat Sultana, Department of Zoology, University of Sindh Jamshoro and Mr. Salman Qureshi, Chief Warden, The Islamia University of Bahawalpur for facilitating the collections in these areas. We would also like to thank our colleagues for assistance in fieldwork.

REFERENCES

- Bao, Y. H., & Peng, X. J., 2002. Six New Species of Jumping Spiders (Araneae: Salticidae) from the Hui-Sun Experimental Forest Station, Taiwan. Zoological Stud., 41(4): 403-411.
- Bauer, T., Freudenschuss, M., Grabolle, A., Aaoao, O., Ooeaaiuphn, I., & Aoaaieea, A., 2015. *Plexippoides flavescens* (O. Pickard-Cambridge, 1872) and *Menemerus marginatus* (Kroneberg, 1875), new records for Pakistan (Aranei: Salticidae). *Arthropoda Sel.*, **24(1)**: 87-90.
- Ghazanfar, M., Hussain, M., Hashim, M., & Ul, A., 2016. Check list of spider (Araneae) fauna of Pakistan. *J. Entomol. Zool. Stud.*, **4(1)**: 245-256

- Maddison, W. P., Li, D., Bodner, M., Zhang, J., Xu, X., Liu, Q., & Liu, F., 2014. The deep phylogeny of jumping spiders (Araneae, Salticidae). *ZooKeys.* **440**: 57-87.
- Metzner, H., 2017. Jumping spiders (Arachnida: Araneae: Salticidae) of the world. Available at: http://www.jumping-spiders.com (accessed 20 August 2017).
- Mukhtar, M. K., Khan, S. Y., Jabeen, S., Tahir, H. M., Qadir, A., Ahmad, K. R., Butt, A., & Arshad, M.,2012. A Preliminary Checklist of the Spider Fauna of Sargodha (Punjab), Pakistan. *Pakistan J. Zool.*. **44(5)**:1245-1254.
- Perveen, F., & Khan, N., 2015. Checklist of the first recorded spider fauna in Sheringal, Khyber Pakhtunkhwa, Pakistan. *AdvEntomol.*, **3(03)**:94-100.
- Perveen, F., Jamal, A., Yasmin, S., & Khatak, K.U., 2012. Biodiversity of spidersfauna in the Frontier Region, Peshawar, Pakistan. *J. Entomol. Nematol.*, **4(3)**:22-33
- Parveen, R., Khan, A. A., Mushtaq, S., & Rana, S., 2007. A checklist of the spiders of the Punjab. *Pakistan J. Agri. Sci.*, **44(4)**: 625-626.
- Proszynski, J., 2016. Monograph of Salticidae (Araneae) of the World 1995-2015. Part II: Global Species Database of Salticidae (Araneae). Available at: http://www.peckhamia.com/salticidae/ (accessed 12 September 2017).
- Sial, N., Ruby, T., Malik, S.,& Mushtaq, S., 2012. A checklist of the spiders of Cholistan and neighbouring areas. *Pakistan J. Agri. Sci.*,**49(3)**:301-304.
- World Spider Catalog., 2017. World Spider Catalog. Natural History Museum Bern. Available at: http://wsc.nmbe.ch, version 18.5,(accessed 12 September 2017).