

Effect of selected organic amendments on germination and yield of *Abelmoschus esculentus* (L.) moench grown under Chromium stress

ALIA NASEEM, *SUMERA IQBAL & KHAJISTA JABEEN

Department of Botany, Lahore College for Women University, Lahore, Pakistan

ARTICLE INFORMATION	ABSTRACT
Received: 23-05-2018 Received in revised form: 29-07-2019 Accepted: 11-09-2019	Okra (<i>Abelmoschus esculentus</i> (L.) Moench) was used as test crop in a research trial to assess the impact of organic amendments (compost and farmyard manure) on germination and yield under application of chromium (Cr) stress. The no. of treatments applied was 10 while 3 replicates were used for each treatment. For inducing Cr stress, K ₂ Cr ₂ O ₇ (potassium dichromate) solution of 50 mg/Kg concentration was applied to the soil. Cr treated and untreated soil samples were amended with compost and farmyard manure at 2.5 % and 5 %. It was demonstrated that in okra when compared with control, the sugar content and germination parameters were affected adversely by Cr application. The treatments of compost at 5 percent and farmyard manure at 2.5 percent positively affected the yield in both Cr treated and untreated plants. In comparative analysis compost appeared as more beneficial for okra than farmyard manure. Overall, the application of compost at 5 percent was found to be effective in improving yield characters, chlorophyll and sugar contents of Cr affected okra plants.
*Corresponding Author: Sumera Iqbal: sumeraiqbal2@yahoo.com	Keywords: Organic amendments, Chromium stress, Okra, Compost, Farmyard manure.
Original Research Article	

INTRODUCTION

Okra (*Abelmoschus esculentus* (L.) Moench), widely cultivated as a vegetable throughout the world is a member of Malvaceae (Ahmed *et al.*, 2006). Poor disposal of tanning wastes (Shakir *et al.*, 2012; Rafique *et al.*, 2010), industrial waste water used for crop irrigation (Khan *et al.*, 2013) and human activities like electroplating and mining (Oliveira, 2012; Singh *et al.*, 2002) has significantly enhanced the chromium concentration in soil. For agronomic plants Cr is toxic at about 0.5 to 5.0 mg m L⁻¹ in nutrient solution and 5 to 100 mg kg⁻¹ of available Cr in soil (Hossner *et al.*, 1998).

The organic amendments are defined as almost decomposed plant or animal materials which are capable of improving the physical, chemical and biological characteristics of soil after its addition (Bouajila and Sanaa, 2011; Naramabuye, 2004). Considered as a conventional approach, the composting procedure requires microbial action in aerobic environment for decaying and stabilizing the organic material originated from living sources (Smith & Collins, 2007). Farmyard manure is a decomposed combination of farm animal excreta with litter and straw (Sabir *et al.*, 2015).

The present research was conducted to analyze and compare the impact of various organic amendments (farmyard manure & compost) on

germination, yield and biochemical parameters of Cr stressed okra plants.

MATERIALS AND METHODS

Experimental site and test crop

The experiment was carried out in Botanical garden of Lahore College for Women University, Lahore in 2016 and okra (*Abelmoschus esculentus* (L.) Moench cv. Arka Anamika) seed was provided by Punjab Seed Corporation Lahore.

Soil survey and seeds sterilization

The soil samples were tested by measuring their pH and EC (electrical conductivity). The okra seeds were immersed in 2 percent MgCl₂ (Mercuric chloride) soln. for 10 minutes and washed 5 times thereafter.

Experimental set up

Using randomized complete block design (RCBD), the experimental set up was organized with thirty pots each containing seven kilograms soil and 10 okra seeds. Total no. of treatments applied was 10 while 3 replicates were used for each

treatment. The soil was treated with $K_2Cr_2O_7$ solution of 50 mg/Kg concentration for inducing Cr stress. Cr treated and untreated soil samples were amended with compost and farmyard manure at 2.5 % and 5 %. In control (T1) samples, the stress and amendments were not applied.

For the measurement of germination percentage and seedling vigor index (SVI), analyses was carried out in line with the procedure of Close & Wilson, (2002) and Abdul Baki & Anderson, (1973), respectively. The samples of plants were selected at vegetative and reproductive phases and kept under the freezing conditions (at -35°C). For analyses of chlorophyll, proline and sugar content present in the leaf standard method according to Arnon, (1949), Dubois *et al.*, (1956) and Bates *et al.*, (1973) were used, respectively.

Estimation of yield

After harvest, the yield attributes of okra plants such as fresh weight of each fruit (g) and no. of fruits per plant were recorded for the yield estimation.

Statistical assessment

The experimental data was statistically analyzed by Analysis of Variance and Duncan's Multiple Range Test (Duncan, 1957) with COSTAT (Steel and Torrie, 1980) was used for comparing the values of group means.

RESULTS AND DISCUSSION

At first Cr affects the seed germination, so the tolerance level to this metal would be expressed by the seed germination potential under Cr application (Peralta *et al.*, 2001). As indicated by Fig. 1 & 2, chromium significantly ($p < 0.05$) affected the germination percentage and SVI, compared to control. The results regarding Cr induced decreased germination % are in lines with those of (Samantary & Deo, 2004) and (Ganesh *et al.*, 2006). The rise in protease functions (Zeid, 2001), restriction of amylase reactions and conduction of sugars to embryonic axis caused by Cr stress are considered as the probable reasons for decreased seed germination (Chidambaram *et al.*, 2006; Zeid, 2001). According to results, in Cr treated plants, an insignificant effect of organic amendments on germination parameters was observed relative to unamended Cr affected plants.

The results depicted that in comparison to control, chlorophyll content was insignificantly affected in Cr stressed plants at the vegetative

phase. According to Fig. 3, 4 & 5, compost applied at both levels and 2.5 % farmyard manure elevated the concentrations of Chl a, Chl b and total chlorophyll at vegetative phase of Cr treated plants.

In citrus (Wu & Xia, 2006), lettuce (Ali *et al.*, 2007) and ginger (Ahmad *et al.*, 2009), the addition of compost raised the levels of chlorophyll. According to (Anburani & Manivannan, 2002), the rise in chlorophyll content could be due to the nutrients supplied by farmyard manure. Similar results were observed by Nehra *et al.*, (2001) and Sanwal *et al.*, (2007). In wheat, the organic manure was involved in improving capability of plants to cope with stress and development of chlorophyll (Jan *et al.*, 2014).

According to the results of present study, a pronounced rise in proline content of Cr affected plants was found at both vegetative and reproductive phases, when compared to control. Regarding proline content, similar results were observed by Yilmaz & Parlak, (2011), Nataraj *et al.*, (2009) and Nataraj & Paramar, (2008). Zengin *et al.*, (2005) reported a marked elevation in proline level of bean leaves when subjected to heavy metal (Mercury, cadmium, copper and lead) stress for a period of 10 days. The excess of proline is considered as an adaptive response of plants for the alleviation of adverse effects caused by Cr stress (Rai *et al.*, 2004; Rout & Shaw, 1998). As shown in Fig. 6, at vegetative phase, Cr affected plants responded insignificantly to amendments with respect to proline content. However, reduction in proline content was registered in Cr affected plants at reproductive stage (Fig. 7) by compost application.

Considering sugar content, both the vegetative and reproductive phases of okra were adversely affected by Cr stress relative to control. Similar decrease in sugar content was observed in water lilies (Choo *et al.*, 2006) and sugarcane (Jain *et al.*, 2000). This stress induced decline in sugar could be due to the disturbances in photosynthetic reactions (Sundarmoorthy *et al.*, 2015). Compost and farmyard manure effectively raised the sugar concentrations in Cr affected plants at the vegetative (Fig. 8) and reproductive phases (Fig. 9).

In comparison to control, the influence of Cr was insignificant on the yield attributes of okra. In Cr treated and untreated plants, the treatments with 5 % compost and 2.5 % farmyard manure have played a significant role in elevating yield (Fig. 10) and weight per fruit (g) relative to control (Fig. 11). In okra, the yield (Akanbi, 2002) and soil nutrient status (Haynes and Naidu, 1998; Garg and Bahla, 2008; El-Magd *et al.*, 2006; Akanbi *et al.*, 2000) could be improved by the addition of different rates

of organic amendments. The enhanced phytoavailability of macronutrients as well as the enrichment of physical attributes of soil was observed after the incorporation of farmyard manure (Bhattacharyya *et al.*, 2008).




Fig. 1: Effect of chromium stress and application of organic amendments (compost and FYM) on germination percentage (%) of okra.




Fig. 2: Effect of chromium stress and application of organic amendments (compost and FYM) on seedling vigor index (SVI) of okra.




Fig. 3: Effect of chromium stress and application of organic amendments (compost and FYM) on chlorophyll a content (mg/g) of okra leaves at vegetative stage.




Fig. 4: Effect of chromium stress and application of organic amendments (compost and FYM) on chlorophyll b content (mg/g) of okra leaves at vegetative stage.




Fig. 5: Effect of chromium stress and application of organic amendments (compost and FYM) on total chlorophyll content (mg /g) of okra eaves at vegetative stage.




Fig. 7: Effect of chromium stress and application of organic amendments (compost and FYM) on proline content ((µM /g) of okra leaves at reproductive stage.




Fig. 6: Effect of chromium stress and application of organic amendments (compost and FYM) on proline content (µM /g) of okra leaves at vegetative stage.




Fig. 8: Effect of chromium stress and application of organic amendments (compost and FYM) on sugar content (mg /g) of okra leaves at vegetative stage.




Fig. 9: Effect of chromium stress and application of organic amendments (compost and FYM) on sugar content (mg /g) of okra leaves at reproductive stage.




Fig. 11: Effect of chromium stress and application of organic amendments (compost and FYM) on fresh weight of fruit (g) of okra.




Fig. 10: Effect of chromium stress and application of organic amendments (compost and FYM) on yield / plant (g) of okra.

CONCLUSION

The results of study revealed a pronounced reduction in seed germination percentage, seedling vigor index and sugar content of okra plants after Cr application. The treatments of compost at 5 percent and farmyard manure at 2.5 percent positively affected the yield in both Cr treated and untreated plants. In comparative analysis compost appeared as more beneficial for okra than farmyard manure. Overall, the application of compost at 5 percent was found to be effective in improving yield characters, chlorophyll and sugar contents of Cr affected okra plants.

REFERENCES

Abdul-Baki, A. A. & Anderson, J. D., 1973. Vigor determination in soybean seed by multiple criteria. *Crop Sci.*, 13(6): 630-633.

Ahmed, B. I., Abdulhameed, A., Yusuf, S. R. & Aliyu, M., 2006. Comparative study of the defoliatory activities of *Podagrica sjotedi* and *P. uniforma* (Coleoptera: Chrysomelidae) on two intercropped okra varieties in Bauchi State Nigeria Savannah. *J. Agric.*, 1(1): 12-14.

Ahmad, R., Azeen, M. & Ahmed, N., 2009. Productivity of ginger (*Zingiber officinalis*) by amendment of vermicompost and

- biogas slurry in saline soil. *Pak. J. Bot.*, 41: 3107-3116.
- Akanbi, W. B., 2002. Growth, nutrient uptake and yield of maize and okra as influenced by compost and nitrogen fertilizer under different cropping systems. Ph.D. Thesis. University of Ibadan, Nigeria. 228 pp.
- Akanbi, W. B., Adediran, J. A., Togun, A. O. & Sobulo, R. A., 2000. Effect of organic-base fertilizer on the growth, yield and storage life of tomato (*Lycopersicon esculentum* Mill). *Biosci. Res. Commun.*, 12(4): 439-444.
- Ali, M., Griffiths, J., Williams, P. & Jones, L., 2007. Evaluating the growth characteristics of lettuce in vermicompost and green waste compost. *Eur. J. Soil Biol.*, 43: 316-319.
- Anburani, A. & Manivannan, K., 2002. Effect of integrated nutrient management on growth in brinjal (*Solanum melongena* L. cv. Annamalai). *South Indian Hort.*, 50(4-6): 377-386.
- Arnon, D. I., 1949. Copper enzymes in isolated chloroplasts, polyphenol oxidase in *Beta vulgaris* L. *J. Plant Physiol.*, 24(1): 1-15.
- Bates, L. S., Waldren, R. P. & Teare, I. D., 1973. Rapid determination of free proline for water stress studies. *Plant Soil*, 39(1): 205-207.
- Bhattacharyya, R., Kundu, S., Prakash, V. & Gupta, H. S., 2008. Sustainability under combined application of mineral and organic fertilizers in a rainfed soybean-wheat system of the Indian Himalayas. *Eur. J. Agron.*, 28(1): 33-46.
- Bouajila, K. & Sanaa, M., 2011. Effects of organic amendments on soil physico-chemical and biological properties. *J. Mater. Environ. Sci.*, 2(S1): 485-490.
- Chidambaram, AL.A., Murugan, A., Ganesh, K. S. & Sundaramoorthy, P., 2006. Effect of chromium on growth and cell division of blackgram (*Vigna mungo* L. Hepper). *Plant Arch.*, 6(2): 763-766.
- Choo, T. P., Lee, C. K., Low, K. S. & Hishamuddin, O., 2006. Accumulation of chromium (VI) from aqueous solutions using water lilies (*Nymphaea spontanea*). *Chemosphere*, 62: 961-967.
- Close, D. C. & Wilson, S. J., 2002. Provenance effects on pregermination treatments for *Eucalyptus regnans* and *Eucalyptus delegatensis* seed. *Forest Ecol. Manag.*, 170: 299-305.
- DuBois, M., Gilles, K. A., Hamilton, J. K., Rebers, P. A. & Smith, F., 1956. Colorimetric method for determination of sugars and related substances. *Anal. Chem.*, 28: 350-356.
- Duncan, B. D., 1957. Multiple range tests for correlated and heteroscedastic means. *Biometrics*, 13(2): 359-364.
- El-Magd, A., El-Bassiony, M. & Fawzy, Z. F., 2006. Effect of organic manure with or without chemical fertilizers on growth, yield and quality of some varieties of broccoli plants. *J. Appl. Sci. Res.*, 2(10): 791-798.
- Ganesh, S. K., Sundaramoorthy, P. & Chidambaram, AL. A., 2006. Chromium toxicity effect on blackgram, soybean and paddy. *Pollut. Res.*, 25: 257-261.
- Garg, S. & Bahla, G. S., 2008. Phosphorus availability to maize as influenced by organic manures and fertilizer P associated phosphatase activity in soils. *Bioresour. Technol.*, 99(13): 5773-5777.
- Haynes, R. J. & Naidu, R., 1998. Influence of lime, fertilizer and manure applications on soil organic matter content and soil physical conditions, a review. *Nutr. Cycl. Agroecosys.*, 51(2): 123-137.
- Hossner, L. R., Loeppert, R. H., Newton, R. J. & Szaniszlo, P. J. 1998. Phytoaccumulation of chromium, uranium, and plutonium in plant systems (No. ANRCP—1998-3). Amarillo National Resource Center for Plutonium, TX (United States).
- Jain, R., Srivastava, S. & Madan, V. K., 2000. Influence of chromium on growth and cell division of sugarcane. *Indian J. Plant Physiol.*, 5(3): 228-231.
- Jan, K., Rather, A. M., Boswal, M. V. & Ganie, A. H., 2014. Effect of biofertilizer and organic fertilizer on morphophysiological parameters associated with grain yield with emphasis for further improvement in wheat yield production (Bread wheat=*Triticum aestivum* L.). *Int. J. Agric. Crop Sci.*, 7(4): 178-184.
- Khan, M. Y., Asghar, H. N., Jamshaid, M. U., Akhtar, M. J. & Zahir, Z. A., 2013. Effect of microbial inoculation on wheat growth and phytostabilization of chromium contaminated soil. *Pak. J. Bot.*, 45: 27-34.
- Naramabuye, F. X., 2004. Use of organic amendments as ameliorants for soil acidity in laboratory and fields experiments. Ph.D. Thesis. University of Kwazulu-Natal.
- Nataraj, M., Parmar, A. & Kher, M. M., 2009. Impact of chromium during early germination of fenugreek in interaction with copper, zinc, lead and cadmium. *PRAJÑA – J. Pure Appl. Sci.*, 17: 40-47.

- Nataraj, M. & Parmar, S., 2008. Biochemical response during the germination of raya and fenugreek seeds under heavy metal stress. *J. Cell Tissue Res.*, 8: 1589-1594.
- Nehra, A. S., Hooda, I. S. & Singh, K. P., 2001. Effect of integrated nutrient management on growth and yield of wheat (*Triticum aestivum* L.). *Indian J. Agron.*, 45: 112-117.
- Oliveira, H., 2012. Chromium as an environmental pollutant: insights on induced plant toxicity. *J. Bot.*, doi: 10.1155/2012/375843.
- Peralta, J. R., Gardea-Torresdey, J. L., Tiemann, K. J., Gomez, E., Arteaga, S., Rascon, E. & Parsons, J. G., 2001. Uptake and effects of five heavy metals on seed germination and plant growth in alfalfa (*Medicago sativa*) L. *Bull. Environ. Contam. Toxicol.*, 66(6): 727-734.
- Rafique, U., Ashraf, A., Khan, A. K., Nasreen, S. & Rashid, R., 2010. Toxic chromium from tanneries pollute water resources and soils of Sialkot (Pakistan). *J. Chem. Soc. Pak.*, 32: 644-649.
- Rai, V. P., Vajpayee, P., Singh, S. N. & Mehrotra, S., 2004. Effect of chromium accumulation on photosynthetic pigments, oxidative stress defense system, nitrate reduction, proline level and eugenol content of *Ocimum tenuiflorum* L. *Plant Sci.*, 167(5): 1159-1169.
- Rout, N. P. & Shaw, B. P., 1998. Salinity tolerance in aquatic macrophytes: probable role of proline, the enzymes involved in its synthesis and C₄ type of metabolism. *Plant Sci.*, 136(2): 121-130.
- Sabir, M., Zia-ur-Rehman, M., Hakeem, K. R. & Saifullah., 2015. Phytoremediation of metal-contaminated soils using organic amendments: Prospects and Challenges. Chapter 17. *Soil Remediation and Plants*, Academic Press, pp. 503-523.
- Samantary, S. & Deo, B., 2004. Studies on chromium toxicity in mung bean (*Vigna radiata* L.). *Adv. Plant Sci.*, 17: 189-194.
- Sanwal, S. K., Lakminarayana, K., Yadav, R. K., Rai, N., Yadav, D. S. & Mousumi, B., 2007. Effect of organic manures on soil fertility, growth, physiology, yield and quality of turmeric. *Indian. J. Hortic.*, 64(4): 444-449.
- Shakir, L., Ejaz, S., Ashraf, M., Qureshi, N. A., Anjum, A. A., Iltaf, I. & Javeed, A., 2012. Ecotoxicological risks associated with tannery effluent wastewater. *Environ. Toxicol. Pharmacol.*, 34: 180-191.
- Singh, D. B., Varma, S. & Mishra, S. N., 2002. Putrescine effect to nitrate reductase activity, organic nitrogen, protein, and growth in heavy metal and salinity stressed mustard seedlings. *Biol. Plant.*, 45(4): 605-608.
- Smith, J. L. & Collins, H. P., 2007. Composting. In: *Soil Microbiology, Ecology, and Biochemistry*. 3rd edition. ed. Paul, E. A. Academic Press. Burlington. pp. 483-486.
- Steel, R. G. D. & Torrie, J. H., 1980. *Principles and Procedures of Statistics*. A biometrical approach. 2nd Ed. McGraw Hill Book Company. Tokyo. Japan.
- Sundarmoorthy, P., Sankarganesh, K., Selvaraj, M., Baskaran, L. & Chidambaram, AL.A., 2015. Chromium induced changes in soybean (*Glycine max* L.) metabolism. *World Sci. News*, 16: 145-178.
- Wu, Q. S. & Xia, R. X., 2006. Arbuscular mycorrhizal fungi influence growth, osmotic adjustment and photosynthesis of citrus under well-watered and water stress conditions. *J. Plant Physiol.*, 163: 417-425.
- Yilmaz, D. D. & Parlak, K. U., 2011. Changes in proline accumulation and antioxidative enzyme activities in *Groenlandia densa* under cadmium stress. *Ecol. Indic.*, 11: 417-423.
- Zeid, I. M., 2001. Responses of *Phaseolus vulgaris* to chromium and cobalt treatments. *Biol. Plant.*, 44(1): 111-115.
- Zengin, F. K. & Munzuroglu, O., 2005. Effects of some heavy metals on content of chlorophyll, proline and some antioxidant chemicals in bean (*Phaseolus vulgaris* L.) seedlings. *Acta Biol. Cracov. Ser. Bot.*, 47(2): 157-164.