



Description of *Paratylenchus manilkarii* n. sp., *P. sindhicus* n. sp., and Observation of *Pratylenchus kralli* Ryss, 1982 from Chikoo (*Manilkara zapota* L. van Royen) Orchards of Balochistan and Sindh, Pakistan

Erum Iqbal*, Nasir Mehmood, Nasira Kazi and Shahina Fayyaz

National Nematological Research Centre, University of Karachi, Karachi, Pakistan

ABSTRACT

During the surveys of chikoo (*Manilkara zapota* L. van Royen) plantation at Hub, Balochistan and Gadab, Sindh a number of plant and soil nematodes were encountered. Soil sample analysis revealed two and a known plant parasitic nematode species viz., *Paratylenchus manilkarii* n. sp., *P. sindhicus* n. sp., and *Pratylenchus kralli* Ryss, 1982 as a new record species. *P. manilkarii* n. sp., is characterized by the lateral field with four incisures; stylet 28-29 μ m long; vulval lips protruding with vulval flap; lip region rounded or truncated without submedian lobes, tail ventrally curved with pointed or acute terminus. *P. sindhicus* n. sp., is characterized by stylet 20-21 μ m long; vulval lips not protruding with advulval flap; female head with small submedian lobes; tail ventrally curved with finely to broadly rounded terminus. Detailed taxonomical studies of new species have been incorporated herein with measurements, descriptions, illustrations and microphotographs along with brief description, measurements and microphotographs of the new record species.

Article Information

Received 15 March 2019

Revised 02 April 2019

Accepted 12 May 2019

Available online 22 November 2019

Authors' Contribution

EI and NK identified the specimens and prepared manuscript. NM did survey, processed samples and assisted in drawing. SF supervised the research and reviewed the manuscript.

Key words

Description, new species, *Paratylenchus*, chikoo, Pakistan

INTRODUCTION

Chikoo is called as Sapota in English. It is also called as nose berry, mud apples, Sapodilla and Sapodilla plum. It is a brown fruit with a rough skin and sweet brown pulp inside. Each fruit may have 4 or 5 black seeds. The seeds easily sprout into plants and grow into trees in suitable warm climates. The tree can only survive in warm, typically tropical environments, dies easily when temperature drops below freezing. From germination, the sapodilla tree will usually take from five to eight years to bear fruit. The sapodilla trees yield fruit twice a year, though flowering may continue year round.

Manilkara zapota, commonly known as the sapodilla (a long-lived, evergreen tree) is native to southern Mexico, Central America and the Caribbean. It was introduced to the Philippines during Spanish colonization. It is grown in large quantities in Pakistan, India, Thailand, Malaysia, Cambodia, Indonesia, Vietnam, Bangladesh and Mexico. In Pakistan sapodilla is cultivated in Karachi, Hyderabad and Gadab, Sind and in Hub, Lasbella district, Balochistan. It is

cultivated on about 564 hectares with production at 2018 metric tons. The plant prefers warm and moist weather and can grow both in dry and humid areas. The coastal climate is best suited for its cultivation (Anon, 2016). Investigations of plant parasitic and soil nematodes in Pakistan have been carried out by several scientists and researchers on various agricultural crops including cereals, fruit and vegetables (Maqbool and Shahina, 2001; Zarina and Shahina, 2013).

In Pakistan the first record of sapodilla nematodes was made by (Saeed and Ashrafi, 1973, 1974) Saeed completed his thesis entitled "Studies on some stylet bearing nematodes associated with sapodilla (*Achras zapota* L.), with special reference to *Hemicriconemoides mangiferae* (Siddiqi, 1961)", the first thesis on sapodilla in Pakistan. Among several species found associated with sapodilla (*Achras zapota* L. now *Manilkara zapota* L. van Royen), five species were reported as new records for Pakistan viz., *Helicotylenchus indicus*, *H. multicinctus*, *Hoplolaimus columbus*, *Psilenchus hilarus*, *Rotylenchulus reniformis* and *Tylenchorhynchus martini* (Saeed, 1974). Later on Saeed and his co-workers published some papers on nematodes of sapodilla during 1973-1987. As very little work has been done on this plant, the present study was conducted with the aim to further investigate the association of nematodes of sapodilla.

* Corresponding author: erum_i@yahoo.com
0030-9923/2020/0001-0001 \$ 9.00/0
Copyright 2020 Zoological Society of Pakistan

MATERIALS AND METHODS

During the nematological studies a total of 40 soil and root samples of sapodilla (*Manilkara zapota* L. van Royen) were taken; 25 samples from 8 locations of Hub, Lasbela, Balochistan and 15 samples collected from three locations of Gadap town, Sindh during October, 2018. The samples were placed in polythene bags and sealed tightly with a rubber band with relevant information including date of collection, locality, and crop detail was mentioned on each bag.

Soil and root samples were processed by the methods of Cobb (1918) and Baermann (1917) funnel method. After qualitative and quantitative analysis of nematodes process of killing, fixing, and slow dehydration were done for permanent slide mounts (Hooper *et al.*, 2005). Identification of nematodes was made through measurements given by De Man (1884) formula with an ocular micrometer under a compound microscope and identification was based on the systematics given by Siddiqi (2000). Illustrations were prepared with the help of a drawing tube attached to the compound microscope and photographed of nematodes were made with a Nikon DS, Film camera, attached to the same Nikon Eclipse E400 microscope.




Fig. 1. *Paratylenchus manilkarii* n.sp. A, whole female body; B, oesophageal region; C, vulval and reproductive structure; D-E, tail region.




Fig. 2. *Paratylenchus manilkarii* n.sp. A, whole female body; B, oesophageal region; C, vulval part; D, vulva with tail region (scale A= 10 µm; B-D=100 µm).

Paratylenchus manilkarii n.sp.

(Figs. 1, 2; Table 1)

Description

Female

Body slender, vermiform, ventrally arcuate when heat relaxed. Cuticular annulation distinct 1-1.5 µm wide at mid body. Lateral field marked by four incisures occupying about 1/3 of the body width. Female head rounded or truncate, sub median lobes indistinct, not offset from the body contour. Cephalic sclerotization weak. Stylet robust, averaging 28-29µm long, conus 75% of the total stylet length; knobs small 1.5-2.0 x 3.0-4 µm rounded, slightly posteriorly directed. Dorsal oesophagel gland orifice about 6µm behind the stylet base. Median pharyngeal bulb elongate, bearing distinct large valve. Isthmus slender encircled by nerve ring. Basal bulb pyriform 5-6 x 8-9 µm. Cardia small conoid, inconspicuous. Excretory pore anterior to the level of basal bulb or 67 µm from anterior end. Hemizonid just anterior to excretory pore. Gonad short, prodelphic, spermatheca short rounded empty. Vulva a transverse slit occupying about half of the body

width. Vulval lips slightly protruding with advulval flaps. Post uterine sac absent. Oocytes arranged in a single row. Anus indistinct. Tail slender, conoid, ventrally curved, annulated, gradually tapers to sub acute terminus.

Table I. Morphometric data for *Paratylenchus manilkarii* n. sp. (All measurements are in μm except L).

| Morphological characters | Holotype female | Paratype females (n=4) |
|--------------------------------|-----------------|----------------------------|
| L (mm) | 0.36 | 0.37 \pm 0.01(0.36-0.40) |
| a | 22.5 | 21.8 \pm 0.49 (21-22.5) |
| b | 4.5 | 4.42 \pm 0.08 (4.3-4.5) |
| c | 15 | 14.75 \pm 0.43 (14-15) |
| c' | 3.0 | 2.95 \pm 0.08 (2-3) |
| V | 78.6 | 78.32 \pm 0.50 (77.7-79) |
| Lip region height | 1.0 | 1.05 \pm 0.08 (1.0-1.2) |
| Lip region width | 4.0 | 3.87 \pm 0.21 (3.5-4) |
| Stylet | 28 | 28.3 \pm 0.41 (28-29) |
| Median bulb length | 14 | 14.5 \pm 0.5 (14-15) |
| Median bulb width | 5 | 5.5 \pm 0.5 (5-6) |
| Median bulb valve length | 5 | 4.87 \pm 0.21 (4.5-5) |
| Median bulb valve width | 3 | 2.95 \pm 0.08 (2-3) |
| Anterior end to excretory pore | 67 | 68.8 \pm 1.43 (67-70.5) |
| Pharynx length | 80 | 79.25 \pm 0.82 (78-80) |
| Maximum body width | 16 | 15.6 \pm 0.41 (15-16) |
| Vulval body diameter | 12 | 12 \pm 0.35 (11.5-12.5) |
| Vulva to tail terminus | 72.4 | 72.8 \pm 0.75 (72-74) |
| Anal body width | 8 | 8.2 \pm 0.25 (8-8.5) |
| Tail length | 24 | 23.6 \pm 0.41(23-24) |

Male

Not found.

Type host and locality

The type material was collected from rhizospheric soil samples of sapodilla (*Manilkara zapota*) from Hub, Lasbela, Balochistan, Pakistan.

Type specimens

Holotype and paratype females are deposited in the Nematode Collection of National Nematological Research Centre, University of Karachi, Karachi Pakistan.

Diagnosis and relationship

Paratylenchus manilkarii n. sp., is characterized by the lateral field with four incisures; stylet 28-29 μm long;

vulval lips protruding with vulval flap; lip region rounded or truncated without submedian lobes, continuous; excretory pore located at the anterior end of basal bulbs; gonad prodelphic with small spermatheca; tail ventrally curved with pointed or acute terminus.

According to the grouping scheme of *Paratylenchus* given by Ghaderi *et al.* (2014), this new species belongs to group 3 (stylet < 40 μm ; 4 lateral lines; advulval flaps present) and is most similar to *P. concavus* (Eroshenko, 1978). The new species differs from *P. concavus* in longer body (L= 360-365 vs 280-300) μm ; greater a, b and c values (a=21-22.5 vs 17-20; b= 4.5-4.7 vs 3.9-4.1; c=15-16 vs 9-10), slightly longer stylet (28-29 vs 25-28) μm . Cuticular annulation distinct vs fine cuticular annulations, ovary small vs long.




Fig. 3. *Paratylenchus sindhicus* n.sp. A, whole female body; B, whole male body; C, oesophageal region; D, vulval and reproductive structure; E, female tail region; F, male tail region.

Paratylenchus sindhicus n.sp.

(Figs. 3, 4; Table II)

Description

Female

Body ventrally curved to open C- shaped when heat relaxed. Cuticle finely annulated, annules 0.5-1 μm wide at mid body. Lip region narrow, truncated with

small submedian lobes; not offset from body. Cephalic sclerotization weak. Stylet well developed, almost straight with rounded to laterally directed knobs; the conus about three fourth of the total stylet length. Orifice of dorsal gland opening at 4-5 μ m posterior to stylet knobs. Pharynx criconematoid with well developed corpus with valvular apparatus large and muscular. Relatively short, narrow isthmus surrounded by nerve ring. Basal bulb pyriform, set off from intestine. Cardia small 2-3 μ m in length Excretory pore at mid or slightly posterior to mid of pharyngeal basal bulb. Gonad prodelphic, short and outstretched, in one female it is reflexed; spermatheca oval containing spheroid sperms. Vulva a broad transverse slit with distinct but relatively small advulval flaps. Post uterine sac absent. Post-vulval region tapering gradually. Anus distinct or vestigial. Tail ventrally curved tapering with conical pointed to slightly rounded tail tip.




Fig. 4. *Paratylenchus sindhicus* n.sp. A, whole female body; B, oesophageal region; C, vulva with tail region; D, male tail region. (Scale A= 10 μ m; B-D=100 μ m).

Male

Body slender, similar to female in general appearance but with smaller body size. Cephalic frame work less developed, narrower than female, continuous from body. Pharynx rudimentary, procorpus, metacarpus and basal bulb inconspicuous, Gonad about 65-70

μ m long; testis outstretched with small spermatozoa. Cloacal opening not elevated. Gubernaculum about 3 μ m long. Bursa absent. Tail terminus similar to that of female, tapering gradually to a bluntly pointed to rounded tip.

Table II. Morphometric data for *Paratylenchus sindhicus* n. sp. (All measurements are in μ m except L).

| Morphological characters | Holotype female | Paratypes | |
|----------------------------------|-----------------|-----------------------------|------------|
| | | Female (n=54) | Male (n=2) |
| L (mm) | 0.30 | 0.29-0.35 (0.32 \pm 0.02) | 0.27,0.32 |
| a | 25 | 23-28 (25 \pm 1.51) | 27,32 |
| b | 4.2 | 4.2-4.8 (4.4 \pm 0.18) | - |
| c | 21.4 | 18-21 (20.4 \pm 0.87) | 16.8,16.8 |
| c' | 2.3 | 2.0-2.5 (2.2 \pm 0.16) | 2,3 |
| V/T | 82 | 79.8-82.4 (81.0 \pm 0.84) | 33,20 |
| Lip region height | 2 | 1-2 (1.52 \pm 0.45) | 3,3 |
| Lip region width | 2 | 2-3 (2.5 \pm 0.42) | 2, 2 |
| Stylet | 21 | 20-21 (20.5 \pm 0.49) | 20,20 |
| Median bulb length | 13 | 13-16 (14.6 \pm 1.08) | - |
| Median bulb width | 5 | 5-7 (5.9 \pm 0.70) | - |
| Valvular apparatus length | 3 | 3-4 (3.5 \pm 0.45) | - |
| Valvular apparatus width | 2 | 2-3 (2.6 \pm 0.41) | - |
| Excretory pore from anterior end | 63 | 61-71 (63.6 \pm 2.5) | 60,56 |
| Pharynx length | 70 | 70-76 (71.4 \pm 2.57) | - |
| Maximum body width | 12 | 12-15 (12.6 \pm 0.98) | 10,10 |
| Vulval body width | 10 | 10-13 (11.6 \pm 1.04) | - |
| Vulva to tail terminus | 50 | 50-70 (57.8 \pm 6.88) | - |
| Anal/cloacal body width | 6 | 6-8 (6.8 \pm 0.72) | 6,8 |
| Tail length | 14 | 14-17 (15.5 \pm 0.93) | 18,16 |
| Spicule | - | - | 16,15 |
| Gubernaculum | - | - | 3,3 |
| St / L% | 7 | 5.8-6.7 (6.3 \pm 0.41) | 7.4,6.2 |

Type host and locality

The type material was collected from rhizospheric soil samples of saponilla (*Manilkara zapota*) from Gadap

town, Sindh, Pakistan

Type specimens

Holotype female, 54 paratype females and 2 paratype males were deposited in the Nematode Collection of National Nematological Research Centre, University of Karachi, Karachi, Pakistan.

Diagnosis and relationship

Paratylenchus sindhicus n. sp., is characterized by the combination of characters: stylet 20-21 μm long; vulval lips not protruding with advulval flap; female head with small submedian lobes; excretory pore located at the middle of basal bulb; gonad prodelphic with round to oval spermatheca filled with sperms; tail ventrally curved with finely to broadly rounded terminus.

According to the grouping scheme of *Paratylenchus* species given by Ghaderi *et al.* (2014), this new species belongs to group 5 (stylet $\leq 40\mu\text{m}$; lateral lines indistinct; advulval flaps present). There is only one species in this group, namely *Paratylenchus leiodermis* (Raski, 1975). The new species differs from it by smaller body length ($L=0.03\text{-}0.35$ vs $0.34\text{-}0.42\text{mm}$), in higher $st/L\%$ ratio ($5.8\text{-}6.7$ vs $4.5\text{-}5$); excretory pore located anteriorly ($61\text{-}71$ vs $70\text{-}84\mu\text{m}$); cuticular annulation (cuticle with distinct annules vs smooth or indistinct), head with small submedian lobes vs bluntly rounded head almost hemispherical.

Paratylenchus kralli Rass, 1982

(Figs. 5, 6; Table III)

Table III. Morphometric data of *Paratylenchus kralli* Ryss, 1983. (All measurements are in μm except L).

| Morphological characters | Female (n=8) | |
|--------------------------|-----------------|-----------|
| | Mean \pm SD | Range |
| L (mm) | 0.37 \pm 0.02 | 0.34-0.40 |
| a | 20.6 \pm 1.88 | 20.2-24.6 |
| b | 6.1 \pm 0.41 | 5.5-6.5 |
| b' | 4.2 \pm 0.25 | 4.0-4.7 |
| c | 18.6 \pm 1.0 | 17.5-20.3 |
| c' | 1.9 \pm 0.19 | 1.6-2.1 |
| V% | 75.4 \pm 0.93 | 74.5-77 |
| Stylet | 13.7 \pm 0.37 | 13-14 |
| Excretory pore | 63.2 \pm 3.1 | 59-68 |
| Oesophageal length | 87.7 \pm 2.94 | 85-93.6 |
| Tail length | 20.2 \pm 1.36 | 18.4-22 |
| Anal body width | 10.6 \pm 0.69 | 10-12 |
| Maximum body width | 17.3 \pm 1.37 | 16-20 |
| Post uterine sac | 23 \pm 1.63 | 20-25 |




Fig. 5. *Paratylenchus kralli* (Rass, 1982). A, whole female body; B, oesophageal region; C, vulval region; D-E, different female tail regions.




Fig. 6. *Paratylenchus kralli* (Rass, 1982). A, oesophageal region; B, vulval region; C-D, different tail regions. (Scale A-D = 100 μm).

*Brief description**Female*

Body slender and vermiform, slightly curved ventrally when killed. Body annuli 0.8-1.0 μm wide at mid body. The head region continuous with 3 lips. Stylet small robust and strongly sclerotized. Stylet conus 50% of the entire stylet length; stylet shaft tubular, basal knobs forwardly directed and wide.

Pharynx with rounded median bulb and rather small glandular lobe that overlaps the intestine ventrally. Isthmus slender and encircling by nerve ring. Hemizonid located just above the excretory pore at the level of pharyngeal-intestinal junction. Hemizonion not observed. Reproductive tract mono-prodelphic, germinative zone outstretched. Spermatheca rounded, filled with small spermatozoa. Post uterine branch long, more than corresponding body diameter. The vulva position is high (75-78%). The tail shape conoid with smooth terminus, bearing 17-23 annules. Phasmid pore like and located at one third from the tail tip.

Male

Not found.

Remarks

Specimens were collected from soil around rhizospheric region of sapodilla (*Manilkara zapota*) from Gadab town, Sindh. The present specimens agree well in all dimensions and morphological characters with those described by Ryss (1982).

Statement of conflict of interest

The authors declare no conflict of interest.

REFERENCES

- Anonymous, 2016. *Potential of chiku cultivation*. Newspaper Dawn.com
- Baermann, G., 1917. A simple method for attaching *Ankylostomum* (nematode) larvae to soil samples. *Dutch Med. Timesheet*, **57**: 131-137.
- Cobb, N.A., 1918. *Estimating the nema population of soil*. Agricultural Technical Circular, US Department of Agriculture, pp. 48.
- De Man, J.G., 1884. *The free living, soil and marine nematode fauna of Netherlands*. Leiden, pp. 1-206. <https://doi.org/10.5962/bhl.title.46884>
- Eroshenko, A.S., 1978. Pathogenic nematodes of pine plantations in the South of Sakhalin Island. *Fitogel'mintologicheskije issledovaniya*, **32**: 33-37 [in Russian].
- Ghaderi, R., Kashi, L. and Karegar, A., 2014. Contribution to the study of the genus *Paratylenchus* Micoletzky, 1922 sensu lato (Nematoda: Tylenchulidae). *Zootaxa*, **3841**: 151-187. <https://doi.org/10.11646/zootaxa.3841.2.1>
- Hooper, D.J., Hallmann, J. and Subbotin, S.A., 2005. Methods of extraction, processing and detection of plant and soil nematodes. In: *Plant parasitic nematodes in subtropical and tropical agriculture* (eds. M. Luc, R.A. Sikora and J. Bridge). 2nd Edition. CABI Publishing, Wallingford, UK, pp. 53-86. <https://doi.org/10.1079/9780851997278.0053>
- Maqbool, M.A. and Shahina, F., 2001. *Systematics and distribution: Biodiversity of nematode fauna in Pakistan*. National Nematological Research Centre, University of Karachi, Karachi, Pakistan, pp. 179.
- Raski, D.J., 1975. Revision of the genus *Paratylenchus* Micoletzky, 1922 and descriptions of species. Part I of 3 parts. *J. Nematol.*, **7**: 15-34.
- Ryss, A., 1982. New phytonematode species of the genus *Pratylenchus* in Estoniiti. *BioloSgia*, **31**: 22-29.
- Saeed, M. and Ashrafi, S.H., 1973. On the occurrence of some plant parasitic nematodes with special reference to new hosts in West Pakistan. *Pakistan J. scient. indust. Res.*, **16**: 128-129.
- Saeed, M., 1974. *Studies on some stylet-bearing nematodes associated with sapodilla (Achras zapota L.) with special reference to Hemicriconemoides mangiferae Siddiqi, 1961*. Ph. D. thesis, Department of Botany, University of Karachi, Karachi, Pakistan, pp. 137.
- Saeed, M., Khan, H.A. and Saeed, V.A., 1976. Plant parasitic nematodes in Sindh. *Iran J. Pl. Pathol.*, **12**: 3-12.
- Saeed, M. and Ghaffar, A., 1979. A survey of stylet bearing nematodes in Karachi. *Nematol. Mediterr.*, **7**: 127-128.
- Saeed, M. and Ghaffar, A., 1986. Seasonal population fluctuations of *Hemicriconemoides mangiferae* and other nematodes associated with sapodilla (*Achras zapota*) in Karachi. *Pakistan J. Nematol.*, **4**: 67-74.
- Saeed, M. and Ghaffar, A., 1987. Vertical and horizontal distribution of nematodes associated with sapodilla (*Achras zapota*) in Karachi. *Sarhad J. Agric.*, **3**: 333-340.
- Siddiqi, M.R., 1961. Studies on species of *Criconematoidea* (Nematoda: Tylenchida). *Proc. Helminthol. Soc. Washington*, **28**: 19-34.
- Siddiqi, M.R., 2000. *Tylenchida: Parasites of plants and insects*. 2nd Edition, CABI Publishing, Wallingford, UK, pp. 833. <https://doi.org/10.1079/9780851992020.0000>
- Zarina, B. and Shahina, F., 2013. *Annotated bibliography on nematology in Pakistan*. 2nd Edition. National Nematological Research Centre, University of Karachi, Karachi, Pakistan, pp. 850.