



Quality Evaluation of Reference Gene Expression on Different Tissues in Adults of Tropical Gar *Atractosteus tropicus*

Luis Daniel Jiménez Martínez¹, Vicente Morales García², Carlos Alfonso Frias Quintana³, Alejandra del Carmen Castillo Collado¹, Gloria Gertrudys Asencio Alcudia⁴, Carina Shianya Alvarez Villagomez⁴, Emyr Saul Peña Marín^{4,5}, Bartolo Concha Frias¹ and Carlos Alfonso Alvarez-Gonzalez^{4*}

¹Laboratorio de Biología Molecular, DAMJ-UJAT, Dirección: Carretera Estatal Libre Villahermosa-Comalcalco Km. 27+000 s/n Ranchería Ribera Alta, C.P. 86205, Jalpa de Méndez, Tabasco, Mexico.

²Laboratorio de Docencia, DAMC-UJAT, Dirección: Ranchería Sur cuarta sección Comalcalco, Tabasco, Mexico.

³Laboratorio de Investigación en Biotecnología Acuicola (LIBA), Tecnológico Nacional de México Campus Boca del Río (ITBoca). Boca del Río, Veracruz, Mexico.

⁴Laboratorio de Acuicultura Tropical, DACBIOL-UJAT, Carretera Villahermosa-Cárdenas Km 0.5, C.P.86139 Villahermosa, Tabasco, Mexico.

⁵Cátedra-CONACyT, CDMX, Mexico.

ABSTRACT

Tropical gar (*Atractosteus tropicus*) is an ancestral subtropical fish species in southeastern Mexico, which has great potential as a model species for physiological, biomedical and genomic studies. The quantification of gene expression through RT-qPCR is one of the most commonly used techniques, due to its precision, sensitivity and high performance, particularly in gene expression to compare between cells, tissues and organs; as well as different populations, stages of development, metabolism, among other conditions. This study analyzed the stability and normalization of six commonly used reference genes such as alpha elongation factor (*ef1- α*), beta-actin (*actb*), 18S ribosomal RNA (*18s rrna*), beta-2-microglobulin (*b2m*), tubulin alpha (*α -tub*) and glyceraldehyde 3-phosphate dehydrogenase (*gapdh*) in the different tissues of the intestine, muscle, gill, stomach, brain and liver in adult males of *A. tropicus* from reared in captivity through three BestKeeper, geNorm and NormFinder algorithms. Based on our results we can conclude that in the three BestKeeper, NormFinder and geNorm algorithms, the most stable genes are *ef1*, followed by *18s rrna* and *actb* where the gene stability will depend on specific tissue to analyze in tropical gar *A. tropicus* adults.

INTRODUCTION

Tropical gar (*Atractosteus tropicus*) is a tropical fish, characterized by being a very important carnivorous species in southeastern Mexico (Márquez-Couturier *et al.*, 2006). In the last 30 years, several studies have been carried out with the species, from the aspects of reproduction, larviculture, digestive physiology, feeding and nutrition. This has allowed advances in *A. tropicus* aquaculture techniques. Nutrition and energy requirements

studies let maximizing growth in less time by improving small-scale culture to seek profitability (Lee, 2002).

In this sense, *A. tropicus* is considered an ancestral species since it has kept its physiognomy intact by not suffering genetic duplication that separated them from the diversification of the Teleostean lineage 320-350 million years ago (Braasch *et al.*, 2016). That is why the interest of carrying out molecular biology works to understand and expand the knowledge of the different physiological and metabolic processes of the species.

The quantification of gene expression through RT-qPCR is one of the most commonly used techniques, due to its precision, sensitivity and high performance. There are two methods: absolute and relative quantification

* Corresponding author: alvarez_alfonso@hotmail.com
0030-9923/2021/0001-0001 \$ 9.00/0
Copyright 2021 Zoological Society of Pakistan

Article Information

Received 13 September 2020

Revised 03 November 2020

Accepted 20 November 2020

Available online 08 April 2021

Authors' Contribution

LDJM and CAFQ presented the concept. CAA-G, BCF, LDJM and CAFQ suggested the methodology. LDJM, GGAA, CSAV and ACCC curated data. ACCC, GGAA, VMC, LDJM, ESPM, CAA-G, BCF, CSAV and CAFQ wrote the manuscript. CAFQ, CAA-G and BCF supervised the study. CAFQ, ACCC, GGAA, ESPM, BCF, CSAV and CAA-G did formal analysis. CAA-G acquired funds for the study.

Key words

Atractosteus tropicus, Reference gene, Gene expression, Stability, Relative expression

(Schmittgen and Livak, 2008). The absolute quantification is determined by defining the exact number of copies by extrapolating the value from a standard or calibration curve while the relative quantification is not necessary a calibration curve, changes in the expression of a gene in response to a treatment are analyzed, comparing and relativizing another untreated control and normalizing with an endogenous control or reference gene whose expression does not change in response to any kind of treatment or condition (Wan *et al.*, 2010). A reference gene must be reliable to conduct a good relative expression study (Dundas and Ling, 2012). Ideal reference genes must be stably expressed in various cells, tissues and organs; different populations; different stages of development; different stages of the cell cycle; and different treatment circumstances (Li *et al.*, 2019). Currently, for the analysis of the stability of the reference gene there are different algorithms such as BestKeeper (Pfaffl *et al.*, 2004), geNorm (Vandesompele *et al.*, 2002) and NormFinder (Andersen *et al.*, 2004). These are tools using different algorithms where BestKeeper uses peer correlation analysis of each internal gene with an optimal normalization factor, NormFinder adjusts the data to a mathematical model, which allows to compare intra and intergroup variation and the calculation of expression stability and finally geNorm uses comparisons between pairs and the geometric average of the Cq values to establish the most stable genes (Wang *et al.*, 2012).

Currently, the alpha elongation factor (*efl- α*), beta-actin (*actb*), 18S ribosomal RNA (*18s rna*), beta 2 microglobulin (*b2m*), tubulin alpha (*α -tub*) and glyceraldehyde 3-phosphate dehydrogenase (*gapdh*) genes are considered the most stable and used as reference genes. The *efl- α* gene encodes a critical protein that acts in the formation of the cytoskeleton of cells (Ingerslev *et al.*, 2006); *actb* gene encodes an essential protein of the cytoskeleton that is an essential component of cells to maintain the necessary activities of life and the performance of an important role in cell secretion, phagocytosis, migration, cytoplasmic transmission and cytoplasmic segregation and recombination (Guo *et al.*, 2013). The *18s rna* gene is an essential component of eukaryotic cells and part of RNA ribosomes (Wang *et al.*, 2018). The *b2m* gene encodes a protein found on the surface of virtually all body cells and is released by cells into the blood, especially B cells (Winchester *et al.*, 2003; McCurley and Callard, 2008). The *α -tub* gene encodes a type of tubulin that exists as a dimer in the cell with tubulin and is involved in important physiological functions such as cell division and differentiation, substance transport, and signal transduction (Fortes *et al.*, 2016). Finally, the *gapdh* gene encodes a protein that acts in glycolysis for

energy creation (Nicholls *et al.*, 2012). These reference genes have been studied in marine fish such as *Paralichthys olivaceus* (Zheng and Sun, 2011), *Danio rerio* (Casadei *et al.*, 2011; McCurley and Callard, 2008), *Gasterosteus aculeatus* (Hibbeler *et al.*, 2008), *Solea senegalensis*, and *Hippoglossus hippoglossus* (Infante *et al.*, 2008; Øvergård *et al.*, 2010), *Salmo salar* (Kortner *et al.*, 2011), *Gadus morhua* (Olsvik *et al.*, 2008) and in freshwater fish such as *Siniperca chuatsi* (Zhou *et al.*, 2010) *Cyprinus carpio* (Tang *et al.*, 2012) *Oreochromis niloticus* (Yang *et al.*, 2013) *Ctenopharyngodon idella* (Su *et al.*, 2011) and *Oncorhynchus mykiss* (Salem *et al.*, 2015). The objective of the study was to determine the best reference genes in the expression in different tissues in *A. tropicus*, knowing the most stable gene is of vital importance for future relative expression studies in the areas of physiology, biomedicine and genomics in this species.

MATERIALS AND METHODS

Fish

A total of 20 male adults *A. tropicus* were obtained. Individuals (550-580 g and from 30 to 35 cm total length) from captivity at the facilities of the Tropical Aquaculture Laboratory of the DACBIOL-UJAT. The organisms were kept in polyethylene tanks measuring 1.94 m in diameter and 0.70 m in height. The diet provided was based on balanced feed for trout and contained 46% protein and 16% lipids with pellet sizes ranging from 5.5 to 9.0 mm (El Pedregal® Silver Cup, Toluca, Mexico).

Sampling, total RNA extraction and cDNA synthesis

Subsequently, individuals of *A. tropicus* were euthanized by thermal shock (-4 °C) according to the methodology of Matthews and Varga (2012) and dissected to obtain six tissues: intestine, muscle, gill, stomach, brain and liver. Extraction of RNA was performed from tissues pooled per replicate using the Trizol Reagent (Invitrogen, Carlsbad, USA) obtaining an integrity value of 9.7. cDNA was synthesized using one microgram of RNA and random primer with an iScript™ Select 170-8896 cDNA synthesis kit (Bio-Rad, Hercules, California, USA) following the manufacturer's instructions.

Quantitative polymerase chain reaction (qPCR)

Resulting cDNA from adult tissues were diluted in 200 μ L-distilled water. The qPCR reactions were performed in a 96-well CFX96 Real-Time System Thermal Cycle (Model C1000, California, USA) thermocycler. The reaction mixture included 10 μ L of Eva Green master mix, 2- μ L cDNA and 0.2 μ M of each primer. The specific primers used in this analysis are given alpha elongation factor (*efl- α*), beta-actina (*actb*)

(from sequences of *Atractosteus tropicus*), 18S ribosomal RNA (*18s rrna*) (from sequences of *Lepisosteus osseus*), beta-2-microglobulin (*b2m*), Tubulin alpha (*α-tub*) and glyceraldehyde 3-phosphate dehydrogenase (*gapdh*) (from sequences of *Lepisosteus oculatus*) (Table I). The thermal program included 2 min at 95 °C, followed by 38 cycles at 95 °C for 10 s, 60 °C for 30 s and extension at 70° C for 5 s. All reactions were performed in duplicate. A standard curve for each pair of primers was generated to estimate amplification efficiencies based on known amounts of cDNA (four serial dilutions corresponding to cDNA transcribed from 100 to 0.1 ng of total RNA). In the melting curve analysis, it was determined that the melting temperature peak varied between 81.5 and 83 °C, which corresponded to the product obtained by these primers. In addition, the absence of primer dimers and nonspecificities was confirmed.

Stability analysis of candidate reference genes

The original cycle thresholds (cq values) of the six candidate reference genes were obtained from Bio-Rad CFX-96 Manager, and the data was sorted by Excel to assess differences in the expression levels of six candidate reference genes. Subsequently, the stability analysis of the candidate reference genes was carried out using three software packages, including BestKeeper v1 (<https://www.gene-quantification.de/bestkeeper.html#download>), geNorm integrated in qBasePlus (<https://www.gene-quantification.de/hkg.html#genorm>) and NormFinder v20 (<https://www.moma.dk/normfinder-software/>). The results of the three software packages were compared and analyzed to determine which gene is the most appropriate reference gene according to the methodology of Li *et al.* (2019).

RESULTS

Analysis of specificity and reliability of qPCR primers

The primers for qPCR of the six candidate reference genes elongation factor alpha (*ef1-α*), beta-actin (*actb*), 18S ribosomal RNA (*18s rrna*), beta-2-microglobulin (*b2m*), Tubulin alpha (*α-tub*) and glyceraldehyde 3-phosphate dehydrogenase (*gapdh*) were subjected to PCR analysis specific ordinary before qPCR. The results showed that the six reference genes produced a single band (Fig. 1). The melting curves of the six candidate reference genes are all single peaks, indicating that the primers' specificity is right, and no primer-dimers are present (Fig. 2). Besides, standard curves were performed to calculate amplification efficiency (E) and correlation coefficients (R²) according to the Cq values of each candidate reference gene amplified by qPCR using the *A. tropicus* brain cDNA (Fig. 3).




Fig. 1. Agarose gel with PCR products of six candidate reference genes of *Atractosteus tropicus*. six candidate reference genes of *Atractosteus tropicus* for qPCR. (A) alfa elongation factor (*ef1-α*), (B) beta-actin (*actb*), (C) 18S ribosomal RNA (*18s rrna*), (D) beta-2-microglobulin (*b2m*), (E) Tubulin alpha (*α-tub*) and (F) Glyceraldehyde 3-phosphate dehydrogenase (*gapdh*).

Table I. Primer sequences and amplification parameters of six candidate reference genes of *Atractosteus tropicus* used in qPCR analysis.

| Gene | Primer sequences (5′–3′) | Product size (bp) | Amplification efficiency (%) | R ² | GenBank accession numbers |
|----------|--|-------------------|------------------------------|----------------|---------------------------|
| 18s rrna | F: GGTAACGGGGAATCAGGGTT R: TCCAATTACAGGGCCTCGAA | 156 | 100.18 | 0.9974 | AF188369.1 |
| gapdh | F: GGAATCAACGGATTTGGCCG R: TCACCTCCCCATGAAAACGG | 163 | 97.40 | 0.9972 | XM_006642348.2 |
| b2m | F: TTTACCTGGACTGGGGGCTA R: GCGAGGCGCCATAAATCAAC | 139 | 94.23 | 0.9924 | XM_015346206.1 |
| actb | F: GAGCTATGAGCTGCCTGATGG R: GTGGTCTCATGAATGCCACAGG | 119 | 97.10 | 0.9956 | KT351351.1 |
| α-tub | F: TCAGCCTCTTTTGTTCAGGCT R: GCATGTGATGAGCAAAGACCA | 181 | 94.35 | 0.9977 | XM_015359451.1 |
| ef1-α | F: CCTGCAGGACGTCTACAAGATCG R: GACCTCAGTGGTCACGTTGGA | 120 | 99.82 | 0.9891 | KT351350.1 |




Fig. 2. Melting curve analyses of six candidate reference genes of *Atractosteus tropicus* for qPCR. (A) alfa elongation factor (*efl-α*), (B) beta-actin (*actb*), (C) 18S ribosomal RNA (*18s rrna*), (D) beta-2-microglobulin (*b2m*), (E) Tubulin alpha (*α-tub*) and (F) Glyceraldehyde 3-phosphate dehydrogenase (*gapdh*).




Fig. 3. Standard curves for the determination of efficiency (E) and correlation coefficients (R²) of six candidate reference genes of *Atractosteus tropicus* for qPCR. (A) alfa elongation factor (*efl-α*), (B) beta-actin (*actb*), (C) 18S ribosomal RNA (*18s rrna*), (D) beta-2-microglobulin (*b2m*), (E) Tubulin alpha (*α-tub*) and (F) Glyceraldehyde 3-phosphate dehydrogenase (*gapdh*).

BestKeeper gene analysis

The stability analysis shows BestKeeper SD and CV values of the six genes of references in the different tissues analyzed (Table II). Likewise, the results obtained in this study are shown in order of the most stable at least stable *efl-α*, *actb*, *18s rrna*, *b2m*, *α-tub* and *gapdh* in the intestine, *efl-α*, *actb*, *α-tub*, *b2m*, *18s rrna* and *gapdh* in the case of muscle, *actb*, *efl-α*, *18s rrna*, *gapdh*, *b2m* and *α-tub* in gill, *18s rrna*, *efl-α*, *b2m*, *actb*, *gapdh* and *α-tub* in stomach, *efl-α*, *18s rrna*, *actb*, *b2m*, *α-tub* and *gapdh* in

brain and finally *efl-α*, *b2m*, *gapdh*, *actb*, *18s rrna* and *α-tub* liver.

Normfinder gene analysis

According to the results of the NormFinder analysis, they show both the stability value and the standard error of each candidate gene in the different tissues analyzed in adults of *A. tropicus* (Table III). Likewise, the results obtained in this study show that the most stable genes are *18s rrna*, *b2m* and *actb* in the case of intestine, *efl-α*, *b2m* and

Table II. BestKeeper analysis results of six candidate reference genes of *Atractosteus tropicus*.

| Gen name | Intestine | | | Muscle | | | Gill | | | Stomach | | | Brain | | | Liver | | |
|----------------|----------------|----------|------|----------------|----------|------|----------------|----------|------|----------------|----------|------|----------------|----------|------|----------------|----------|------|
| | SD (\pm cq) | CV (%cq) | r | SD (\pm cq) | CV (%cq) | r | SD (\pm cq) | CV (%cq) | r | SD (\pm cq) | CV (%cq) | r | SD (\pm cq) | CV (%cq) | r | SD (\pm cq) | CV (%cq) | r |
| 18s rrna | 0.12 | 0.53 | 0.99 | 1.25 | 4.82 | 0.84 | 0.40 | 1.71 | 0.54 | 0.03 | 0.13 | 0.96 | 0.14 | 0.57 | 0.93 | 0.31 | 1.56 | 1.00 |
| gapdh | 3.66 | 14.78 | 1.00 | 1.00 | 3.60 | 0.47 | 0.43 | 1.91 | 0.09 | 0.47 | 1.74 | 0.79 | 0.35 | 1.70 | 0.77 | 0.23 | 0.21 | 0.77 |
| b2m | 0.38 | 1.34 | 0.83 | 0.35 | 1.02 | 0.77 | 0.48 | 2.01 | 1.00 | 0.22 | 0.99 | 0.44 | 0.25 | 0.93 | 0.81 | 0.21 | 0.93 | 0.68 |
| actb | 0.12 | 0.36 | 0.98 | 0.10 | 0.31 | 0.81 | 0.01 | 0.26 | 0.82 | 0.41 | 1.47 | 1.00 | 0.20 | 0.86 | 0.74 | 0.29 | 1.12 | 1.00 |
| α - tub | 0.54 | 1.99 | 0.81 | 0.12 | 0.44 | 1.00 | 0.61 | 2.45 | 0.72 | 0.78 | 3.24 | 0.09 | 0.29 | 1.46 | 0.90 | 0.42 | 1.64 | 0.99 |
| ef1- α | 0.10 | 0.49 | 0.53 | 0.01 | 0.002 | 0.62 | 0.08 | 0.27 | 0.51 | 0.10 | 0.55 | 1.00 | 0.13 | 0.49 | 1.00 | 0.12 | 0.66 | 0.78 |

SD, Standard; CV, coefficient of variance, r, coefficient of correlation.

Table III. NormFinder analysis results of six candidate reference genes of *Atractosteus tropicus*.

| Gen name | Intestine | | Muscle | | Gill | | Stomach | | Brain | | Liver | |
|----------------|-----------------|----------------|-----------------|----------------|-----------------|----------------|-----------------|----------------|-----------------|----------------|-----------------|----------------|
| | Stability value | Standard error |
| 18s rrna | 0.004 | 0.0055 | 0.067 | 0.0060 | 0.706 | 0.0052 | 0.085 | 0.0042 | 0.689 | 0.0145 | 0.233 | 0.0086 |
| gapdh | 0.971 | 0.0086 | 1.545 | 0.0086 | 0.117 | 0.0040 | 0.613 | 0.0048 | 0.048 | 0.0067 | 0.011 | 0.0045 |
| b2m | 0.004 | 0.0065 | 0.014 | 0.0058 | 0.117 | 0.0042 | 0.504 | 0.0088 | 0.290 | 0.0157 | 0.014 | 0.0048 |
| actb | 0.152 | 0.0089 | 2.190 | 0.0080 | 0.753 | 0.0068 | 0.085 | 0.0043 | 0.336 | 0.0092 | 0.026 | 0.0080 |
| α - tub | 0.164 | 0.0293 | 0.163 | 0.0094 | 0.929 | 0.0059 | 1.421 | 0.0065 | 0.050 | 0.0086 | 0.051 | 0.0172 |
| ef1- α | 1.368 | 0.0178 | 0.014 | 0.0056 | 0.706 | 0.0172 | 0.085 | 0.0043 | 0.046 | 0.0055 | 0.266 | 0.00290 |

18s rrna in muscle, gapdh, b2m, 18s rrna in gill, 18s rrna, actb and ef1- α in stomach ef1- α , gapdh and α -tub in brain and finally gapdh, b2m and actb in liver.

geNorm gene analysis

The results of the geNorm analysis of each candidate reference gene in the different tissues analyzed in adults of *A. tropicus* show that the M values of the six candidate reference genes are all less than 1.5, indicating that the seven genes are suitable genes of reference (Fig. 4). The stability of the expression of the seven candidate reference genes in both is classified in descending order according to the principle that the lower the M value, the better the stability of the gene expression: 18s rrna, α -tub, actb, b2m, ef1- α , and gapdh in intestine, ef1- α , actb, 18s rrna, b2m, gapdh, α -tub in the case of muscle, actb, ef1- α , 18s rrna, gapdh, α -tub and b2m in gill, 18s rrna, ef1- α , gapdh, α -tub, b2m and actb in stomach, ef1- α , gapdh, actb, b2m, 18s rrna and tuba in brain and finally 18s rrna, ef1- α , gapdh, actb, α -tub and b2m in liver.

DISCUSSION

The qPCR is an effective method based on the polymerase chain reaction (PCR), which can amplify

and quantify DNA molecules or specific complementary DNA (cDNA). This method allows us to access reliable data and precise information on the genetic expression of the cells under study, and frequently it is combined with a retro-transcription reaction (RT-qPCR). This method has become a more suitable choice for performing a rapid and quantitative examination for specific gene expression (Wang and Zhang, 2012). Nevertheless, selecting a suitable reference gene is the precondition for analysis of the relative expression of a target gene in quantitative real-time PCR (Wang et al., 2018).

Considering the above-mentioned, the most important characteristic of any reference gene candidate is the stability of its expression, regardless of in which tissues, developmental restrictions or physiological states are expressed (Yang et al., 2013). Likewise, Schaeck et al. (2016) suggest using at least two reference genes for normalization and that these genes are previously validated using a larger set of reference genes (n=10) to identify the most stable genes for each tissue/ cell type and each experimental condition. In this sense, algorithms such as BestKeeper, NormFinder and GenNorm (Wang et al., 2012) are used to evaluate the stability of these genes. Thanks to these algorithms, the stability of the reference gene candidate can be identified and evaluated for the

correct normalization being therefore, an essential tool for an adequate selection. However, due to the characteristics of each of these platforms, it is necessary to select what gathers the requirements of the research work (De Spiegelaere *et al.*, 2015).




Fig. 4. geNorm analysis results of six candidate reference genes of *Atractosteus tropicus*.

Regarding our results, in the case of intestine and stomach the most stable gene was *18s rrna*, which correspond with the work done in species such as Nile tilapia (*Oreochromis niloticus*), carp (*Ctenopharyngodon idella*), Asian sea bass (*Lateolabrax maculatus*) and American catfish (*Ictalurus punctatus*) (Yang *et al.*, 2013; Su *et al.*, 2011; Wang *et al.*, 2018; Small *et al.*, 2008), presenting good stability in other tissues such as heart, liver, muscle, spleen, skin and kidney without being affected by any treatment.

It is important to mention that *18s rrna* is considered an appropriate gene in embryogenesis in Zebra fish (*Danio rerio*), fathead minnow (*Pimephales promelas*), Channel catfish (*Ictalurus punctatus*) and Atlantic salmon (*Salmo salar*) (McCurley and Callard, 2008; Filby and Tyler, 2007; Small *et al.*, 2008; Jorgensen *et al.*, 2006; Kortner *et al.*, 2011). Additionally, *18S rRNA* gene has been detected

in ancestral species such as longnose gar (*Lepissosteus osseus*), green sturgeon (*Acipenser medirostris*) in muscle and in bowfin (*Amia calva*) in fin (Krieger and Fuerst, 2002). Studies on the expression of growth hormone in larvae of alligator gar (*Atractosteus spatula*) used the *18s rrna* gene to normalize the relative expression (Cahu *et al.*, 2004; Revol *et al.* 2005; Panserat and Kaushik, 2010). It is important to highlight that for *A. tropicus* this gene can be used for nutrigenomic studies with tissues such as the intestine and stomach since it has good stability.

In the case of muscle, brain and liver, the most stable gene was *ef1-α* for *A. tropicus* adults. These results are consistent with those reported in species such as common carp (*Cyprinus carpio*), reporting greater stability in the liver, brain, hypothalamus, heart and kidneys (Tang *et al.*, 2012), in *O. niloticus* (Yang *et al.*, 2013) in muscle and heart. Similarly, *ef1-α* has been reported in common lancelet (*Branchiostoma lanceolatum*), Senegal sole (*Solea senegalensis*), Atlantic halibut (*Hippoglossus hippoglossus*), grass carp (*Ctenopharyngodon idella*), Korean rockfish (*Sebastes schlegeli*) (Carlos *et al.*, 2008; Infante *et al.*, 2008; Tang *et al.*, 2012; Wang and Zhang, 2012; Liman *et al.*, 2013), showing high stability in the different tissues analyzed in these species. On the other hand, *ef1-α* would be the ideal reference gene in studies involving the expression of genes involved in metabolism and neurofunctional (Diotel *et al.*, 2010; Zheng *et al.*, 2013), as well as in rainbow trout (*Oncorhynchus mykiss*) when analyzing the gene expression in liver and muscle tissues using different diets (Kolditz *et al.*, 2008). Finally, for *A. tropicus*, *EF1-α* was an excellent reference gene for analysis of relative gene expression of lipogenic genes during the initial ontogeny for *A. tropicus* (Jiménez-Martínez *et al.*, 2019) and in juveniles feed with diets supplemented with different concentrations of β -glucans (Nieves-Rodríguez *et al.*, 2018).

In gills, *actb* gene was highly stable in adults of *A. tropicus*, which are agreed with species such as Turbot (*Scophthalmus maximus*) (Dang and Sun, 2011) and bastard halibut (*Paralichthys olivaceus*) (Zheng and Sun, 2011), been the best reference gene not only for gill, also for liver, spleen, kidney, heart muscle, brain and intestine. Likewise, *actb* is considering the best reference gene in different tissues in teleosts (Deloffre *et al.*, 2012). It is important to point-out that gill tissue is the first line of defense against pathogens suspended in the aquatic environment, therefore, cells need to be fastly replicated (Hibbeler *et al.*, 2008). Finally, *ACTB* has been used as reference gene for toxicology, osmorrugulation and hypoxia studies (Wong *et al.*, 2001; Lin *et al.*, 2004; Bridle *et al.*, 2006).

In the case of *ef1-α* and *b2m* they were the least stable

genes in our study, however in bastard halibut (*Paralichthys olivaceus*) stands out, in which it mentions that *α -tub* is the best reference gene for spleen, heart, muscle and gill tissues (Zheng and Sun, 2011). While *b2m* alone in the geNorm program was one of the most stable in branchia *A. tropicus* similar to those occurring in larval development and tissues in the zebrafish *Danio rerio* under treatments such as different chemicals (McCurlley and Callard, 2008).

Based on BestKeeper, NormFinder, and geNorm analyses, the most stable genes for *A. tropicus* were *efl- α* , followed by *18s rrna* and *actb*, so they are adequate as reference genes and can be used individually; however, it is advisable to use any of them to quantify the relative gene expression in *A. tropicus*.

ACKNOWLEDGMENTS

This study was supported by the projects SAGARPA-2011-08-164673 and CB-2016-01-282765. The author thanks to the Consejo Nacional de Ciencia y Tecnología and Programa Institucional de Superación Académica for the scholarship awarded.

Statement of conflict of interest

The authors have declared no conflict of interest.

Ethical approval

All applicable international, national, and/or institutional guidelines for the care and use of animals were followed by the authors.

REFERENCES

- Andersen, C.L., Jensen, J.L. and Ørntoft, T.F., 2004. Normalization of real-time quantitative reverse transcription-PCR data: a model-based variance estimation approach to identify genes suited for normalization, applied to bladder and colon cancer data sets. *Cancer Res.*, **64**: 5245-525. <https://doi.org/10.1158/0008-5472.CAN-04-0496>
- Braasch, I., Gehrke, A.R., Smith, J.J., Kawasaki, K., Manousaki, T., Pasquier, J. and Catchen, J., 2016. The spotted gar genome illuminates vertebrate evolution and facilitates human-teleost comparisons. *Nat. Genet.*, **48**: 427-437. <https://doi.org/10.1038/ng.3526>
- Bridle, A.R., Morrison, R.N. and Nowak, B.F., 2006. The expression of immune-regulatory genes in rainbow trout, *Oncorhynchus mykiss*, during amoebic gill disease (AGD). *Fish Shellf. Immunol.*, **20**: 346-364. <https://doi.org/10.1016/j.fsi.2005.05.014>
- Cahu, C., Rønnestad, I., Grangier, V. and Infante, J.Z., 2004. Expression and activities of pancreatic enzymes in developing sea bass larvae (*Dicentrarchus labrax*) in relation to intact and hydrolyzed dietary protein; involvement of cholecystokinin. *Aquaculture*, **238**: 295-308. <https://doi.org/10.1016/j.aquaculture.2004.04.013>
- Carlos, I., Makoto, M., Esther, A., Michael, R. and Manuel, M., 2008. Selection of housekeeping genes for gene expression studies in larvae from flatfish using real-time PCR. *BMC Mol. Biol.*, **9**: 1-28. <https://doi.org/10.1186/1471-2199-9-28>
- Casadei, R., Pelleri, M.C., Vitale, L., Facchin, F., Lenzi, L., Canaider, S., Strippoli, P. and Frabetti, F., 2011. Identification of housekeeping genes suitable for gene expression analysis in the zebrafish. *Gene Expr. Patterns*, **11**: 271-276. <https://doi.org/10.1016/j.gep.2011.01.003>
- Dang, W. and Sun, L., 2011. Determination of internal controls for quantitative real time RT-PCR analysis of the effect of *Edwardsiella tarda* infection on gene expression in turbot (*Scophthalmus maximus*). *Fish Shellf. Immunol.*, **30**: 720-728. <https://doi.org/10.1016/j.fsi.2010.12.028>
- De Spiegelaere, W., Dern-Wieloch, J., Weigel, R., Schumacher, V., Schorle, H., Nettersheim, D., Bergmann, M., Brehm, R., Kliesch, S., Vandekerckhove, L. and Fink, C., 2015. Reference gene validation for RT-qPCR, a note on different available software packages. *PLoS One*, **10**: 122-515. <https://doi.org/10.1371/journal.pone.0122515>
- Deloffre, L.A., Andrade, A., Filipe, A.I. and Canario AV., 2012. Reference genes to quantify gene expression during oogenesis in a teleost fish. *Gene*, **506**: 69-75. <https://doi.org/10.1016/j.gene.2012.06.047>
- Diotel, N., Le Page, Y., Mouriec, K., Tong, S.K., Pellegrini, E., Vaillant, C. and Chung, B.C., 2010. Aromatase in the brain of teleost fish: Expression, regulation and putative functions. *Front. Neuroendocrinol.*, **31**: 172-192. <https://doi.org/10.1016/j.yfrne.2010.01.003>
- Dundas J. and Ling M., 2012. Reference genes for measuring mRNA expression. *Theory Biosci.*, **131**: 215-223. <https://doi.org/10.1007/s12064-012-0152-5>
- Filby A.L. and Tyler C.R., 2007. Appropriate 'housekeeping' genes for use in expression profiling the effects of environmental estrogens in fish. *BMC Mol. Biol.*, **8**: 1-10. <https://doi.org/10.1186/1471-2199-8-10>
- Fortes, M.A.S., Marzuca-Nassr, G.N., Vitzel, K.F., da Justa Pinheiro, C.H., Newsholme, P. and Curi, R., 2016. Housekeeping proteins: How useful are they

- in skeletal muscle diabetes studies and muscle hypertrophy models? *Anal. Biochem.*, **504**: 38-40. <https://doi.org/10.1016/j.ab.2016.03.023>
- Guo, C., Liu, S., Wang, J., Sun, M.Z. and Greenaway, F.T., 2013. ACTB in cancer. *Clin. Chim. Acta*, **417**: 39-44. <https://doi.org/10.1016/j.cca.2012.12.012>
- Hibbeler, S., Scharsack, J.P. and Becker, S., 2008. Housekeeping genes for quantitative expression studies in the three-spined stickleback *Gasterosteus aculeatus*. *BMC Mol. Biol.*, **9**: 9-18. <https://doi.org/10.1186/1471-2199-9-18>
- Infante, C., Matsuoka, M.P, Asensio, E., Cañavate, J.P., Reith, M. and Machado, M., 2008. Selection of housekeeping genes for gene expression studies in larvae from flatfish using real-time PCR. *BMC Mol. Biol.*, **9**: 18-28. <https://doi.org/10.1186/1471-2199-9-28>
- Ingerslev, H.C., Pettersen, E.F., Jakobsen, R.A., Petersen, C.B. and Wergeland, H.I., 2006. Expression profiling and validation of reference gene candidates in immune relevant tissues and cells from Atlantic salmon (*Salmo salar* L.). *Mol. Immunol.*, **43**: 1194-1201. <https://doi.org/10.1016/j.molimm.2005.07.009>
- Jiménez-Martínez, L.D., Álvarez-González, C.A., De la Cruz-Hernández, E., Tovar-Ramírez D., Galaviz, M.A., Camarillo-Coop, S., Martínez-García, R. and Peña E., 2019. Partial sequence characterization and ontogenetic expression of genes involved in lipid metabolism in the tropical gar (*Atractosteus tropicus*). *Aquact. Res.*, **50**: 162-172. <https://doi.org/10.1111/are.13879>
- Jorgensen, S.M., Kleveland, E.J., Grimholt, U. and Gjoen, T., 2006. Validation of reference genes for real-time polymerase chain reaction studies in Atlantic salmon. *Mar. Biotechnol.*, **8**: 398-408. <https://doi.org/10.1007/s10126-005-5164-4>
- Kolditz, C., Paboeuf, G., Borthaire, M., Esquerre, D., Sancristobal, M., Lefevre, F. and Medale, F., 2008. Changes induced by dietary energy intake and divergent selection for muscle fat content in rainbow trout (*Oncorhynchus mykiss*), assessed by transcriptome and proteome analysis of the liver. *BMC Genom.*, **9**: 506. <http://doi.org/10.1016/j.aquaculture.2009.05.013>
- Kortner, T.M., Valen, E.C., Kortner, H., Marjara, I.S., Krogdahl, Å. and Bakke, A.M., 2011. Candidate reference genes for quantitative real-time PCR (qPCR) assays during development of a diet-related enteropathy in Atlantic salmon (*Salmo salar* L.) and the potential pitfalls of uncritical use of normalization software tools. *Aquaculture*, **318**: 355-363. <https://doi.org/10.1016/j.aquaculture.2011.05.038>
- Krieger, J. and Fuerst, P., 2002. Evidence of multiple alleles of the nuclear 18S ribosomal RNA gene in sturgeon (Family: *Acipenseridae*). *J. appl. Ichthyol.*, **18**: 290-297. <https://doi.org/10.1046/j.1439-0426.2002.00377.x>
- Lee, S.M., 2002. Apparent digestibility coefficients of various feed ingredients for juvenile and grower rockfish (*Sebastes schlegeli*). *Aquaculture*, **207**: 79-95. [https://doi.org/10.1016/S0044-8486\(01\)00751-7](https://doi.org/10.1016/S0044-8486(01)00751-7)
- Li, J.Y, Chen, W.Z., Yang, S.H., Xu, C.L., Huang, X., Chen, C. and Xie H., 2019. Screening of reference genes in real-time PCR for *Radopholus similis*. *PeerJ.*, **7**: 6233-6253. <https://doi.org/10.7717/peerj.6253>
- Liman, M., Wenji, W., Conghui, L., Haiyang, Y., Zhigang, W., Xubo, W. and Quanqi, Z., 2013. Selection of reference genes for reverse transcription quantitative real-time PCR normalization in black rockfish (*Sebastes schlegeli*). *Mar. Gen.*, **11**: 67-73. <https://doi.org/10.1016/j.margen.2013.08.002>
- Lin, C., Tsai, R. and Lee, T., 2004. Expression and distribution of Na, K-ATPase in gill and kidney of the spotted green pufferfish, *Tetraodon nigroviridis*, in response to salinity challenge. *Comp. Biochem. Physiol.*, **138**: 287-295. <https://doi.org/10.1016/j.cbpb.2004.04.005>
- Márquez-Couturier, G., Álvarez-González, C.A., Contreras-Sánchez, W., Hernández-Vidal, U., Hernández-Franyutti, A., Mendoza-Alfaro, R. and Goytortua-Bores, E., 2006. Advances in food and nutrition alligator gar *Atractosteus tropicus*. In: *Advances in aquaculture nutrition* (eds. E. Cruz-Suarez, D. Ricque, M. Tapia, M. Nieto, D. Villareal, A. Puello and A. García). Monterrey, Nuevo León, México: Autonomous University of Nuevo Leon. pp. 446–523.
- Matthews, M. and Varga, Z., 2012. Anesthesia and euthanasia in zebrafish. *ILAR J.*, **53**: 192-204. <https://doi.org/10.1093/ilar.53.2.192>
- McCurlay, A.T. and Callard, G.V., 2008. Characterization of housekeeping genes in zebrafish: Male female differences and effects of tissue type, developmental stage and chemical treatment. *BMC Mol. Biol.*, **9**: 90-102. <https://doi.org/10.1186/1471-2199-9-102>
- Nicholls, C., Li, H. and Liu, J.P., 2012. GAPDH: A common enzyme with uncommon functions. *Clin. exp. Pharmacol. Physiol.*, **39**: 674-679. <https://doi.org/10.1111/j.1440-1681.2011.05599.x>
- Nieves-Rodríguez, K.N, Alvarez-González, C.A,

- Peña, E., Vega-Villasante, F., Martínez-García, R., Camarillo-Coop, S., Tovar-Ramírez, D., Guzmán-Villanueva, L.T., Andree, K.B. and Gisgert E., 2018. Effect of β -glucans in diets on growth, survival, digestive enzyme activity, and immune system and intestinal barrier gene expression for tropical gar (*Atractosteus tropicus*) Juveniles. *Fishes*, **3**: 1-13. <https://doi.org/10.3390/fishes3030027>
- Olsvik, P.A., Søfteland, L. and Lie, K.K., 2008. Selection of reference genes for qRT-PCR examination of wild populations of Atlantic cod *Gadus morhua*. *BMC Res. Notes*, **1**: 47-56. <https://doi.org/10.1186/1756-0500-1-47>
- Øvergård, A.C., Nerland, A.H. and Patel, S., 2010. Evaluation of potential reference genes for real time RT-PCR studies in Atlantic halibut (*Hippoglossus Hippoglossus* L.); during development, in tissues of healthy and NNV-injected fish, and in anterior kidney leucocytes. *BMC Mol. Biol.*, **11**: 36-51. <https://doi.org/10.1186/1471-2199-11-36>
- Panserat, S. and Kaushik, S., 2010. Regulation of gene expression by nutritional factors in fish. *Aquac. Res.*, **41**: 751-762. <https://doi.org/10.1111/j.1365-2109.2009.02173.x>
- Pfaffl, M.W., 2001. A new mathematical model for relative quantification in real-time RT-PCR. *Nucl. Acids Res.*, **29**: 45-50. <https://doi.org/10.1093/nar/29.9.e45>
- Pfaffl, M.W., Tichopad, A., Prgomet, C. and Neuvians T.P., 2004. Determination of stable housekeeping genes, differentially regulated target genes and sample integrity: BestKeeper-Excel-based tool using pair-wise correlations. *Biotechnol. Lett.*, **26**: 509-515. <https://doi.org/10.1023/B:BILE.0000019559.84305.47>
- Revol, A., Rodríguez, M.d.L.G., Montenegro, V.H., Aguilera, C., Saldaña, H.B. and Mendoza R., 2005. Cloning of the growth hormone cDNA of alligator gar *Atractosteus spatula* and its expression through larval development. *Comp. Biochem. Physiol.*, **140**: 423-429. <https://doi.org/10.1016/j.cbpb.2005.02.004>
- Salem, M., Paneru, B., Al-Tobasei, R., Abdouni, F., Thorgaard, G.H., Rexroad, C.E and Yao, J., 2015. Transcriptome assembly, gene annotation and tissue gene expression atlas of the rainbow trout. *PLoS One*, **10**: 1-27. <https://doi.org/10.1371/journal.pone.0121778>
- Schaeck, M., De Spiegelaere, W., De Craene, J., Van Den Broeck, W., De Spiegelaere, B., Burvenich, C. and Decostere, A., 2016. Laser capture microdissection of intestinal tissue from sea bass larvae using an optimized RNA integrity assay and validated reference genes. *Sci. Rep.*, **6**: 210-292. <https://doi.org/10.1038/srep21092>
- Schmittgen, T.D. and Livak, K.J., 2008. Analyzing real-time PCR data by the comparative CT method. *Nat. Prot.*, **3**: 1101-1108. <https://doi.org/10.1038/nprot.2008.73>
- Small, B.C., Murdock, C.A., Bilodeau-Bourgeois, A.L., Peterson, B.C. and Waldbieser, G.C., 2008. Stability of reference genes for real-time PCR analyses in channel catfish (*Ictalurus punctatus*) tissues under varying physiological conditions. *Comp. Biochem. Physiol.*, **151**: 296-304. <https://doi.org/10.1016/j.cbpb.2008.07.010>
- Su, J., Zhang, R., Dong, J. and Yang, C., 2011. Evaluation of internal control genes for qRT-PCR normalization in tissues and cell culture for antiviral studies of grass carp (*Ctenopharyngodon idella*). *Fish Shellf. Immunol.*, **30**: 830-835. <https://doi.org/10.1016/j.fsi.2011.01.006>
- Tang, Y.K., Yu, J.H., Xu, P., Li, J.L., Li, H.X. and Ren, H.T., 2012. Identification of housekeeping genes suitable for gene expression analysis in Jian carp (*Cyprinus carpio* var. jian). *Fish Shellf. Immunol.*, **33**: 775-779. <https://doi.org/10.1016/j.fsi.2012.06.027>
- Vandesompele, J., De Preter, K., Pattyn, F., Poppe, B., Van Roy, N., De Paepe, A. and Speleman, F., 2002. Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol.*, **3**: 1-12. <https://doi.org/10.1186/gb-2002-3-7-research0034>
- Wan, H., Zhao, Z., Qian, C., Sui, Y., Malik, A.A. and Chen, J., 2010. Selection of appropriate reference genes for gene expression studies by quantitative real-time polymerase chain reaction in cucumber. *Anal. Biochem.*, **399**: 257-261. <https://doi.org/10.1016/j.ab.2009.12.008>
- Wang, H., Wen, H., Li, Y., Zhang, K. and Liu Y., 2018. Evaluation of potential reference genes for quantitative RT-PCR analysis in spotted sea bass (*Lateolabrax maculatus*) under normal and salinity stress conditions. *PeerJ.*, **6**: 5631-5647. <https://doi.org/10.7717/peerj.5631>
- Wang, Q., Ishikawa, T., Michiue, T., Zhu, B.L., Guan, D.W and Maeda, H., 2012. Stability of endogenous reference genes in postmortem human brains for normalization of quantitative real-time PCR data: Comprehensive evaluation using geNorm, NormFinder, and BestKeeper. *Int. J. Legal Med.*, **126**: 943-952. <https://doi.org/10.1007/s00414-012-0774-7>

- Wang, Y. and Zhang, S., 2012. EF1 α is a useful internal reference for studies of gene expression regulation in amphioxus, *Branchiostoma japonicum*. *Fish Shellf. Immunol.*, **32**: 1068-1073. <https://doi.org/10.1016/j.fsi.2012.03.001>
- Winchester, J.F., Salsberg, J.A. and Levin N.W., 2003. Beta-2 microglobulin in ESRD: An in-depth review. *Adv. Ren. Replace Ther.*, **10**: 279-309. <https://doi.org/10.1053/j.arrt.2003.11.003>
- Wong, C.K., Yeung, H., Woo, P. and Wong, M.H., 2001. Specific expression of cytochrome P4501A1 gene in gill, intestine and liver of tilapia exposed to coastal sediments. *Aquat. Toxicol.*, **54**: 69-80. [https://doi.org/10.1016/S0166-445X\(00\)00173-9](https://doi.org/10.1016/S0166-445X(00)00173-9)
- Yang, C.G., Wang, X.L., Tian, J., Liu, W., Wu, F., Jiang, M. and Wen, H., 2013. Evaluation of reference genes for quantitative real-time RT-PCR analysis of gene expression in Nile tilapia (*Oreochromis niloticus*). *Gene*, **527**: 183-192. <https://doi.org/10.1016/j.gene.2013.06.013>
- Zheng, J.L., Luo, Z., Zhu, Q.L., Tan, X.Y., Chen, Q.L., Sun, L.D. and Hu, W., 2013. Molecular cloning and expression pattern of 11 genes involved in lipid metabolism in yellow catfish *Pelteobagrus fulvidraco*. *Gene*, **531**: 53-63. <https://doi.org/10.1016/j.gene.2013.08.028>
- Zheng, W.J. and Sun, L., 2011. Evaluation of housekeeping genes as references for quantitative real time RT-PCR analysis of gene expression in Japanese flounder (*Paralichthys olivaceus*). *Fish Shellf. Immunol.*, **30**: 638-645. <https://doi.org/10.1016/j.fsi.2010.12.014>
- Zhou, R.X., Meng, T., Meng, H.B., Cheng, D.X., Bin, S.Y., Cheng, J. and Zhang, J.S., 2010. Selection of reference genes in transcription analysis of gene expression of the Mandarin fish, *Siniperca chuasti*. *Zool. Res.*, **31**: 141-146. <https://doi.org/10.3724/SP.J.1141.2010.02141>