



# Effect of Artificial Saliva Supplementation on Growth, Rumen Characteristics and Fecal Odor of Growing Naeemi Lambs Fed Total Mixed Ration

Mutassim M. Abdelrahman<sup>1</sup>, Ibrahim Alhidary<sup>1</sup>, Mohsen Alobre<sup>1</sup>, Riyadh S. Aljumaah<sup>1</sup> and Rifat Ullah Khan<sup>2\*</sup>

<sup>1</sup>Department of Animal Production, Faculty of Food and Agriculture Sciences, King Saud University, P.O. Box 2460, Riyadh 11451, Saudi Arabia

<sup>2</sup>College of Veterinary Sciences, Faculty of Animal Husbandry and Veterinary Sciences, The University of Agriculture, Peshawar, Pakistan

## ABSTRACT

We investigated the effect of feeding a complete feed as a total mixed ration (TMR) with different levels of artificial saliva (AS) on the rumen pH, epithelial tissues pigmentation, fecal odor and status of some trace minerals. Forty-five growing lambs were used in an 84-day trial. Lambs were randomly assigned to five dietary treatments as follows: TMR, AS1 (TMR +15 kg AS/Ton; 22.5 g/lamb/day), AS2 (TMR + 30 kg AS/Ton; 45 g/lamb/day), AS3 (TMR + 45 kg AS/ Ton; 67.5 g/lamb/day) and AS4 (TMR +90 kg AS/Ton; 135 g/lamb/day). The growth performance was not affected by the supplementation of AS. The addition of AS improved the rumen color and maintained the rumen pH of the lambs. The fecal H<sub>2</sub>S gas significantly ( $P<0.05$ ) higher in AS4 group compared to the TMR. However, serum iron (Fe) content was significantly ( $P<0.05$ ) higher in TMR fed group alone. From the findings of the present study, we concluded that AS supplementation improved rumen color, however, fecal H<sub>2</sub>S production was increased which needed to be further investigated.

## Article Information

Received 29 October 2021

Revised 07 November 2021

Accepted 24 November 2021

Available online 03 March 2022  
(early access)

## Authors' Contribution

MA, IH and MSA designed and outlined the experiment, wrote the manuscript and follow up of the feeding trial. RJ assisted in data collection and statistical analysis, and reviewed the article. MO execute the experiment and analyzed the samples. RUK edited the paper. All authors critically reviewed the data, wrote and revised the final version of the manuscript.

## Key words

Artificial saliva, Growing lambs, Iron, Rumen color, Rumen pH

## INTRODUCTION

Feeding a full feed as a total mixed ration (TMR) containing highly fermentable carbohydrate (CHO) may raise the risk of subacute rumen acidosis by lowering ruminal pH, compromising ruminant animal health and production (Alhidary *et al.*, 2016a; Alharthi *et al.*, 2021a, b). TMR fibre stimulates rumination, chewing activity, and saliva production, which helps to buffer rumen pH and maintain an ideal environment for rumen bacteria (Alhidary *et al.*, 2016b). Feeding TMR low in fiber content and small particle size will lead to decreased chewing activity and salivary production, ultimately lowering the rumen pH. Moreover, it is used to determine the chemical and physical characteristics of roughage as a source of effective fiber (Lee *et al.*, 2004). The negative or positive effects of feeding complete feed may be caused by the levels used and the chemical composition of the ingredients, especially when byproducts and premixes are used. For this reason, special consideration must be given

to using byproducts due to their negative effects, especially when used at high levels. For example, the dark-brownish color of rumen epithelium tissues may be caused by many factors including high iron intake, assuming that the pigmentation is an iron-flavonoid complex. Alhidary *et al.* (2016a) reported that TMR plus alfalfa can affect the production of a pigment that attaches firmly to the keratinized confirmed layer of the rumen epithelium tissues. The risk of rumen acidosis (low pH) can be reduced by controlling the rumen fermentation process by feeding high fiber or using feed additives such as artificial saliva (AS), which are designed to reduce the negative effects of feeding highly fermented CHO on rumen characteristics.

Salivary secretions are considered to be the primary natural buffering system that maintains rumen function by controlling the rumen environment, especially rumen pH. In addition, saliva plays an important role as a fluid medium for feed mastication, swallowing, and transport of ingesta through the rumen and reticulum. Krause and Oetzel (2006) and Jouany (2006) reported that the normal range of ruminal pH of 6.0–7.2 is the optimum environment for rumen microbial activities, which are mainly regulated by saliva. Thus, AS may play an important role in controlling rumen pH in cases of unbalanced rations and low NDF

\* Corresponding author: rukhan@aup.edu.pk  
0030-9923/2022/0001-0001 \$ 9.00/0  
Copyright 2022 Zoological Society of Pakistan

dietary levels. AS is produced in the form of powder containing a mixture of chemical compounds ( $\text{NaHCO}_3$ ,  $\text{KCl}$ ,  $\text{CaCl}_2$ ,  $\text{Na}_2\text{HO}_4 \cdot 12\text{H}_2\text{O}$ ,  $\text{NaCl}$ , and  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ) in appropriate proportions (McDougall, 1948).

The objective of this trial was to study the effects of feeding four levels of AS on the rumen characteristics, trace minerals, and fecal odor status of growing lambs fed complete feed as a TMR.

## MATERIALS AND METHODS

### *Ethical approval*

This study was approved by the Ehtical Committee on Animal Rights, King Saud University, Riyadh, Saudi Arabia.

### *Animals and study design*

A total of 45 growing lambs ( $25 \pm 3$  kg average body weight; 90 days old) were used in an 84-days trial. Lambs were weighed, ear tagged, vaccinated against clostridial diseases, and treated for internal and external parasites. Thereafter, the lambs were distributed randomly to group feeding pens, with three lambs per pen for fourteen days adaptation period for the new diets. On day 1 of the experimental period, the pens were randomly assigned to one of the five dietary treatments (nine lambs per treatment on the basis of weight). The dietary treatments were as follows: TMR, AS1 (TMR+15 kg AS/Ton, 22.5 g/lamb/day), AS2 (TMR+30 kg AS/Ton, 45 g/lamb/day); AS3 (TMR+45 kg AS/Ton, 67.5 g/lamb/day) and AS4 (TMR+90 kg AS/Ton, 135 g/lamb/day). Aartificial saliva is produced in a form of powder containing a mixture of chemical compounds ( $\text{NaHCO}_3$ ,  $\text{KCl}$ ,  $\text{CaCl}_2$ ,  $\text{Na}_2\text{HO}_4 \cdot 12\text{H}_2\text{O}$ ,  $\text{NaCl}$ , and  $\text{MgSO}_4 \cdot 7\text{H}_2\text{O}$ ) in appropriate proportions.

### *Performance traits*

The weights of offered feed and feed refusals were measured weekly, and the feed intake was calculated on a dry matter basis per pen. Lambs were weighed before morning feeding at 0730 h on day 1 of the study and at the end of the study. The gain to feed ratio for each lamb was calculated and expressed as body weight gain per kg of dry matter intake.

### *Serum minerals*

Blood samples were collected from each lamb via jugular venipuncture at the end of the study. Serum was obtained by centrifugation at  $2400 \times g$  for 15 min at 4 °C. Serum trace mineral concentrations were analyzed at the beginning and the end of the trial using ICP-MS (Inductively Coupled Plasma-Mass Spectrometer: Perkin Elmer, USA).

### *Rumen pH*

The pH values of ruminal fluid from three lambs/treatment were measured at 30-min intervals for 60 d using rumen pH/temperature sensor (Horiba Twin pH, Spectrum Technologies, Inc. 60544, Plainfield, IL, USA), and measurements were stored using a data logger and analyzed after downloading. The pH value of ruminal fluid was recorded every 30 min, and hourly means were calculated. Samples for rumen pH were taken on weekly basis after three h of feeding the AS. The average values for the whole experimental perios of each treatment were compared.

### *Rumen color*

At the end of the trial, four lambs from each treatment were randomly slaughtered, and rumen tissue samples were collected from the dorsal sac and used for color measurement. The color of rumen tissues was determined using a Minolta Chroma Meter (Konica Minolta, CR-400, Japan) with a CIELAB Color System for the color values ( $L^*$  = value designates lightness).

### *Fecal odor gases*

Before the end of the feeding trial, three lambs were selected from each treatment for the determination of fecal odor gases. Feces and urine were collected daily for 4 d and mixed well, and then homogenized samples were taken. Fecal samples were mixed with urine in a glass tube (50 g feces with 25 mL urine; 12 tubes per treatment). Gas concentrations were measured using Gastec detector tubes designed for  $\text{H}_2\text{S}$ ,  $\text{NH}_3$ , and acetic acid (Gastec Corp., Kitagawa, Japan).

### *Statistical analysis*

All data was analyzed as a Complete Randomized Design with repeated measurements using the PROC Mixed and GLM Models of SAS. Means were separated using Tukey test. Significant differences were considered at  $P < 0.05$ .

## RESULTS

### *Feed intake, body weight gain, and feed conversion ratio*

Table I shows the effect of treatments on the growth performance and feed efficiency of growing Naeemi lambs fed different levels of AS. Lambs fed TMR and AS had no significant effect on body weight gain, feed intake and FCR.

### *Rumen fluid pH*

Table II shows the rumen pH of the TMR and the treatment groups. As a general trend, addition of AS at all

levels increased ( $P > 0.05$ ) the rumen pH compared with TMR.

**Table I. Effect of dietary different levels of artificial saliva on the performance of growing lambs fed complete feed as TMR.**

Measurements treatments	TFI (Kg)	BWG (Kg)	FCR
AB	140.51a	18.08b	7.94a
TMR	142.59a	26.83a	5.32b
AS1	133.31d	22.14c	6.03b
AS2	135.23d	22.07c	6.15b
AS3	129.70b	21.54c	6.04b
AS4	125.73b	19.44b	7.29a
SEM	1.86	0.74	0.23
P value	0.001	0.000	0.000

AB, traditional feeding; TMR, Total mixed ration; AS1, artificial level 1; AS2, artificial level 2; AS3, artificial level 3; AS4, artificial level 4. TFI, total feed intake; BWG, body weight gain and FCR, feed conversion ratio; SEM, Standard error of the means. Column with different superscripts with small letters show significant differences

**Table II. Time distribution throughout the day of rumen pH changes for lambs fed artificial saliva.**

pH	Traditional AB	TMR	T1 AS1	T2 AS2	T3 AS3	T4 AS4
less than 5.3	0	870	0	0	0	0
5.3–5.8	0	570	15	1065	0	968
5.8–7	1440	0	1425	375	1440	473

For abbreviations, see Table I.

#### Rumen color

The effects of treatment on the color of rumen tissues are presented in Table III. As a general trend, feeding all AS levels caused an improvement in color values (L values) compared with TMR alone. The rumen tissue color of lambs from the AS1 group showed a significant ( $P < 0.05$ ) improvement in color values compared with TMR and other levels of AS2, 3, and 4.

#### Fecal odor

Regarding the unacceptable fecal odors linked to acetic acid,  $H_2S$ , and  $NH_3$  emissions, there was no significant effect of treatment on acetic acid emission levels, but significant differences were reported for  $H_2S$  emissions as a result of the treatments. A significantly lower fecal  $H_2S$  level was reported for lambs fed TMR, and a higher level was reported for lambs from AS4. Furthermore, for  $NH_3$  levels, no significant differences were found between the control and the treatment groups (Table IV).

**Table III. Minimum, maximum, and average rumen fluid pH during the day in treatments supplemented with artificial saliva.**

	Traditional AB	TMR	T1 AS1	T2 AS2	T3 AS3	T4 AS4
Mean	6.05	5.28	5.88	5.68	6.05	5.70
Minimum	4.85	4.66	5.19	4.72	4.91	4.94
Maximum	7.01	6.62	6.38	6.82	6.90	7.18
ST.V	0.41	0.33	0.18	0.42	0.32	0.47

For abbreviations, see Table I. ST.V, standard of variation.

**Table IV. Effect of dietary different levels of artificial saliva on fecal odor gases of growing lambs fed complete feed as TMR.**

Measurements treatment	Acetic acid (ppm)	$H_2S$ (ppm)	$NH_3$ (ppm)
AB	1.69	31.25b	110.00
TMR	1.63	151.25c	50.00
AS1	1.05	96.25c	106.25
AS2	1.19	190.00c	81.25
AS3	1.56	360.00d	60.00
AS4	2.00	782.50a	50.00
SEM	0.17	22.00	8.85
P value	0.650	0.010	0.149

For abbreviations, see Table I. SEM, Standard error means. Columns with different superscripts with small letters showed significant differences.

#### Trace elements

Table V shows the effect of the treatments on trace minerals concentrations in the serum of the experimental lambs. The results showed a significant difference between all treatment groups in terms of Fe concentrations. Fe concentration was significantly ( $P < 0.05$ ) higher in TMR compared to the treatments.

## DISCUSSION

Feeding ruminant animal complete feed with high fermentable carbohydrate increases the levels of ruminal organic acids such as volatile fatty acids (VFAs) and lactic acid accumulation, which may cause acidosis by decreasing pH. As a consequence, the pH of the rumen drops. Ruminal pH levels of 5.6 and lower are used to characterize subacute ruminal acidosis (SARA) and ruminal acute acidosis (RAA) (Nagaraja and Tigemeyer, 2007). Immediately after feeding, rumen pH normally drops, but it causes problems when it is low for more than 3 h. This can have a negative effect on animal health and

**Table V. Effect of dietary different levels of artificial saliva on blood serum mineral concentrations of growing lambs fed complete feed as TMR.**

Trace elements	Treatments						SEM	P value
	AB	TMR	AS1	AS2	AS3	AS4		
Mn, $\mu\text{g/L}$	2.95 <sup>b</sup>	4.13 <sup>a</sup>	4.10 <sup>a</sup>	4.02 <sup>a</sup>	3.76 <sup>a</sup>	4.31 <sup>a</sup>	0.13	0.001
Fe, $\text{m}\text{g/L}$	1.32	1.40	1.43	1.48	1.34	1.50	0.02	0.23
Co, $\mu\text{g/L}$	0.42	0.48	0.49	0.48	0.47	0.48	0.01	0.15
Cu, $\text{m}\text{g/L}$	1.13	0.93	1.00	0.96	1.02	1.48	0.08	0.60
Zn, $\text{m}\text{g/L}$	0.58 <sup>a</sup>	0.68 <sup>b</sup>	0.77 <sup>b</sup>	0.71 <sup>b</sup>	0.67 <sup>b</sup>	1.02 <sup>a</sup>	0.04	0.02
Se, $\mu\text{g/L}$	112.58	129.61	151.67	120.89	134.93	157.36	7.33	0.54

For abbreviations, see Table I. <sup>a, b</sup> Means with different superscripts within each row were significantly different. SEM, Standard error of means.

productivity and cause dark coloration of the rumen epithelial tissues (Shusterman, 1992). This coloration may also be caused by a high intake of iron from plant byproducts such as PKC, which contain high levels of minerals, especially iron (Alimon, 2004) and phenols. The iron reacts with phenols when the pH drops in the rumen fluid to form a blackening ferric-polyphenol compound (Hamada *et al.*, 1969), which may lead to discoloration of the rumen epithelial tissues. The complete feed confirmed the formation of black stain in the rumen tissues, which makes it unacceptable to consumers (Hamada *et al.*, 1969; McGuire *et al.*, 1985). On the other hand, incomplete microbial degradation of dietary nutrients such as protein and carbohydrates in complete feed as a result of pH decreases which may increase odorous compounds in the manure (Zhu and Jacobson, 1999). These can harm human health and cause environmental pollution. This trial examined the effect of using AS as a rumen buffering feed additive to control rumen pH and avoid acidosis, and consequently to enhance the rumen microbial fermentation process to reduce fecal odors.

In this study, the lambs fed TMR had pH values in the rumen fluid as follows: 5.3 for 870 min after feeding and from 5.3 to 5.8 for 570 min per day, with a mean value of 5.28 (Tables II, III). This can be linked to the very dark color in rumen tissues ( $L = 29.1$ ). The addition of AS caused an increase in pH values compared to the TMR group and improved rumen tissue color. This result confirms that feeding TMR can reduce rumen fluid pH and form blackening ferric-polyphenol compounds that result from the inclusion of palm kernel meal in the complete feed. As a general trend, the addition of AS at all levels increased the rumen pH compared with TMR, but was almost the same as that of the AB group.

Furthermore, feeding all AS levels caused an improvement in color values (L values) compared with TMR alone, but was lower than that of rumen tissues in the

AB group. The rumen tissue color of lambs from the AS1 group showed a significant ( $P < 0.05$ ) improvement in color values compared with TMR and other levels of AS2, 3, and 4, but tissue from AB lambs showed a significantly brighter coloring compared with the other groups. The darker color, lower rumen fluid, and other performance measures of growing lambs fed TMR consisting of PKC compared with traditional feeding have been reported in several recent studies (Alhidary *et al.*, 2016a, b, 2017; Abdelrahman *et al.*, 2017, 2019). In order to reduce serious health disorders caused by rumen acidosis that can negatively affect ruminant animal performance, it is crucial to search for a good alternative to improve rumen pH (Aschenbach *et al.*, 2011; Owens and Basalah, 2016). Different commercial buffering additives (Mao *et al.*, 2017) have been introduced to the market to help with maintenance of proper rumen pH levels. These include AS, sodium bicarbonate, calcified seaweed, and malic acid, which can be added to TMR (Alhidary *et al.*, 2017; Abdelrahman *et al.*, 2019). A recent study conducted by Alhidary *et al.* (2019), supported our finding that the addition of different buffers to TMR of growing lambs can improve their growth performance, rumen pH, fermentation process, and meat quality. Furthermore, Abdelrahman *et al.* (2019) conducted an experiment using malic acid as a rumen buffering agent in TMR consisting of palm kernel meal for growing lambs and measured its effect on rumen characteristics. They concluded that malic acid caused a significant improvement in rumen pH and rumen tissue coloring by decreasing the lactic acid accumulation compared with regular TMR. These findings are in agreement with the results of this trial. The main problem with using different buffering agents is determining the proper level that can be used without negative effects on ruminant animal health and productivity.

On the other hand, a higher accumulation of iron in the rumen tissues of lambs fed TMR, AS2, and AS4

compared with those of AB, AS1, and AS3, which may indicate that high iron intake and accumulation in the rumen tissues resulting from the formation of ferric-polyphenol compounds play a crucial role in dark coloring, in addition to other factors. Regarding the trace mineral status of growing lambs as a result of feeding TMR with or without AS at different levels, feeding lambs complete feed improved blood serum trace mineral levels compared with traditional feeding because of the well-balanced mineral additives in the complete feed. There was no significant effect of the treatment on trace mineral levels in the blood serum, except for Mn and Zn. All of the trace mineral values were within normal levels according to Puls (1990). Moreover, the Mn and Zn levels in the blood serum of lambs from the AB group were significantly lower than those in the other dietary groups (Table V).

Fecal gas odors are an acute environmental problem (Singh and Rashid, 2017) and have an adverse effect on human health and the environment (Ushida *et al.*, 2003; Singh and Rashid, 2017). Manipulation of the rumen fermentation process by using feed additives is a potential solution for reducing emissions of these gases. In this study, the effects of using AS on the fecal unacceptable smells, which are linked to acetic acid, H<sub>2</sub>S, and NH<sub>3</sub> emissions, were analyzed. There was no significant effect of treatment on acetic acid emission levels, but significant differences were reported for H<sub>2</sub>S emissions as a result of the treatments. A significantly lower fecal H<sub>2</sub>S was reported for lambs fed AB, but a higher value was reported for lambs from AS3 and AS4. Lambs from the TMR, AS1, and AS2 groups were not significantly different ( $P > 0.05$ ; Table IV), but numerically lower values were found for lambs from the AS1 group. Furthermore, for NH<sub>3</sub> levels, no significant differences were found, but there was a trend of variation between all treatment groups. To our knowledge, no studies have investigated the effect of using AS on fecal gas emissions, especially in ruminant animals. The numerical differences detected between all treatment groups in term of fecal gases gives a general indication of improvement as a result of using AS as a feed additive to improve nutrient digestibility, especially protein and carbohydrate.

## CONCLUSION

AS supplementation with complete feed as a TMR improved rumen fluid pH and rumen tissue coloring, but a negative effect on the growth performance of lambs was found at high levels of AS supplementation. Moreover, AS1 and AS3 supplementation reduced the iron accumulation in rumen epithelial tissues, which may indicate that high iron intake plays a role in dark rumen

pigmentation by forming blackening ferric-polyphenol compounds, in addition to other possible factors that are not yet well-defined. Including Level 1 AS (15 kg/ton: 22.5 gm/ lamb/day) is highly recommended to improve the rumen fermentation process by increasing rumen pH value to  $>5.8$ . Moreover, this can reduce the dark pigmentation of rumen epithelial tissues and fecal NH<sub>3</sub> emissions.

## ACKNOWLEDGMENTS

The authors would like to extend their sincere appreciation to the Deanship of Scientific Research at King Saud University for funding this work through research group RG -1441-408.

## REFERENCES

- Abdelrahman, M., Alhidary, I.A., Albaadani, H.H., Alobre, M., Khan, R.U. and Aljumaah, R.S., 2019. Effect of palm kernel meal and malic acid on rumen characteristics of growing Naemi lambs fed total mixed ration. *Animals*, **9**: 408. <https://doi.org/10.3390/ani9070408>
- Abdelrahman, M.M., Alhidary, I., Alyemni, A.H., Khan, R.U., Bello, A.R., Al-Saiady, M.Y., and Amran, R.A., 2017. Effect of alfalfa hay on rumen fermentation patterns and serum biochemical profile of growing Naemi lambs with *ad libitum* access to total mixed rations. *Pakistan J. Zool.*, **49**: 1519–1522. <https://doi.org/10.17582/journal.pjz/2017.49.4.sc6>
- Alharthi, A.H., Al-Baadani, H.H., Al-Badawi, M.A., Abdelrehman, M.A., Alhidary, I.A. and Khan, R.U., 2021b. Effects of sunflower hulls on productive performance, digestibility indices and rumen morphology of growing Awassi lambs fed with total mixed rations. *Vet. Sci.*, **8**: 174-184. <https://doi.org/10.3390/vetsci8090174>
- Alharthi, A.S., Alobre, M.M., Abdelrahman, M.M., Al-Baadani, H.H., Swelum, A.A., Khan, R.U. and Alhidary, I.A., 2021a. The effects of different levels of sunflower hulls on reproductive performance of yearly ewes fed with pelleted complete diets. *Agriculture*, **11**: 959. <https://doi.org/10.3390/agriculture11100959>
- Alhidary, I., Abdelrahman, M.M., Alyemni, A.H., Khan, R.U., Al-Mubarak, A.H. and Albaadani, H.H., 2016a. Characteristics of rumen in Naemi lamb: morphological characteristics in response to altered feeding regimen. *Acta Histochem.*, **118**: 331–337.

- <https://doi.org/10.1016/j.acthis.2016.03.002>
- Alshamiry, F.A., Abdelrahman, M.M., Alyemni, A.H., Khan, R.U., Al-Saiady, M.Y., Amran, R.A. and Alshamiry, F.A., 2016. 2016b. Effect of alfalfa hay on growth performance, carcass characteristics, and meat quality of growing lambs with ad libitum access to total mixed rations. *Rev. Bras. Zootec.*, **45**: 302–308. <https://doi.org/10.1590/S1806-92902016000600004>
- Alhidary, I.A., Abdelrahman, M.M., Aljumaah, R.S., Alyemni, A.H., Ayadi, M.A., and Al-Saiady, M.Y., 2017. Rumen discoloration of growing lambs fed with diets containing different levels of neutral detergent fiber. *Pakistan J. Zool.*, **49**: 1847–1855. <https://doi.org/10.17582/journal.pjz/2017.49.5.1847.1855>
- Alhidary, I.A., Abdelrahman, M.M., and Elsabagh, M., 2019. A comparative study of four rumen buffering agents on productive performance, rumen fermentation and meat quality in growing lambs fed a total mixed ration. *Animal*, **13**: 2252–2259. <https://doi.org/10.1017/S1751731119000296>
- Alimon, A.R., 2004. The nutritive value of palm kernel cake for animal feed. *Palm Oil Dev.*, **40**: 12–14.
- Aschenbach, J.R., Penner, G.B., Stumpff, F., and Gäbel, G., 2011. Ruminant Nutrition Symposium: Role of fermentation acid absorption in the regulation of ruminal pH. *J. Anim. Sci.*, **89**: 1092–1107. <https://doi.org/10.2527/jas.2010-3301>
- Erwanto, I.R., Zakaria, W.A., and Prayuwidayati, M., 2012. The use of ammoniated zeolite to improve rumen metabolism in ruminant. *Anim. Prod.*, **13**: 138–142.
- Ghaemnia, L., Bojarpour, M., Mirzadeh, K.H., Chaji, M., and Eslami, M., 2010. Effects of different levels of zeolite on digestibility and some blood parameters in Arabic lambs. *J. Anim. Vet. Adv.*, **9**: 779–781. <https://doi.org/10.3923/javaa.2010.779.781>
- Hamada, T., Maeda, S., and Kameoka, K., 1969. Effects of minerals on formation of color in the rumen epithelium of kids. *J. Dairy Sci.*, **53**: 588–591. [https://doi.org/10.3168/jds.S0022-0302\(70\)86257-9](https://doi.org/10.3168/jds.S0022-0302(70)86257-9)
- Jouany, J.P., 2006. Optimizing rumen functions in the close-up transition period and early lactation to drive dry matter intake and energy balance in cows. *Anim. Reprod. Sci.*, **96**: 250–264. <https://doi.org/10.1016/j.anireprosci.2006.08.005>
- Krause K.M., and Oetzel G.R., 2006. Understanding and preventing subacute ruminal acidosis in dairy herds: A review. *Anim. Feed Sci. Tech.*, **126**: 215–236. <https://doi.org/10.1016/j.anifeedsci.2005.08.004>
- Lee, W.S., Lee, B.S., Lee, S.C., Lee, S.S., Lee, S.Y., Lee, D.Y., and Ha, J.K., 2004. Effects of rice straw and rice hull supplement on rumination and chewing behavior in Hanwoo steers. *Kor. J. Anim. Sci.*, **46**: 49–54. <https://doi.org/10.5187/JAST.2004.46.1.049>
- Mao, S., Huo, W., Liu, J., Zhang, R., and Zhu, W., 2017. In vitro effects of sodium bicarbonate buffer on rumen fermentation, levels of lipopolysaccharide and biogenic amine, and composition of rumen microbiota. *J. Sci. Fd. Agric.*, **97**: 1276–1285. <https://doi.org/10.1002/jsfa.7861>
- McDougall, E.I., 1948. Studies on ruminant saliva. 1. The composition and output of sheep's saliva. *Biochem. J.*, **43**: 99–109. <https://doi.org/10.1042/bj0430099>
- McGuire, S.O., Miller, W.J., Gentry, R.P., Neathery, M.W., Ho, S.Y., and Blackmon, D.M., 1985. Influence of high dietary iron as ferrous carbonate and ferrous sulfate on iron metabolism in young calves. *J. Dairy Sci.*, **68**: 2621–2628. [https://doi.org/10.3168/jds.S0022-0302\(85\)81146-2](https://doi.org/10.3168/jds.S0022-0302(85)81146-2)
- Nagaraja, T.G., and Titgemeyer, E.C., 2007. Ruminal acidosis in beef cattle: the current microbiological and nutritional outlook. *J. Dairy Sci.*, **90**: E17–E38. <https://doi.org/10.3168/jds.2006-478>
- Nik-Khan, A., Sadeghi, A. A., 2002. *Natural clinoptilolite-tuff effects on health homo-immuno parameters in newborn calves*. Zeolite '02, 6<sup>th</sup> International Conference, Occurrence, Properties and Utilisation of Natural Zeolites, pp. 253-265.
- Nockels, C.F., Kintner L.D., and Pfander, W.H., 1966. Influence of ration on morphology, histology and trace mineral content of sheep rumen papillae. *J. Dairy Sci.*, **49**: 1068–1074. [https://doi.org/10.3168/jds.S0022-0302\(66\)88019-0](https://doi.org/10.3168/jds.S0022-0302(66)88019-0)
- Owens, F.N. and Basalan, M., 2016. *Ruminal fermentation*. In Rumenology (ed. D Millen, MDB Arrigoni and RDL Pacheco), Springer International Publishing Switzerland, Cham, Switzerland. pp. 63–102. [https://doi.org/10.1007/978-3-319-30533-2\\_3](https://doi.org/10.1007/978-3-319-30533-2_3)
- Puls, R., 1988. *Mineral levels in animal health. Diagnostic data*. Sherpa International, USA.
- Sadeghia, A.A., and Shawrang, P., 2006. The effect of natural zeolite on nutrient digestibility, carcass traits and performance of Holstein steers given a diet containing urea. *Anim. Sci.*, **82**: 163–167. <https://doi.org/10.1079/ASC200524>
- Shusterman, D., 1992. Critical review: The health significance of environmental odor pollution. *Arch. environ. Hlth.*, **47**: 76–87. <https://doi.org/10.1080/0>

- 0039896.1992.9935948
- Singh A., and Rashid, M., 2017. Impact of animal waste on environment, its managerial strategies and treatment protocols to reduce environmental contamination. *Vet. Sci. Res. J.*, **8**: 1-2. <https://doi.org/10.15740/HAS/VSRJ/8.1and2/1-12>
- Sofi, S., Singh, J., Rafiq, S., and Rashid, R., 2017. Fortification of dietary fiber ingredients in meat application: A review. *Int. J. Biochem. Res. Rev.*, **19**: 1–14. <https://doi.org/10.9734/IJBCRR/2017/36561>
- Ushida, K., Hashizume, K., Miyazaki, K., Kojima, Y., and Takakuwa, S., 2003. Isolation of *Bacillus* sp. as a volatile sulfur-degrading bacterium and its application to reduce the fecal odor of pig. *Asian Aust. J. Anim. Sci.*, **16**: 1795–1798. <https://doi.org/10.5713/ajas.2003.1795>
- Yang, W.Z., and Beauchemin, K.A., 2009. Increasing physically effective fiber content of dairy cow diets through forage proportion versus forage chop length: Chewing and ruminal pH. *J. Dairy Sci.*, **92**: 1603–1615. <https://doi.org/10.3168/jds.2008-1379>
- Yazdani, A.R., and Hajilari, D., 2009. Application of natural zeolite on blood characteristics, physiological reactions and feeding behaviors of finishing Holstein beef steers. *Ind J. Anim. Res.*, **43**: 295–299.
- Zhu, J., and Jacobson, L.D., 1999. Correlating microbes to major odorous compounds in swine manure. *J. Environ. Quart.*, **28**: 737–744. <https://doi.org/10.2134/jeq1999.00472425002800030001x>

Online First Article