



Enhanced Phytase Production from Indigenous *Bacillus subtilis* KT004404 by Response Surface Methodology to be Used as Poultry Feed Supplement

Aqsa Javaid, Masoom Yasinzai* and Khunsa Saeed Kiyani

Suleman Bin Abdullah Bin Aba-Alkhail Centre for Interdisciplinary Research in Basic and Applied Sciences, International Islamic University, Islamabad, Pakistan 44000

ABSTRACT

The aim of this experimental study was the optimization of phytase production from *Bacillus subtilis* KT004404 through statistical model (Response Surface Methodology) by utilizing centrally composite design (CCD). The optimal phytase activity obtained after CCD design was 132.32 U/mL at 35 °C, pH 6.5, inoculum density of 2.5mL after incubating for 24h at 75 rpm. Extracellular enzymes were then partially purified in 2 steps (ammonium sulfate precipitation and gel filtration chromatography using Sephadex G-50). The molecular weight of extracellular phytase from *B. subtilis* KT004404 estimated by SDS-PAGE was 30 kDa. The K_m and V_{max} of *B. subtilis* KT004404 phytase were 0.175 mM and 250 U/mL (phytase activity) respectively. The optimum temperature for partially purified extracellular phytase was 40°C, and the optimum pH range for extracellular phytase activity was pH 6.5-7.0. The extracellular enzyme activity was not significantly affected by Cu^{2+} and Mn^{2+} metal ions while the inhibitory effect of Ca^{2+} , Co^{2+} , Zn^{2+} , Hg^{2+} and Fe^{2+} was recorded with 27.12%, 32%, 57.31%, 14.76% and 13.54% residual activity.

Article Information

Received 20 September 2021
Revised 05 March 2022
Accepted 29 March 2022
Available online 02 August 2022
(early access)

Authors' Contribution

MY supervised the work and provided major guidance during drafting of the paper. AJ conducted all the experimental work including isolation, optimization and manuscript write up. KSK contributed during the lab experimentation and performed all the spectroscopy in the lab.

Key words

Bacillus subtilis KT004404, Phytase, Poultry, Centrally composite design (CCD), Response surface methodology (RSM)

INTRODUCTION

Phosphorus is stored in plants in the form of phytic acid particularly in grains, legumes, and cereals which represents about 75%-80% of the stored phosphorus present in nature (Al-Kandari *et al.*, 2021). Phytic acid has significant chelating potential under normal physiological conditions due to which it limits the bioavailability of essential mineral complexes such as zinc, calcium, iron, and magnesium (Singh, 2008). Moreover, it also inhibits the activity of digestive enzymes by binding with amino acids and proteins (Cowieson *et al.*, 2006). Phytic acid due to its anti-nutrient potential also causes phosphorus pollution via manure (Toor *et al.*, 2005). Phosphorous pollution in the

soil can be reduced between 30% to 50% by adding the phytase supplementation in feed (Alias *et al.*, 2017).

The bioavailability of phosphorous along with other minerals can be enhanced by supplementing the feed with phytic acid-degrading enzyme phytase (EC 3.1.3.8, myo-inositol hexakisphosphate phosphohydrolase). Phytase have significant application in agriculture, transgenic plant technology, and horticulture (Laboure *et al.*, 1993). Phytase enzyme catalyzes stepwise hydrolysis of myo-inositol hexakisphosphate and release phosphorous to be utilized in various biochemical reactions (Vats and Banerjee, 2004).

Bacteria, fungi, and yeast are reported to produce various classes of phytase. Different phytases differ in their structure, catalytic activity, and optimal conditions (Kumar *et al.*, 2017). There is no single phytase which can meet the diverse environmental and commercial need therefore, there is an increased interest in screening the microorganism with ability to produce novel phytase to improve the nutritive value of feed containing phytic acid (Alias *et al.*, 2017). Due to broad industrial application, the isolation, production and optimization of phytase has attracted significant attention.

The use of enzymes in poultry feed has posed many challenges because of thermal treatment during the

* Corresponding author: rector@iiu.edu.pk, aqsa.phdns27@iiu.edu.pk
0030-9923/2022/0001-0001 \$ 9.00/0



Copyright 2022 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

pelleting process, which causes decline or complete loss of enzyme activity due to denaturation (Kumar *et al.*, 2016). Enzymes exhibiting optimal activity at high temperatures and acidic pH are significantly important for diverse applications in the poultry feed industry (Naves *et al.*, 2012; Singh and Satyanarayana, 2015).

Traditional method for the optimization of enzymes known as one factor at a time (OFAT) have numerous downsides such as requirement of running large number of experimental runs and lack of information about the interaction of different parameters. However, statistical, or mathematical modeling for optimization of parameters for enzyme production offers multiple advantages over traditional OFAT method (Kumari and Bansal, 2021). Response surface methodology (RSM) is widely utilized statistical method for optimizing and analyzing the interactions between different experimental variables during optimization reaction (Jain and Singh, 2017; Shahryari *et al.*, 2018; Pable *et al.*, 2019; Jatuwong *et al.*, 2020; Kumari and Bansal, 2021)

The main objective of the current study was directed towards the selection of media components for optimal phytase production from *B. subtilis* KT004404 by using RSM and characterization of purified phytase enzyme to analyze the potential use as poultry feed supplement.

MATERIALS AND METHODS

Culture conditions and maintenance

The already identified and isolated *B. subtilis* KT004404 strain from hot water spring from Kashmir (Pakistan) was obtained from International Islamic University, Islamabad, Pakistan, and cultures were maintained at 4 °C in form of glycerol stocks.

Bacterial inoculum preparation

B. subtilis was cultivated in 50 mL of nutrient broth (Sigma-Aldrich Co., St Louis, USA) in Erlenmeyer flasks (250 mL), 0.2% dextrose at 37 °C and 150 rpm. The culture fluid was used as seed inoculum (CFU 1.39×10^6 mL⁻¹) after 24 h of incubation in the nutrient broth.

Phytase plate assay

The ability of *B. subtilis* KT004404 to exogenously produce phytase enzyme was identified by confirmatory tests proposed by Yanke *et al.* (1998). Phytase screening media composition contained glucose 1% (w/v), ammonium sulfate 0.2% (w/v), urea 2% (w/v), citric acid 0.3% (w/v), sodium citrate 0.2% (w/v), MgSO₄·7H₂O, 0.1% (w/v), sodium phytase 0.5% (w/v), iron sulfate 0.02% (w/v). The phytase-containing media was autoclaved at 121 °C for about 15 min at 15 psi pressure conditions

to ensure sterilization. Flooding plates with 2% cobalt chloride solution with 6.25% ammonium molybdate and 0.42% ammonium vanadate (1:1). Diameters of zones of clearance, an indicator of the phytase production were measured through vernier caliper scale.

Quantitative assay for phytase activity

Phytase activity was determined by colorimetric method using ferrous sulfate-molybdenum blue method (Murphy and Riley, 1962). 100 µL phytase solution and 900 µL (1 mM) substrate solution were incubated for 30 min. After 30-min incubation, the stop solution 1 mL of TCA (5%) was added followed by addition of 1 mL ferrous sulfate. 4mM solution of potassium dihydrogen phosphate was used to plot standard curve. One phytase unit is the activity of phytase that generates 1 µmol of inorganic phosphorus per min from an excess of the substrate at 37 °C. Protein concentration in filtrate was determined by Lowry method (Waterborg *et al.*, 1996).

Selecting source of nitrogen and carbon via one variable at a time

Nitrogen and carbon sources of energy were screened in a preliminary study, by the one-factor optimization method (OFT). Agro-industrial substrates such as sorghum, linseed, cornmeal, soybean, millet, sesame, wheat bran, rice husk, canola meal, sorghum were selected for analyzing the best carbon source in production media for enhanced phytase production (Sadh *et al.*, 2018). Similarly, during submerged fermentation, various nitrogen substrates like, peptone, casein, yeast extract, malt and beef extract, and ammonium nitrate were added in the medium (1% w/w in basal media) as nitrogen sources. Initially, production media was inoculated with 1 mL of inoculum (OD₆₀₀ = 1.5, CFU = 1.39×10^6 mL⁻¹) in order to maintain a constant metabolic activity in all the experiments.

Submerged fermentation

50 ml erlenmeyer shake flasks were used to produce phytase containing different nitrogen and carbon sources of energy for bacteria. The basal medium contained wheat bran (5g/L), calcium chloride (0.2%), magnesium sulfate heptahydrate (0.2g/L), and distilled H₂O to make 1 L production media. Sterilization of the flask was done at 121 °C for 15 min at 15 lbs pressure. 1 mL (v/v) inoculum was added in the basal media and incubated at 37 °C for 24 h.

Experimental design and data analysis

Fermentation parameters optimization via RSM using centrally composite design

Response surface methodology using CCD was

used for optimization of phytase production. Varying concentrations of carbon and nitrogen substrates were used in the production medium for phytase production. Phytase activity in the medium (U/mL) was the dependent response while five independent parameters were selected to determine optimal variable combination namely temperature (X_1), medium pH (X_2), Inoculum size (X_3), the incubation period (X_4), and shaking intensity (X_5). Three different levels of all selected independent variables were studied and total of 50 experimental runs were analyzed as shown in Figure 1. The flasks containing basal medium were sterilized at 121 °C for 15 min, then inoculated with bacterial (w/v) inoculum followed by 24h incubation at varying temperature range. Combined effects of five variables on phytase production by *B. subtilis* was studied within a specific range as shown in Table I.




Fig. 1. Qualitative assay for detecting extracellular phytase production by *Bacillus subtilis* KT004404 on phytase-screening medium (PSM) with Na-phytate. Zones of clearance formed around the wells (A) and (B) loaded with bacterial filtrate while well (C) was loaded with control and showed no clearance zone.

Statistical software ‘Design Expert 6.0’ Stat-Ease, Inc., Minneapolis, USA, was used for analyzing the experimental design and to generate data (Table II).

The production of phytase in the medium was analyzed by a second-order polynomial equation for a 5-factor system fit to experimental values (phytase activity) which were obtained as following regression Equation 1.

$$Y = \beta_0 + \sum \beta_i x_i + \sum \beta_{ii} x_i^2 + \sum \sum \beta_{ij} x_i x_j \dots (1)$$

Y (predicted response variable), β_0 represents intercepts, β_i linear coefficients, β_{ii} is the squared effect; β_{ij} is the interaction coefficient. Multiple regression equation for five factors system in the full quadratic model to describe response surface is as follow (Equation 2):

$$Y = \beta_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_3 X_3 + \beta_4 X_4 + \beta_5 X_5 + \beta_{11} X_1^2 + \beta_{22} X_2^2 + \beta_{33} X_3^2 + \beta_{44} X_4^2 + \beta_{55} X_5^2 + \beta_{12} X_1 X_2 + \beta_{13} X_1 X_3 + \beta_{14} X_1 X_4 + \beta_{15} X_1 X_5 + \beta_{23} X_2 X_3 + \beta_{24} X_2 X_4 + \beta_{25} X_2 X_5 + \beta_{34} X_3 X_4 + \beta_{35} X_3 X_5 + \beta_{45} X_4 X_5 \dots (2)$$

Where Y represents Phytase activity, β_0 for offset term, $\beta_1, \beta_2, \beta_3$ show linear coefficients, $\beta_{11}, \beta_{22}, \beta_{33}$ show squared coefficients, $\beta_{12}, \beta_{23}, \beta_{13}$ are interaction coefficients which represent the following levels of Independent variable:

$X_1, X_2, X_3, X_4, X_5, X_{21}, X_{22}, X_{23}, X_{24}, X_{25}, X_1 X_2, X_1 X_3, X_1 X_4, X_1 X_5, X_2 X_3, X_2 X_4, X_2 X_5, X_3 X_4, X_3 X_5, X_4 X_5$

To validate the model, the estimated phytase activity was compared with predicted software values. Statistical analysis including standard deviation, ANOVA and Duncan multiple range test on the obtained results were applied using IBM SPSS Statistics Ver.17 (IBM corporation, USA).

Table I. Coded levels of process variables and experimental range for phytase production by *Bacillus subtilis*.

Process variables	Units	Range		
		-1	0	+1
X_1 -Temperature	°C	20	35	50
X_2 -Medium pH	pH	4	6.5	9
X_3 -Inoculum size	%	0.1	2.5	5
X_4 -Incubation time	h	24	60	96
X_5 -Shaking intensity	rpm	50	75	100

Partial purification of phytase

The crude filtrate was subjected to ammonium sulfate precipitation at different percentage saturation (20%, 30%, 40%, 50%, 60%, 70%, 80% and 90%) with constant stirring. The precipitates were collected by centrifugation (15,000×g, 20 min) and then suspended in 0.2M acetate buffer (pH 6.0). Desalting of the enzyme was done by dialyzing the enzyme overnight in 50 mM acetate buffer (pH 6.0) followed by Sephadex G-50 gel filtration chromatography column (12/60, Pharmacia), equilibrated with 0.2M acetate buffer with pH 6.0. Enzyme fractions were eluting with 0.2M Na-acetate buffer at 0.2 cm³/min flow rate. The gel electrophoresis was performed on fractions collected by passing through 15% SDS-PAGE to determine the size of the enzyme. The gel slabs were stained with Coomassie stain (Neira-Vielma *et al.*, 2018).

Table II. Centrally composite design (CCD) of production parameters along with experimental and predicted phytase activity.

Run	X ₁ Temperature °C	X ₂ pH	X ₃ Inoculum size %	X ₄ Incubation time h	X ₅ ^b Rpm	Phytase production ^a U/mL	
						^c Actual	^d Predicted
1	35	6.5	2.55	60	75	120	113
2	20	9	0.1	96	50	98	61.4
3	20	4	5	24	50	132	97.42
4	35	6.5	2.55	60	75	165	113.10
5	20	4	0.1	96	50	99	81.82
6	50	4	0.1	96	50	76	69.01
7	50	9	0.1	24	100	46	37.46
8	20	4	0.1	96	100	59	59.41
9	35	6.5	2.55	60	75	110	113.10
10	20	4	5	24	100	89	76.26
11	35	0.55	2.55	60	75	89	88.11
12	35	6.5	2.55	24	75	132	113.10
13	50	9	5	96	50	79	49.30
14	50	4	0.1	96	100	69	71.87
15	20	4	5	96	50	85	91.83
16	50	4	5	96	100	75	67.88
17	50	9	0.1	24	50	65	26.50
18	20	9	5	24	50	63	55.32
19	50	9	0.1	96	100	52	67.88
20	35	6.5	2.55	60	15.56	82	102.24
21	35	12.4	2.55	60	75	28	33.19
22	70.6	6.5	2.55	60	75	38	18.70
23	35	6.5	2.55	60	75	99	102.21
24	50	4	5	24	50	62	64.25
25	50	9	5	96	100	42	46.27
26	50	9	5	24	50	32	38.64
27	20	9	0.1	24	100	43	21.40
28	35	6.5	2.55	-25.6229	75	16	26.88
29	20	4	0.1	24	100	43	45.24
30	50	4	5	96	50	46	71.65
31	20	9	5	24	100	41	34.54
32	50	4	0.1	24	100	42	45.43
33	35	6.5	2.55	60	75	79	121.23
34	20	9	5	96	50	38	53.62
35	35	6.5	2.55	60	134.46	98	81.55
36	35	6.5	2.55	145.623	75	79	56.41
37	50	4	5	24	100	61	69.70
38	20	9	0.1	96	100	32	39.44

Table continued on next page.....

Run	X ₁ Temperature °C	X ₂ pH	X ₃ Inoculum size %	X ₄ Incubation time h	X ₅ ^b Rpm	Phytase production ^a U/mL	
						^c Actual	^d Predicted
41	35	6.5	-3.277	60	75	0	13.76
42	20	4	0.1	24	50	55	60.02
43	20	9	0.1	24	50	21	35.81
44	35	6.5	2.55	60	75	96	111.21
45	-0.676	6.5	2.55	60	75	0	23.60
46	20	4	5	96	100	41	63.05
47	50	4	0.1	24	50	41	34.84
48	50	9	0.1	96	50	21	64.54
49	50	9	5	24	100	21	43.23
50	35	6.5	8.37711	60	75	42	32.53

^a Mean of three values, SD within 10%; ^b rpm, revolution per minute; ^c Actual response: Mean phytase activity (U/mL/min) obtained as triplicate; ^d Predicted response: Response predicted by the model.

Characterization of purified enzyme

The temperature effect on phytase activity was measured at 20°C, 30°C, 40°C, 50°C, 60 °C, and 70°C by incubating enzyme fractions with sodium phytate for 20 min. The optimum pH for phytase activity was determined at different pH range of 4,5,6,7,8,9,10. Various buffers were used to determine the optimum pH such as 0.06 M glycine-HCl (for pH range 2.0-3.6), 0.2M Na-acetate (pH range 3.6-5.8), phosphate-borate (pH range 6.0-8.0). The effect of the presence of metal ions in media were investigated to determine the activation or inhibitory effect of metallic ions on the phytase activity. The effect of different metal ions (Na⁺, Mn²⁺, Hg²⁺, Fe²⁺, Zn²⁺, Mg²⁺, Co²⁺, Ca²⁺) on the phytase activity was determined by incubating the enzyme fractions with 10 mM salt solutions of NaCl, MnCl₂, HgCl₂, FeSO₄, ZnCl₂, MgSO₄, CoCl₂, CaCl₂. The optimized enzyme was further analyzed for immobilization and stability studies on different inert medias with the prospect of being used as poultry feed supplement.

Enzyme kinetics and substrate specificity

The kinetic parameters of phytase hydrolysis of Na-phytate (at pH 5, 0.2 M Na-acetate buffer, and 40 °C) were studied to determine enzyme-substrate affinity. Various substrates used to determine the enzyme substrate specificity included sodium phytate, D-glucose-1-phosphate, glycerophosphate, fructose-6-phosphate, and p-nitrophenyl phosphate. The rate of Na-phytate hydrolysis was studied at different substrate concentrations (0.05 to 3 mM). K_m and V_{max} were found by Line weaver–Burk plot (Dokuzparmak *et al.*, 2017). All experimental data were means of triplicate determinations.

RESULTS

Phytase activity of *B. subtilis*

Figure 1 show zone of clearance (49 ± 1.414 mm) on phytate screening plates indicating and confirming the production of phytase enzyme by *B. subtilis*.

Our experiment has shown that wheat bran has the highest phytase production as compared to other carbon sources (71.21 U/ml). Peptone as nitrogen source gave the highest activity of phytase (80.45 U/mL) when compared with ammonium nitrate (62.3 U/mL) during fermentation.

Optimization of phytase production

Five important parameters (temperature, production media, pH, inoculation density, incubation period, and shaking flask intensity) were selected through preliminary studies were considered for centrally composite design analysis by RSM. These parameters were analyzed for the best suitable combination for optimal enzyme production. The highest phytase activity in the experimental design was 132.32 U/mL at 35 °C, pH 6.5, inoculum density of 2.5mL after incubating for 24 h at 75 rpm.

The results in Table II exhibit that predicted values for phytase production and experimental values by *B. subtilis* KT004404 are not significantly different. These results were analyzed by ANOVA. The second-order polynomial equation for phytase production as a function of temperature, pH, inoculum density, incubation time, and rpm was fitted to the model led to following regression Equation 3.

$$Y = -49.91 + 2.81X_1 + 12.06X_2 + 28.22X_3 + 1.42X_4 + 0.312X_5 + 0.105X_1X_2 - 0.05X_1X_3 + 0.005X_1X_4 + 0.016X_1X_5 - 0.72X_2X_3 + 0.010X_2X_4 + 0.001X_2X_5 - 0.077X_3X_4 - 0.021X_3X_5 - 0.002X_4X_5 - 0.072X_1^2 - 1.48X_2^2 - 2.649X_3^2 - 0.009X_4^2 - 0.005X_5^2 \dots (3)$$

Where X₁ -temperature for enzyme production, X₂

-pH of the medium, X_3 -inoculation density, X_4 -Incubation period, and X_5 -Shake flask rpm. The 3D response surface plots were generated to study the interaction of physicochemical parameters on enzyme production.

Centrally Composite Design was used to analyze the effect of five independent variables for optimizing the production of Phytase. Variability in temperature, pH, incubation substantially effect phytase production during fermentation.




Fig. 2. Response surface curve for production of phytase as a function of temperature incubation (A), pH-incubation (B), inoculum size-pH (C), temperature-inoculum size (D).

Validation of phytase production

As predicted by the model the combination of 5 different types of variables were determined for optimized phytase production. Higher similarity index was found between the predicted and experimental values for phytase production by *B. subtilis*. Predicted values were in agreement with experimentally determined phytase activity. The coefficient of determination (R^2) for the model was 0.705 which implies that there is 70.5% compatibility of predicted and experimental data. Phytase activity was measured in triplicates in all predicted model 50 experimental runs. Signal to noise ratio is measure by model precision. A ratio above 4 is desirable for reliable results. A ratio of 6.030 shows a suitable signal hence, this model can be used to navigate the design space. The model Prob > F (0.01) and F value (3.43) indicates significance of the model and specify that only 0.1% chance that model F value equivalent to this would occur due to noise (Tables III, IV).

Table III. ANOVA for response surface quadratic model (CCD).

Source	Sum of squares	df	Mean square	F value	p value	
Model	44859.50	20	2242.97	3.47	0.001	significant
A-Temp	45.97	1	45.97	0.0711	0.002*	
B-pH	5773.77	1	5773.77	8.93	0.005*	
C-inoculum size	674.26	1	674.26	1.04	0.315	
D-incubation	1669.32	1	1669.32	2.58	0.01*	
E-shaking int.	859.50	1	859.50	1.33	0.25	
AB	504.03	1	504.03	0.7799	0.3844	
AC	108.78	1	108.78	0.1683	0.6846	
AD	306.28	1	306.28	0.4739	0.4967	
AE	1287.78	1	1287.78	1.99	0.1687	
BC	639.03	1	639.03	0.9888	0.3283	
BD	30.03	1	30.03	0.0465	0.8308	
BE	0.2812	1	0.2812	0.0004	0.9835	
CD	1498.78	1	1498.78	2.32	0.1386	
CE	81.28	1	81.28	0.1258	0.7254	
DE	116.28	1	116.28	0.1799	0.6746	
A ²	14684.04	1	14684.04	22.72	< 0.0001	
B ²	4778.35	1	4778.35	7.39	0.0109	
C ²	14052.24	1	14052.24	21.74	< 0.0001	
D ²	8866.73	1	8866.73	13.72	0.0009	
E ²	762.63	1	762.63	1.18	0.2863	
Residual	18741.32	29	646.25			
Lack of fit	13833.82	22	628.81	0.8969	0.6112	not significant
Pure error	4907.50	7	701.07			
Cor total	63600.82	49				

* Statistically significant at 99% probability level; ^a df, degrees of freedom; ^b F-value, F distribution showing variation between sample means.

Table IV. ANOVA for regression analysis.

Std. Dev.	25.42	R ²	0.7053
Mean	63.06	Adjusted R ²	0.721
C.V. %	40.31	Predicted R ²	0.769
		Adeq precision	6.0298

Partial purification and optimization of enzyme activity

The optimum temperature of *B. subtilis* phytase activity was found to be 40°C, while the enzyme remains stable until 70°C. The partially purified phytase exhibited

stability at acidic pH, while the optimal enzymatic activity was found between pH 6.5-7.0 (enzyme activity of 267U/mg) and completely inactive above 9.0 (Figs. 3, 4).




Fig. 3. One factor plots for the effect of temperature (A) and pH (B) on phytase production.




Fig. 4. Lineweaver Burk plot drawn to calculate V_{max} and K_m values.

The residual activity of phytase was assayed following pre-incubation with different metal ions. Few metal ions after incubation were tested to have non inhibitory effect like Mn^{2+} and Cu^{2+} with residual activities as 185.8, 215.21 U/mL, respectively, where as some had a slight inhibitory effect like Ca^{2+} (27.12% residual activity), Co^{2+} (32%), Zn^{2+} (57.31%), Hg^{2+} (14.76% residual activity) Fe^{2+} (13.54%) of its original activity, respectively.




Fig. 5. SDS-PAGE Lane 1: Protein ladder (PAGE-MASTER Protein Standard) Lane 2: Partially Purified Phytase.

Purified phytase

The phytase was partially purified using ammonium persulfate precipitation and Sephadex G-50 column via chromatography. The purification of phytase increased approximately 87-folds as compared to crude extract with a 46.27 % yield. Purified phytase showed a high specific activity of 408.2 U/mg protein. Production of extracellular phytase in *B. subtilis* was marked with approximately 87 folds purification and 407 phytase U/mg of total protein (Table V).

The molecular weight of phytase obtained from Sephadex G- 50 column determined by SDS-PAGE was 30 kDa (Fig. 5). A single band was observed during the SDS-PAGE hence the assumption that *B. subtilis* phytase consists of identical subunits.

Table V. Purification of phytase from *Bacillus subtilis* KT004404.

Steps	^a Total activity (U/mL)	^b Total protein (mg)	^c Specific activity (U/mg)	^d Yield (%)	^e Purification (fold)
Enzyme production before RSM ^f	45.3	4.8	9.4	-	1
Crude extract of enzyme (after RSM)	132.32	6.3	21.00	100	2.3
Ammonium sulfate precipitation	78.34	1.7	46.08	59.20	2.19
Elusion Chromatography Sephadex G-50	61.23	0.4	153.07	46.27	7.2

^aTotal activity or phytase activity as the amount of phytase which can liberate the phosphorus from catalyzed phytate in one minute; ^bTotal protein content determined by Lowry method using BSA as standard; ^cSpecific activity = Total activity/ Total protein; ^dYield: Recovered enzyme activity; ^ePurification fold: Increase in specific activity; ^fMedia parameters (35 °C, pH-7.0, 100rpm, Na-phytate as substrate).

Kinetics of enzyme activity

Kinetic parameters of phytase enzyme were studied at varying Na-phytate concentrations. The optimal reaction velocity (V_{max}) and Michaelis constant (K_m) of *B. subtilis* phytase on sodium phytate was calculated via Lineweaver-Burk plot 250 U/mL of protein and were 0.175 mM of sodium phytate, respectively. K_m value is significantly higher than those reported for phytase from *A. vacuum* – 40 μ mol but lower than 1.77mM in *Geobacillus* sp., and 2.19 mM *B. subtilis*. Value of K_m is sensitive towards conditions such as pH, temperature, and ionic strength. The phytase activity reaches saturation in presence of a small amount of substrate concentration if the K_m value is small while a large K_m demonstrates requirement of higher sodium phytate concentrations to reach V_{max} . When binding site of enzyme is saturated with substrate it demonstrates the maximum velocity of the specific reactions.

DISCUSSION

A large number of studies have reported the phytase production from different microbial species with potential use in poultry feed (Oh *et al.*, 2001; Zhang *et al.*, 2011; Jorquera *et al.*, 2011; Singh and Satyanarayana, 2015). Kumari *et al.* (2016) reported the production of phytase from thermophilic mould *Sporotrichum thermophile* as a potential candidate to be used in poultry feed. B propeller phytase from *Bacillus* spp. isolated from high temperature habitat has also been reported to have a significant importance in the feed industry (Choi *et al.*, 2001; Tye *et al.*, 2002; Wulandari *et al.*, 2015). These studies support the idea of optimizing the production of phytase from thermostable *B. subtilis* KT004404 for large scale enzyme production to be used as poultry feed supplement (Powar and Jagannathan, 1982; Shimizu, 1992; Brown *et al.*, 2011; Singh and Satyanarayana, 2015). Bacterial production of enzyme during submerged fermentation has multiple benefits such as easy growth, fast replication, and easier genetic manipulation for production of recombinant enzymes as compared to other sources such as plants,

fungi and animals (Yang *et al.*, 2017). *B. subtilis* is also reported to be benign as it doesn't possess the pathogenic or toxigenic traits which makes it as a better candidate to be used as source of exogenous enzyme (Mingmongkolchai and Panbangred, 2018). Although many studies have reported the use of bacterial and fungal species for phytase production with potential industrial application, however, this is the first study which has reported the use of statistical model (RSM) for optimization of parameters for phytase production from *B. subtilis* KT004404.

The amount of phytase production obtained through statistical model (response surface methodology) was 132.32 U/mL at 35 °C, pH 6.5, inoculum density of 2.5mL after incubating for 24h at 75 rpm. Multiple studies have utilized the centrally composite design (CCD) of response surface methodology (RSM) for optimization of production parameters for exogenous enzymes through different microbial resources to determine the interaction and impacts of various production parameters (Vohra and Satyanarayana, 2003; Pable *et al.*, 2019; Kumari and Bansal, 2021). Thermostable enzymes have attracted profound interest in feed industry due to high temperature treatment during feed preparation and pelleting which affects the catalytic activity of exogenous enzymes (Vasudevan *et al.*, 2019).

Enzymes that exhibit catalytic stability at acidic pH are of significant importance in poultry due to acidic pH of gizzard (pH=3.4) and intestine (pH=6.4) (Svihus, 2011). The phytase produced from *B. subtilis* KT004404 reported during the current studies exhibited stable catalyzing potential between 4.0-7.0 pH due to which it is assumed that acidic pH of digestive tract of poultry will not be a limiting factor for in vivo phytase activity (Fengying *et al.*, 2011).

In the current studies the optimal phytase production was found in the medium containing wheat bran as principal carbon source (Kumari and Bansal, 2021). These results are in agreement with previously conducted studies in which the wheat bran is reported to be an efficient source of carbon for bacterial metabolism for exogenous

enzyme production (Wang *et al.*, 2011; Gupta *et al.*, 2014; Shahryari *et al.*, 2018). Bala *et al.* (2014) reported highest phytase yield when wheat bran (source of carbon), and ammonium persulfate (as a nitrogen source) were added in the fermentation medium during SSF using *H. nigrescens* BJ82. Several other studies also reported ammonium persulfate as suitable nitrogen source medium (Grothe *et al.*, 1999; Bajaj and Singh, 2010). A major advantage of using the agricultural residues as compared to expensive synthetic media is a significant reduction in production cost for enzyme production at commercial scale specially in agricultural country such as Pakistan (Jabeen *et al.*, 2019). Temperature is reported to be one of the most crucial factor for bacterial propagation and to maintain catalytic activity of enzymes (Vohra and Satyanarayana, 2003; Ajith *et al.*, 2019). In the present study, the optimal enzymatic activity of phytase produced by *B. subtilis* KT004404 was achieved at 35 °C. Our findings are in agreement with number of studies which have reported the optimal temperature for phytase activity produced from different microbial sources within temperatures range from 33–45 °C (Quan *et al.*, 2001; Gunashree and Venkateswaran, 2008; Gaind and Singh, 2015; Sanni *et al.*, 2019).

Demir *et al.* (2018) characterized 43.5 kDa phytase from *Lactobacillus coryniformis* MH121153 which exhibited optimal activity at 5.0 pH and 60 °C. Choi *et al.* (2001) reported the isolation of 44 kDa phytase from *Bacillus* sp. KHU-10 with optimal activity at 40 °C and optimal pH between 6.5–8.5 (Choi *et al.*, 2001). Similarly, Fujiti *et al.* (2003) also reported the optimal pH for microbial phytase between 4.5–5.5 at 50 °C. The effect of different metal ions on the enzyme activity depends on the microbial species. The current study shows that phytase activity was declined in presence of metal ions. Similar results have been reported in numerous previously conducted studies (Roy *et al.*, 2012; Yu and Chen, 2013). Phytase V_{max} values reported in the literature for *A. vacuum*. (55.9 nmol s⁻¹), *Rhizopus oligosporus* 32 nmol s⁻¹, *Bacillus licheniformis* 4.7mM *A. niger* 54.35 U/mL and *Geobacillus* sp. TF16 526.28 U/mg (Casey and Walsh, 2003; Chadha *et al.*, 2004; Ekren and Metin, 2013; Fasimoye *et al.*, 2014; Dokuzparmak *et al.*, 2017) support values obtained during current studies ($K_m = 0.175$ mM and $V_{max} = 250$ U/mL).

Plant based feed stuff such as soybean, corn, and rapeseed contain a high proportion of phytic acid. Poultry birds are unable to indigenously catalyze the phytic acid to release phosphorous (Abbasi *et al.*, 2019). Therefore, farmers have to rely on expensive inorganic phosphorous source to meet the phosphorus requirement of poultry which results in increased cost of feeding in poultry sector

(Woyengo and Nyachoti, 2011). Several studies have reported positive impacts of using exogenous phytase supplements on the growth performance during poultry trials (Cowieson and Adeola, 2005; Selle *et al.*, 2009; Ravindran, 2013). Due to the ability of phytase produced from *B. subtilis* KT004404 to retain catalytic activity within a wide range of pH and temperature, it offers a potential use in the feed industry for improving the nutrient utilization and growth performance of poultry birds.

CONCLUSION

Bacillus subtilis KT004404 has been proven to be a potential candidate for producing exogenous phytase. As compared to OFAT, the RSM based CCD model resulted in 2.3-fold increase in phytase production in the optimized culture media. The results of the current study indicated that the temperature, pH and concentration of wheat bran are significantly important parameters for producing bacterial phytase. Due to the ability of phytase to withstand temperature and acidic pH condition, it appears to have a potential use in poultry industry for improving nutritional properties of feed.

ACKNOWLEDGEMENTS

We like to acknowledge Prof. Dr. Abdul Hameed (late) for his support and guidance during this study. We also like to acknowledge HEC for indigenous PhD scholarships that helped in execution of this study (HEC-PIN no: 315-2692-2BS3-032). Higher Education Commission Pakistan played no direct role in collection, analysis, interpretation and publication of this work.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Abbasi, F., Fakhur-un-Nisa, T., Liu, J., Luo, X., and Abbasi, I.H.R., 2019. Low digestibility of phytate phosphorus, their impacts on the environment, and phytase opportunity in the poultry industry. *Environ. Sci. Pollut. Res.*, **26**: 9469-9479. <https://doi.org/10.1007/s11356-018-4000-0>
- Ajith, S., Ghosh, J., Shet, D., ShreeVidhya, S., Punith, B., and Elangovan, A., 2019. Partial purification and characterization of phytase from *Aspergillus foetidus* MTCC 11682. *AMB Express*, **9**: 1–11. <https://doi.org/10.1186/s13568-018-0725-x>
- Alias, N., Shunmugam, S., and Ong, P.Y., 2017. Isolation and molecular characterization of

- phytase producing bacteria from Malaysia hot springs. *J. Fundam. appl. Sci.*, **9**: 852-865. <https://doi.org/10.4314/jfas.v9i2s.53>
- Al-Kandari, S., Bhatti, M.E., Aldughpassi, A., Al-Hassawi, F., Al-Foudari, M., and Sidhu, J.S., 2021. Development of functional foods using psyllium husk and wheat bran fractions: Phytic acid contents. *Saudi J. biol. Sci.*, **10**: 220-223. <https://doi.org/10.1016/j.sjbs.2021.03.037>
- Bajaj, B.K., and Singh, N.P., 2010. Production of xylanase from an alkalitolerant *Streptomyces* sp. 7b under solid-state fermentation, its purification, and characterization. *Appl. Biochem. Biotechnol.*, **162**: 1804-1818. <https://doi.org/10.1007/s12010-010-8960-x>
- Bala, A., Sapna, J.J., Kumari, A., and Singh, B., 2014. Production of an extracellular phytase from a thermophilic mould *Humicola nigrescens* in solid state fermentation and its application in dephytinization. *Biocatal. Agric. Biotechnol.*, **3**: 259-264. <https://doi.org/10.1016/j.bcab.2014.07.002>
- Brown, M.E., Walker, M.C., Nakashige, T.G., Iavarone, A.T., and Chang, M.C., 2011. Discovery and characterization of heme enzymes from unsequenced bacteria: application to microbial lignin degradation. *J. Am. chem. Soc.*, **133**: 18006-18009. <https://doi.org/10.1021/ja203972q>
- Casey, A., and Walsh, G., 2003. Purification and characterization of extracellular phytase from *Aspergillus niger* ATCC 9142. *Bioresour. Technol.*, **86**: 183-188. [https://doi.org/10.1016/S0960-8524\(02\)00145-1](https://doi.org/10.1016/S0960-8524(02)00145-1)
- Chadha, B.S., Harmeet, G., Mandeep, M., Saini, H.S., and Singh, N., 2004. Phytase production by the thermophilic fungus *Rhizomucor pusillus*. *World J. Microbiol. Biotechnol.*, **20**: 105-109. <https://doi.org/10.1023/B:WIBI.0000013319.13348.0a>
- Choi, Y.M., Suh, H.J., and Kim, J.M., 2001. Purification and properties of extracellular phytase from *Bacillus* sp. KHU-10. *J. Protein Chem.*, **20**: 287-292. <https://doi.org/10.1023/A:1010945416862>
- Cowieson, A.J., and Adeola, O., 2005. Carbohydrases, protease, and phytase have an additive beneficial effect in nutritionally marginal diets for broiler chicks. *Poult. Sci.*, **84**: 1860-1867. <https://doi.org/10.1093/ps/84.12.1860>
- Cowieson, A., Acamovic, T., and Bedford, M., 2006. Phytic acid and phytase: Implications for protein utilization by poultry. *Poult. Sci.*, **85**: 878-885. <https://doi.org/10.1093/ps/85.5.878>
- Demir, Y., Dikbaş, N., and Beydemir, Ş., 2018. Purification and biochemical characterization of phytase enzyme from *Lactobacillus coryniformis* (MH121153). *Mol. Biotechnol.*, **60**: 783-790. <https://doi.org/10.1007/s12033-018-0116-1>
- Dokuzparmak, E., Sirin, Y., Cakmak, U., and Saglam, E.N., 2017. Purification and characterization of a novel thermostable phytase from the thermophilic *Geobacillus* sp. TF16. *Int. J. Fd. Prop.*, **20**: 1104-1116. <https://doi.org/10.1080/10942912.2016.1203930>
- Ekren, G.S., and Metin, K., 2013. Production, purification and characterization of phytase from phytase producing fungus. *Curr. Opin. Biotechnol.*, **24**: S81. <https://doi.org/10.1016/j.copbio.2013.05.234>
- Fasimoye, F.O., Olajuyigbe, F.M., and Sanni, M.D., 2014. Purification and characterization of a thermostable extracellular phytase from *Bacillus licheniformis* PFBL-03. *Prep. Biochem. Biotechnol.*, **44**: 193-205. <https://doi.org/10.1080/10826068.2013.812565>
- Fengying, G., Jie, G., Hui, R., and Guoqing, H., 2011. *In vitro* evaluating the activities and stabilities of the multihydrolases produced by *Aspergillus niger* Zju-Y1 in simulated poultry digestive tract pH levels. *Proc. Eng.*, **18**: 405-410. <https://doi.org/10.1016/j.proeng.2011.11.065>
- Fujita, J., Shigeta, S., Yamane, I., Fukuda, H., Kizaki, Y., Wakabayashi, S. and Ono, K., 2003. Production of two types of phytase from *Aspergillus oryzae* during industrial koji making. *J. Biosci. Bioeng.*, **95**: 460-465. [https://doi.org/10.1016/S1389-1723\(03\)80045-2](https://doi.org/10.1016/S1389-1723(03)80045-2)
- Gaind, S., and Singh, S., 2015. Production, purification and characterization of neutral phytase from thermotolerant *Aspergillus flavus* ITCC 6720. *Int. Biodeterior. Biodegrad.*, **99**: 15-22. <https://doi.org/10.1016/j.ibiod.2014.12.013>
- Grothe, E., Moo-Young, M., and Chisti, Y., 1999. Fermentation optimization for the production of poly (β -hydroxybutyric acid) microbial thermoplastic. *Enzyme Microb. Technol.*, **25**: 132-141. [https://doi.org/10.1016/S0141-0229\(99\)00023-X](https://doi.org/10.1016/S0141-0229(99)00023-X)
- Gunashree, B., and Venkateswaran, G., 2008. Effect of different cultural conditions for phytase production by *Aspergillus niger* CFR 335 in submerged and solid-state fermentations. *J. indust. Microbiol. Biotechnol.*, **35**: 1587. <https://doi.org/10.1007/s10295-008-0402-1>
- Gupta, R.K., Gangoliya, S.S., and Singh, N.K., 2014. Isolation of thermotolerant phytase producing fungi and optimisation of phytase production by *Aspergillus niger* NRF9 in solid state fermentation using response surface methodology. *Biotechnol. Bioprocess Eng.*, **19**: 996-1004. <https://doi.org/10.1007/s10295-008-0402-1>

- [org/10.1007/s12257-014-0175-5](https://doi.org/10.1007/s12257-014-0175-5)
- Jabeen, F., Muneer, B., and Qazi, J.I., 2019. Characterization of thermophilic bacteria *Anoxybacillus rupiensis* and cultivation in agroindustrial wastes isolated from hot spring in Chakwal, Pakistan. *Pakistan J. Zool.*, **51**: 1243-1250. <https://doi.org/10.17582/journal.pjz/2019.51.4.1243.1250>
- Jain, J., and Singh, B., 2017. Phytase production and development of an ideal dephytinization process for amelioration of food nutrition using microbial phytases. *Appl. Biochem. Biotechnol.*, **181**: 1485-1495. <https://doi.org/10.1007/s12010-016-2297-z>
- Jatuwong, K., Kumla, J., Suwannarach, N., Matsui, K., and Lumyong, S., 2020. Bioprocessing of agricultural residues as substrates and optimal conditions for phytase production of chestnut mushroom, *Pholiota adiposa*, in solid state fermentation. *J. Fungus.*, **6**: 384. <https://doi.org/10.3390/jof6040384>
- Jorquera, M.A., Crowley, D.E., Marschner, P., Greiner, R., Fernández, M.T., Romero, D., and De La Luz Mora, M., 2011. Identification of β -propeller phytase-encoding genes in culturable *Paenibacillus* and *Bacillus* spp. from the rhizosphere of pasture plants on volcanic soils. *FEMS Microbiol. Ecol.*, **75**: 163-172. <https://doi.org/10.1111/j.1574-6941.2010.00995.x>
- Kumar, V., Yadav, A., Saxena, A., Sangwan, P., and Dhaliwal, H., 2016. Unravelling rhizospheric diversity and potential of phytase producing microbes. *SM J. Biol.*, **2**: 1009. <https://www.jsmcentral.org>
- Kumar, V., Yadav, A.N., Verma, P., Sangwan, P., Saxena, A., Kumar, K., and Singh, B., 2017. β -Propeller phytases: diversity, catalytic attributes, current developments and potential biotechnological applications. *Int. J. Biol. Macromol.*, **98**: 595-609. <https://doi.org/10.1016/j.ijbiomac.2017.01.134>
- Kumari, A., Satyanarayana, T., and Singh, B., 2016. Mixed substrate fermentation for enhanced phytase production by thermophilic mould *Sporotrichum thermophile* and its application in beneficiation of poultry feed. *Appl. Biochem. Biotechnol.*, **178**: 197-210. <https://doi.org/10.1007/s12010-015-1868-8>
- Kumari, N., and Bansal, S., 2021. Statistical modeling and optimization of microbial phytase production towards utilization as a feed supplement. *Biomass Convers. Biorefin.*, pp. 1-11. <https://doi.org/10.1007/s13399-021-01672-x>
- Laboure, A.-M., Gagnon, J., and Lescure, A.M., 1993. Purification and characterization of a phytase (myo-inositol hexakisphosphate phosphohydrolase) accumulated in maize (*Zea mays*) seedlings during germination. *Biochem. J.*, **295**: 413-419. <https://doi.org/10.1042/bj2950413>
- Mingmongkolchai, S., and Panbangred, W., 2018. *Bacillus* probiotics: An alternative to antibiotics for livestock production. *J. appl. Microbiol.*, **124**: 1334-1346. <https://doi.org/10.1111/jam.13690>
- Murphy, J., and Riley, J., 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal. chim. Acta*, **27**: 31-36. [https://doi.org/10.1016/S0003-2670\(00\)88444-5](https://doi.org/10.1016/S0003-2670(00)88444-5)
- Naves, L. de P., Corrêa, A., Bertechini, A., Gomide, E., and dos Santos, C., 2012. Effect of pH and temperature on the activity of phytase products used in broiler nutrition. *Braz. J. Poult. Sci.*, **14**: 181-185. <https://doi.org/10.1590/S1516-635X2012000300004>
- Neira-Vielma, A.A., Aguilar, C.N., Ilyina, A., Contreras-Esquivel, J.C., das Graça Carneiro-da-Cunha, M., Michelena-Álvarez, G., and Martínez-Hernández, J.L., 2018. Purification and biochemical characterization of an *Aspergillus niger* phytase produced by solid-state fermentation using triticale residues as substrate. *Biotechnol. Rep.*, **17**: 49-54. <https://doi.org/10.1016/j.btre.2017.12.004>
- Oh, B.C., Chang, B.S., Park, K.H., Ha, N.C., Kim, H.K., Oh, B.H., and Oh, T.K., 2001. Calcium-dependent catalytic activity of a novel phytase from *Bacillus amyloliquefaciens* DS11. *Biochemistry*, **40**: 9669-9676. <https://doi.org/10.1021/bi010589u>
- Pable, A.A., Shah, S., Kumar, V.R., and Khire, J.M., 2019. Use of plackett-burman design for enhanced phytase production by *Williopsis saturnus* NCIM 3298 for applications in animal feed and ethanol production. *3 Biotech.*, **9**: 1-8. <https://doi.org/10.1007/s13205-019-1764-y>
- Powar, V.K., and Jagannathan, V., 1982. Purification and properties of phytate-specific phosphatase from *Bacillus subtilis*. *J. Bact.*, **151**: 1102-1108. <https://doi.org/10.1128/jb.151.3.1102-1108.1982>
- Quan, C., Zhang, L., Wang, Y., and Ohta, Y., 2001. Production of phytase in a low phosphate medium by a novel yeast *Candida krusei*. *J. Biosci. Bioeng.*, **92**: 154-160. [https://doi.org/10.1016/S1389-1723\(01\)80217-6](https://doi.org/10.1016/S1389-1723(01)80217-6)
- Ravindran, V., 2013. Feedenzymes: The science, practice, and metabolic realities. *J. appl. Poult. Res.*, **22**: 628-636. <https://doi.org/10.3382/japr.2013-00739>
- Roy, M.P., Poddar, M., Singh, K.K., and Ghosh, S., 2012. Purification, characterization and properties of phytase from *Shigella* sp. CD2. *Indian. J. Biochem. Biophys.*, **49**: 266-2671.
- Sadh, P.K., Duhan, S., and Duhan, J.S., 2018. Agro-

- industrial wastes and their utilization using solid state fermentation: A review. *Bioresour. Bioprocess*, **5**: 1-15. <https://doi.org/10.1186/s40643-017-0187-z>
- Sanni, D.M., Lawal, O.T., and Enujiugha, V.N., 2019. Purification and characterization of phytase from *Aspergillus fumigatus* isolated from African giant snail (*Achatina fulica*). *Biocatal. Agric. Biotechnol.*, **17**: 225–232. <https://doi.org/10.1016/j.bcab.2018.11.017>
- Shahryari, Z., Fazelipour, M.H., Setoodeh, P., Nair, R.B., Taherzadeh, M.J., and Ghasemi, Y., 2018. Utilization of wheat straw for fungal phytase production. *Int. J. Recycl. Org. Waste Agric.*, **7**: 345-355. <https://doi.org/10.1007/s40093-018-0220-z>
- Shimizu, M., 1992. Purification and characterization of phytase from *Bacillus subtilis* (natto) N-77. *Biosci. Biotechnol. Biochem.*, **56**: 1266–1269. <https://doi.org/10.1271/bbb.56.1266>
- Selle, P.H., Ravindran, V., and Partridge, G.G., 2009. Beneficial effects of xylanase and/or phytase inclusions on ileal amino acid digestibility, energy utilisation, mineral retention and growth performance in wheat-based broiler diets. *Anim. Feed Sci. Technol.*, **153**: 303-313. <https://doi.org/10.1016/j.anifeeds.2009.06.011>
- Singh, P., 2008. Significance of phytic acid and supplemental phytase in chicken nutrition: A review. *World's Poult. Sci. J.*, **64**: 553–580. <https://doi.org/10.1017/S0043933908000202>
- Singh, B., and Satyanarayana, T., 2015. Fungal phytases: characteristics and amelioration of nutritional quality and growth of non-ruminants. *J. Anim. Physiol. Anim. Nutr.*, **99**: 646-660. <https://doi.org/10.1111/jpn.12236>
- Svihus, B., 2011. The gizzard function, influence of diet structure and effects on nutrient availability. *World's Poult. Sci. J.*, **67**: 207-224. <https://doi.org/10.1017/S0043933911000249>
- Toor, G.S., Cade-Menun, B.J., and Sims, J.T., 2005. Establishing a linkage between phosphorus forms in dairy diets, feces, and manures. *J. environ. Qual.*, **34**: 1380–1391. <https://doi.org/10.2134/jeq2004.0232>
- Tye, A., Siu, F., Leung, T., and Lim, B., 2002. Molecular cloning and the biochemical characterization of two novel phytases from *B. subtilis* 168 and *B. licheniformis*. *Appl. Microbiol. Biotechnol.*, **59**: 190-197. <https://doi.org/10.1007/s00253-002-1033-5>
- Vats, P., and Banerjee, U.C., 2004. Production studies and catalytic properties of phytases (myo-inositolhexakisphosphate phosphohydrolases): An overview. *Enzyme Microb. Tech.*, **35**: 3–14. <https://doi.org/10.1016/j.enzmictec.2004.03.010>
- Vasudevan, U.M., Jaiswal, A.K., Krishna, S., and Pandey, A., 2019. Thermostable phytase in feed and fuel industries. *Bioresour. Technol.*, **278**: 400-407. <https://doi.org/10.1016/j.biortech.2019.01.065>
- Vohra, A., and Satyanarayana, T., 2003. Phytases: Microbial sources, production, purification, and potential biotechnological applications. *Crit. Rev. Biotechnol.*, **23**: 29–60. <https://doi.org/10.1080/713609297>
- Wang, Z.H., Dong, X.F., Zhang, G.Q., Tong, J.M., Zhang, Q., and Xu, S.Z., 2011. Waste vinegar residue as substrate for phytase production. *Waste Manage. Res.*, **29**: 1262-1270. <https://doi.org/10.1177/0734242X11398521>
- Waterborg, J.H., and Matthews, H.R., 1996. *The Lowry method for protein quantitation. The protein protocols handbook* (ed. J.M. Walker), Humana Press, Totowa, New Jersey. pp. 7–9. ISBN: 0-89603-338-4. https://doi.org/10.1007/978-1-60327-259-9_2
- Woyengo, T.A., and Nyachoti, C.M., 2011. Supplementation of phytase and carbohydrases to diets for poultry. *Can. J. Anim. Sci.*, **91**: 177-192. <https://doi.org/10.4141/cjas10081>
- Wulandari, R., Sari, E.N., Ratriyanto, A., Weldekiros, H., and Greiner, R., 2015. Phytase-producing bacteria from extreme regions in Indonesia. *Braz. Arch. Biol. Technol.*, **58**: 711-717. <https://doi.org/10.1590/S1516-89132015050173>
- Yanke, L.J., Bae, H.D., Selinger, L.B., and Cheng, K.J., 1998. Phytase activity of anaerobic ruminal bacteria. *Microbiol.*, **144**: 1565-1573. <https://doi.org/10.1099/00221287-144-6-1565>
- Yang, H., Li, J., Du, G., and Liu, L., 2017. Microbial production and molecular engineering of industrial enzymes: challenges and strategies. *Biotechnol. Microb. Eng.*, **12**: 151-165. <https://doi.org/10.1016/B978-0-12-803725-6.00006-6>
- Yu, P., and Chen, Y., 2013. Purification and characterization of a novel neutral and heat-tolerant phytase from a newly isolated strain *Bacillus nealsonii* ZJ0702. *BMC Biotechnol.*, **13**: 1-7. <https://doi.org/10.1186/1472-6750-13-78>
- Zhang, R., Yang, P., Huang, H., Yuan, T., Shi, P., Meng, K., and Yao, B., 2011. Molecular and biochemical characterization of a new alkaline β -propeller phytase from the insect symbiotic bacterium *Janthinobacterium* sp. TN115. *Appl. Microbiol. Biotechnol.*, **92**: 317-325. <https://doi.org/10.1007/s00253-011-3309-0>