



# Effect of Urbanization and Season on the Spatiotemporal Distribution of Mosquitoes in Bahawalpur, Pakistan

Robeela Shabbir<sup>1\*</sup>, Farkhanda Manzoor<sup>1</sup>, Muhammad Oneeb<sup>2</sup>, Furqan Awan<sup>2</sup> and Hania Ramzan<sup>1</sup>

<sup>1</sup>Department of Zoology, Lahore College for Women University (LCWU), Jail Road, Lahore, Pakistan

<sup>2</sup>Department of Parasitology, University of Veterinary and Animal Sciences (UVAS), Outfall Road, Lahore, Pakistan

## ABSTRACT

Mosquitoes are disease-causing vectors that carry a variety of pathogens. Their diversity and spatiotemporal distribution are highly affected by urbanization and seasons. In this study, the mosquitoes were collected from six tehsils (Hasilpur, Ahmedpur East, Khairpur Tamewali Yazman Bahawalpur City and Bahawalpur Saddar,) of district Bahawalpur from September 2020 to August 2021. All the mosquitoes were morphologically identified by using standard identification keys. The data was analyzed using the Shannon-Wiener diversity Index, one-way ANOVA, and unpaired t-test. The highest number of mosquitoes was found in tehsil Bahawalpur Saddar while the lowest number was observed in tehsil Ahmedpur East. Of all the three genera collected, the genus *Culex* was highly abundant (75.89%), followed by *Anopheles* (19.08%) and *Aedes* (5.03%). *Culex* and *Anopheles* mosquitoes were found in all the six tehsils of Bahawalpur while *Aedes* mosquitoes were found in Bahawalpur City and Bahawalpur Saddar, only. The number of mosquitoes was higher in rural areas as compared to urban areas of four tehsils of district Bahawalpur calculated through t test.

## Article Information

Received 24 July 2023

Revised 05 November 2023

Accepted 17 November 2023

Available online 18 December 2023 (early access)

## Authors' Contribution

FM and MO conceptualized and designed the study. RS performed field work and lab work and prepared the first draft. FA analyzed the data. HR reviewed and revised the manuscript. All authors approved the final manuscript to be submitted.

## Key words

Spatiotemporal, Urbanization, *Culex*, *Anopheles*, *Aedes*

## INTRODUCTION

Pakistan is a subtropical country that is rich in mosquito fauna (Imran *et al.*, 2022). Mosquitoes can cause deadly diseases like dengue (Knowlton *et al.*, 2009), malaria (Lawler and Lanzaro, 2005) West Nile fever (Styer *et al.*, 2011), Chikungunya (Higgs and Vanlandingham 2015), Japanese Encephalitis, etc. (Attaullah *et al.*, 2023).

Bahawalpur, a highly populated district of Pakistan, is also experiencing rapid mosquito-borne disease outbursts due to the high level of urbanization which causes mismanagement and deterioration in solid waste disposal (Mohsin and Chinyama, 2016). Due to rapid urbanization a change in anthropogenic landscape i.e., the development of parks, gardens, swimming pools, urban sewage systems, etc.

is occurring which is affecting the diversity, spatiotemporal distribution, and abundance of mosquitoes (Ferraguti *et al.*, 2016).

The influence of seasons on the breeding habitats of mosquitoes affects mosquito abundance. In the different seasons, temperature plays a key role. Temperature changes the period of maturity of adult mosquitoes (Tun-Lin *et al.*, 2000; Bayoh and Lindsay, 2004). Rainfall in different seasons directly affects their breeding sites (Ceccato *et al.*, 2005; Byun and Webb, 2012). The relative humidity is also one of the factors in the seasonal effects that regulate the oviposition and metabolism of adult mosquitoes (Ceccato *et al.*, 2005; Reiter, 2001). All these climatic variables are therefore responsible for controlling many mosquito-borne diseases. (Wongkoon *et al.*, 2013; Bashar and Tuno, 2014).

The present study aims to determine the specific patterns of diversity, abundance, and community composition of mosquitoes in different tehsils of district Bahawalpur. The effect of urbanization and seasons on the spatiotemporal distribution of mosquitoes is also investigated in this study. This study would be helpful in developing strategy for control of mosquito-borne diseases.

\* Corresponding author: robeela@yahoo.com  
0030-9923/2023/0001-0001 \$ 9.00/0



Copyright 2023 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## MATERIALS AND METHODS

### Study area

Sampling was done from district Bahawalpur (29.418° N and 71.670° E). The total area of this district is 24,830 km<sup>2</sup>. It is located 117 m above sea level. Administratively, it is divided into six tehsils; Bahawalpur City, Yazman, Bahawalpur Saddar, Hasilpur, Khairpur Tamewali, Ahmadpur East, and Khairpur. These tehsils are further divided into 78 rural and 29 urban areas. It is a highly populated district with a total population of 3,668,106. The population density of district Bahawalpur is 153.4 persons per km<sup>2</sup>. The climate is extremely dry and hot during summers and dry and cold during winters. The mean annual rainfall of Bahawalpur district is 169.8 mm. The rainfall usually occurs during the monsoon season i.e., July and August (<https://bahawalpur.org/tehsils/>).

### Collection of mosquitoes

Adult mosquitoes were collected from specific sites on monthly basis from September 2020 to August 2021 from inside the houses outside gutters, cattle sheds, lawns, graveyards, zoos, etc., from different tehsils of Bahawalpur. The adult mosquitoes were collected with a CDC sweeper and mechanical aspirators.

### Identification of mosquitoes

Mosquitoes were identified based on their morphological features using taxonomic keys (Christophers, 1933; Rueda, 2004; Barraud, 1934).

### Statistical analysis

The Shannon diversity index ( $H$ ) was applied to characterize species diversity in the six tehsils of district Bahawalpur. Shannon's index accounts for both the abundance and evenness of the species present (Strong, 2016). Adult mosquito abundance between the six tehsils was compared using a one-way analysis of variance test (ANOVA) where significant differences were observed in an ANOVA test, Tukey's post hoc analysis was used to separate the means. One-way ANOVA also helped to find the relationship between the abundance of mosquitoes with the season (Kim, 2017). The comparison of the abundance of mosquitoes between rural and urban areas of different tehsils of district Bahawalpur was analyzed through a student t-test (Kim, 2015).

## RESULTS

A total of 18,354 mosquitoes were collected from the six tehsils of Bahawalpur. Among all the three genera, the genus *Culex* was highly abundant (75.89%), followed by

*Anopheles* (19.08%) and *Aedes* (5.03%). Genus *Culex* and *Anopheles* were found in all six tehsils. However, *Aedes* mosquitoes were found in two tehsils only i.e., Bahawalpur city and Bahawalpur Saddar (Fig. 1). It was observed that the highest number of mosquitoes were found in the tehsil Bahawalpur Saddar while the lowest number of mosquitoes were found in tehsil Ahmedpur East. A total of five species from the genus *Culex* were identified i.e., *Culex quinquefasciatus* (30.49%), *Culex tritaeniorhynchus* (27.42%), *Culex pipiens* (11.83%), *Culex pseudovishnui* (4.23%), and *Culex vagans* (1.92%). Similarly, five species of the *Anopheles* genus were identified i.e., *Anopheles subpictus* (10.5%), *Anopheles culicifacies* (5.76%), *Anopheles stephensi* (2.8%), *Anopheles pulcherrimus* (1.72%), and *Anopheles nigerrimus* (0.87%). Only two species, *Aedes albopictus* (3.4%) and *Aedes aegypti* (1.6%) were identified from the genus *Aedes* (Fig. 2).




Fig. 1. Map of district Bahawalpur showing different tehsils.




Fig. 2. Total number of mosquitoes in different tehsils of Bahawalpur.

### Species diversity and abundance

In general, there were considerable differences between tehsils in terms of the diversity of species identified ( $n$ ), Shannon diversity ( $H$ ), and Simpson diversity ( $D$ ) (Table I).

**Table I. Diversity index of number of individuals of different species of mosquitoes collected from six tehsils of District Bahawalpur.**

Name of the species	Saddar		Bahawalpur city		Hasilpur		Khairpur		Yazman		Ahmed Pur Sharqia							
	% age	Shannon's index Pi*(logPi)	Simpson's index n(n-1)	% age	Shannon's index Pi*(logPi)	Simpson's index n(n-1)	% age	Shannon's index Pi*(logPi)	Simpson's index n(n-1)	% age	Shannon's index Pi*(logPi)	Simpson's index n(n-1)						
<i>Culex quinquefasciatus</i>	24	0.343	928332	26	0.349	546860	32	0.364	1264500	33	0.366	1116192	34	0.367	1007012	31	0.362	499142
<i>Culex tritaenorrhynchus</i>	28	0.356	1246572	29	0.358	683102	23	0.339	669942	25	0.344	598302	27	0.352	601400	31	0.363	516242
<i>Culex pseudovishnui</i>	5	0.149	39800	3	0.110	8372	5	0.148	29756	4	0.133	17030	4	0.119	10712	3	0.112	5700
<i>Culex Vagans</i>	2	0.079	6642	1.3	0.055	1260	2	0.081	5402	2	0.082	4422	2	0.078	3306	2	0.064	1190
<i>Culex pipiens</i>	11.6	0.249	214832	10	0.233	86730	13	0.265	211140	12	0.256	147840	11	0.246	107256	10	0.235	57360
<i>Anopheles subpictus</i>	6.8	0.183	73712	11	0.245	103362	11	0.246	158006	11	0.243	120756	11	0.241	99540	12	0.252	74256
<i>Anopheles stephensi</i>	1.8	0.071	4830	2	0.084	3906	4	0.124	17822	3	0.113	10920	3	0.100	6642	3	0.098	3782
<i>Anopheles culcifacies</i>	3.7	0.121	21462	5.5	0.159	24492	7	0.181	55932	6	0.176	40602	6	0.169	31152	6	0.168	18632
<i>Anopheles pulcherrimus</i>	1.	0.046	1560	1	0.058	1482	2	0.086	6320	2	0.079	4160	2	0.072	2652	2	0.070	1560
<i>Anopheles negirrimus</i>	0.5	0.026	380	0.8	0.037	462	1	0.049	1406	1	0.046	992	1	0.042	650	1	0.043	420
<i>Aedes albopictus</i>	10	0.236	174306	7	0.192	44732	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aedes aegypti</i>	5	0.158	46872	3	0.096	5700	0	0	0	0	0	0	0	0	0	0	0	0

**Table II. Abundance (Mean ± SD) of mosquitoes in different Tehsils of District Bahawalpur.**

Species	Bahawalpur	Hasilpur	Ahmadpur east	Khairpur	Yazman	Saddar	F	P
<i>C. quinquefasciatus</i>	738.67±17.04 <sup>c</sup>	1128.33±15.28 <sup>a</sup>	705±16.09	105.67±10.5 <sup>b</sup>	1005.3±18 <sup>c</sup>	963.33±12.01 <sup>d</sup>	394.56	0.001
<i>C. tritaenorrhynchus</i>	828.67±16.56 <sup>b</sup>	816±14.73 <sup>c</sup>	717.67±17.04	773.7±17.5 <sup>c</sup>	774±17.09 <sup>c</sup>	1118.7±17.6 <sup>a</sup>	217.62	0.001
<i>C. pseudovishnui</i>	91.33±17.01 <sup>d</sup>	171.67±15.04 <sup>ab</sup>	75.67±11.5	134±15.72 <sup>bc</sup>	102±13.11 <sup>cd</sup>	201.67±14.57 <sup>a</sup>	34.1	0.001
<i>C. pipiens</i>	293±13.11 <sup>c</sup>	460±12 <sup>a</sup>	238.67±16.04 <sup>b</sup>	383.67±13.05 <sup>b</sup>	327.67±11.5 <sup>c</sup>	464±11 <sup>a</sup>	150.3	0.001
<i>C. vagans</i>	35.67±2.52 <sup>c</sup>	74.33±14.5 <sup>b</sup>	35±2 <sup>c</sup>	65.67±5.13 <sup>ab</sup>	58.33±3.51 <sup>b</sup>	82.67±3.06 <sup>a</sup>	26.26	0.001
<i>A. subpictus</i>	324±17.09 <sup>b</sup>	396.3±17.6 <sup>a</sup>	273±14 <sup>c</sup>	346.67±12.06 <sup>b</sup>	316.33±16.5 <sup>b</sup>	272.33±13.5 <sup>c</sup>	28.45	0.001
<i>A. stephensi</i>	64.33±5.13 <sup>d</sup>	134.67±5.03 <sup>a</sup>	64 <sup>d</sup> ±5.29	103.33±7.64 <sup>b</sup>	84±5.29 <sup>c</sup>	17.33±4.51 <sup>cd</sup>	74.73	0.001
<i>A. culcifacies</i>	159.67±16.17 <sup>bc</sup>	237.3±19.5 <sup>a</sup>	137.67±17.01 <sup>c</sup>	201.67±10.5 <sup>ab</sup>	178.67±13.58 <sup>bc</sup>	146.67±14.5 <sup>c</sup>	17.6	0.001
<i>A. pulcherrimus</i>	39.33±1.52 <sup>d</sup>	80.67±3.06 <sup>a</sup>	39.67 <sup>d</sup> ±1.52	64.33±3.06 <sup>b</sup>	53.33±4.16 <sup>c</sup>	40.33±2.52 <sup>d</sup>	108.69	0.001
<i>A. negirrimus</i>	21.67±1.52 <sup>c</sup>	37.67±2.52 <sup>a</sup>	21±1 <sup>bc</sup>	32.67±2.08 <sup>a</sup>	26±3 <sup>b</sup>	20±1 <sup>c</sup>	38.82	0.001
<i>A. albopictus</i>	217.3±30.4 <sup>b</sup>	0±0 <sup>c</sup>	0±0 <sup>c</sup>	0±0 <sup>c</sup>	0±0 <sup>c</sup>	419.3±30 <sup>a</sup>	307.17	0.001
<i>A. aegypti</i>	77.33±15.04 <sup>a</sup>	0±0 <sup>c</sup>	0±0 <sup>c</sup>	0±0 <sup>c</sup>	0±0 <sup>c</sup>	214.33±16.17 <sup>a</sup>	278.61	0.001

The Shannon and Simpson diversity indices for the six tehsils, respectively, ranged from 2.01 to 1.76 and 0.82 to 0.77. The tehsil Saddar has the greatest and tehsil Ahmedpur East had the lowest values for both indices, respectively. Significant variations in mosquito abundance were found in all of the tehsils after analysis of variance (ANOVA) tests were performed. Hasilpur had the highest concentration of *Culex quinquefasciatus* ( $F = 394.56$ ,  $df = 5$ ,  $P < 0.001$ ), followed by Khairpur Tamewali, Yazman, Saddar, Bahawalpur, and Ahmedpur East with the lowest concentration (Table II). *Aedes aegypti* was a rare species in this district that was absent from Hasilpur, Khairpur Tamewali, Yazman, and Ahmedpur East tehsils but highly dispersed in Bahawalpur and Saddar tehsils ( $F = 278.61$ ,  $df = 5$ ,  $P < 0.001$ ). In the other tehsils of Bahawalpur, every species displayed a similar trend of abundance.

#### Relationship of the abundance of mosquitoes with seasons

Interesting findings were obtained via a one-way ANOVA on the seasonal abundance of mosquitoes. The research revealed that each tehsil's mosquito population was at its highest during the monsoon season, followed by fall, and at its lowest during the winter. In comparison to the summer, there were more mosquitoes in the spring. The Hasilpur tehsil had the greatest *Culex quinquefasciatus* count ( $F = 85.81$ ,  $df = 4$ ,  $P = 0.001$ ) (Table III). Its peak levels of abundance were noted in the autumn (Mean = 299,  $SD = 15.72$ ) and monsoon seasons (Mean = 299.33,  $SD = 12.06$ ), with a sharp decline in abundance throughout the winter (Mean = 125,  $SD = 15.01$ ). In the spring, it increases once more (Mean = 237,  $SD = 17$ ), and in the summer, it decreases (Mean = 160,  $SD = 14.1$ ) (Table III). *Anopheles subpictus* of the genus *Anopheles* displayed the same

**Table III. Relationship of abundance of species of mosquitoes with seasons in Tehsil Hasilpur of District Bahawalpur.**

Species	Mean $\pm$ Standard Deviation					F	P
	Autumn	Winter	Spring	Summer	Monsoon		
<i>Culex quinquefasciatus</i>	299 $\pm$ 15.72 <sup>a</sup>	125 $\pm$ 15.01 <sup>c</sup>	237 $\pm$ 17 <sup>b</sup>	160 $\pm$ 14.11 <sup>c</sup>	299.33 $\pm$ 12.06 <sup>a</sup>	85.81	0.001
<i>Culex tritaenorrhynchus</i>	243.67 $\pm$ 11.02 <sup>a</sup>	79.3 $\pm$ 18.5 <sup>c</sup>	149 $\pm$ 14 <sup>b</sup>	109.33 $\pm$ 13.01 <sup>c</sup>	238.67 $\pm$ 11.02 <sup>a</sup>	88.01	0.001
<i>Culex pseudovishnui</i>	46 $\pm$ 3 <sup>b</sup>	15 $\pm$ 3 <sup>c</sup>	32.33 $\pm$ 2.52 <sup>b</sup> <sup>c</sup>	26.66 $\pm$ 1.52 <sup>c</sup>	54.33 $\pm$ 15.04 <sup>a</sup>	14.4	0.001
<i>Culex pipiens</i>	120 $\pm$ 16.52 <sup>a</sup>	37.67 $\pm$ 2.52 <sup>b</sup>	103.33 $\pm$ 13.5 <sup>a</sup>	64 $\pm$ 17.09 <sup>b</sup>	133 $\pm$ 12.53 <sup>a</sup>	26.06	0.001
<i>Culex vagans</i>	20.66 $\pm$ 1.52 <sup>a</sup>	6 $\pm$ 1 <sup>c</sup>	10.33 $\pm$ 2.08 <sup>b</sup> <sup>c</sup>	11.33 $\pm$ 1.52 <sup>b</sup>	25.33 $\pm$ 2.52 <sup>a</sup>	58.49	0.001
<i>Anopheles subpictus</i>	141.33 $\pm$ 14.05 <sup>a</sup>	22 $\pm$ 3 <sup>b</sup>	116 $\pm$ 16 <sup>a</sup>	0 <sup>b</sup>	119.7 $\pm$ 17.5 <sup>a</sup>	79.98	0.001
<i>Anopheles stephensi</i>	45.67 $\pm$ 2.52 <sup>a</sup>	9.76 $\pm$ 2.08 <sup>c</sup>	38.33 $\pm$ 1.52 <sup>b</sup>	0 <sup>b</sup>	41.67 $\pm$ 2.08 <sup>a</sup> <sup>b</sup>	372.42	0.001
<i>Anopheles culicifacies</i>	67.3 $\pm$ 18 <sup>a</sup>	7 $\pm$ 1 <sup>b</sup>	61.7 $\pm$ 17.5 <sup>a</sup>	0 <sup>b</sup>	66.67 $\pm$ 15.5 <sup>a</sup>	19.82	0.001
<i>Anopheles pulcherrimus</i>	19.33 $\pm$ 1.52 <sup>a</sup> <sup>b</sup>	6.33 $\pm$ 1.52 <sup>c</sup>	17.33 $\pm$ 2.52 <sup>b</sup>	0 <sup>b</sup>	23.67 $\pm$ 2.08 <sup>a</sup>	94.29	0.001
<i>Anophels nigerrimus</i>	11.66 $\pm$ 1.52 <sup>a</sup>	2.67 $\pm$ 2.08 <sup>c</sup>	6.3 $\pm$ 1.52 <sup>b</sup>	0 <sup>c</sup>	11.33 $\pm$ 2.08 <sup>a</sup>	30.09	0.001
<i>Aedes albopictus</i>	0	0	0	0	0	0	0
<i>Aedes aegypti</i>	0	0	0	0	0	0	0

**Table IV. Relationship of abundance of species of mosquitoes with seasons in Tehsil Saddar of District Bahawalpur.**

Species	Mean $\pm$ Standard deviation					F	P
	Autumn	Winter	Spring	Summer	Monsoon		
<i>Culex quinquefasciatus</i>	295.67 $\pm$ 14.57 <sup>a</sup>	80 $\pm$ 20 <sup>c</sup>	139.6 $\pm$ 16.17 <sup>b</sup>	167.3 $\pm$ 12.01 <sup>b</sup>	283.33 $\pm$ 15.57 <sup>a</sup>	104.06	0.001
<i>Culex tritaenorrhynchus</i>	320 $\pm$ 17 <sup>a</sup>	98.67 $\pm$ 13.01 <sup>c</sup>	171.6 $\pm$ 14.01 <sup>b</sup>	200.6 $\pm$ 17.04 <sup>b</sup>	325.67 $\pm$ 14.5 <sup>a</sup>	125.1	0.001
<i>Culex pseudovishnui</i>	48 $\pm$ 15.72 <sup>b</sup>	11 $\pm$ 3 <sup>d</sup>	36.67 $\pm$ 2.08 <sup>b</sup> <sup>c</sup>	19.33 $\pm$ 2.52 <sup>d</sup>	81.67 $\pm$ 13.05 <sup>a</sup>	26.41	0.001
<i>Culex pipiens</i>	138.6 $\pm$ 12.01 <sup>a</sup>	35 $\pm$ 3 <sup>c</sup>	69.67 $\pm$ 14.5 <sup>b</sup>	74 $\pm$ 14.53 <sup>b</sup>	149.67 $\pm$ 13.58 <sup>a</sup>	47.26	0.001
<i>Culex vagans</i>	15 $\pm$ 3 <sup>b</sup> <sup>c</sup>	2.67 $\pm$ 2.08 <sup>d</sup>	16.66 $\pm$ 1.52 <sup>b</sup>	10.33 $\pm$ 2.08 <sup>c</sup>	37.33 $\pm$ 2.52 <sup>a</sup>	94.79	0.001
<i>Anopheles subpictus</i>	87.33 $\pm$ 16.01 <sup>a</sup>	16.67 $\pm$ 2.52 <sup>b</sup>	78 $\pm$ 14.73	0 <sup>b</sup>	93 $\pm$ 18.5 <sup>a</sup>	34.25	0.001
<i>Anopheles stephensi</i>	19 $\pm$ 3 <sup>b</sup>	5 $\pm$ 1 <sup>c</sup>	21 $\pm$ 1 <sup>a</sup> <sup>b</sup>	0 <sup>d</sup>	24.33 $\pm$ 2 <sup>a</sup>	11.93	0.001
<i>Anopheles culicifacies</i>	52.33 $\pm$ 13.05 <sup>a</sup>	10.33 $\pm$ 1.52 <sup>b</sup>	37 $\pm$ 3 <sup>a</sup>	0 <sup>b</sup>	48.66 $\pm$ 1.52 <sup>a</sup>	44.48	0.001
<i>Anopheles pulcherrimus</i>	10.33 $\pm$ 1.52 <sup>a</sup>	3.33 $\pm$ 1.52 <sup>b</sup>	13.66 $\pm$ 1.52 <sup>a</sup>	0 <sup>b</sup>	12.67 $\pm$ 2.52 <sup>a</sup>	40.81	0.001
<i>Anopheles negirrimus</i>	5 $\pm$ 1 <sup>b</sup>	2 $\pm$ 1 <sup>c</sup> <sup>d</sup>	4 $\pm$ 1 <sup>b</sup> <sup>c</sup>	0 <sup>d</sup>	9.33 $\pm$ 1.52 <sup>a</sup>	34.75	0.001
<i>Aedes albopictus</i>	146.6 $\pm$ 16.17 <sup>a</sup>	16.33 $\pm$ 2.52 <sup>c</sup>	71 $\pm$ 15.52 <sup>b</sup>	62.33 $\pm$ 15.01 <sup>b</sup>	125 $\pm$ 14.11 <sup>a</sup>	43.51	0.001
<i>Aedes aegypti</i>	72.33 $\pm$ 14.5 <sup>a</sup>	9 $\pm$ 1 <sup>c</sup>	35.67 $\pm$ 2.52 <sup>b</sup>	26.33 $\pm$ 2.08 <sup>b</sup> <sup>c</sup>	73 $\pm$ 13 <sup>a</sup>	31.15	0.001

**Table V. Relationship of abundance of species of mosquitoes with seasons in Tehsil Khairpur East of District Bahawalpur.**

Species	Mean ± Standard deviation					F	P
	Autumn	Winter	Spring	Summer	Monsoon		
<i>Culex quinquefasciatus</i>	289.67±13.58 <sup>a</sup>	109.67±15.18 <sup>d</sup>	222.67±10.5 <sup>b</sup>	152.67±13.01 <sup>c</sup>	284.67±16.62 <sup>a</sup>	98.2	0.001
<i>Culex tritaenorrhynchus</i>	226.3±19.5 <sup>a</sup>	71.67±15.57 <sup>c</sup>	152.67±17.01 <sup>b</sup>	99±14 <sup>c</sup>	226.67±15.5 <sup>a</sup>	56.49	0.001
<i>Culex pseudovishnui</i>	36.33±2.08 <sup>a</sup>	13.66±1.52 <sup>c</sup>	27.33±2.52 <sup>b</sup>	18.67±2.08 <sup>c</sup>	35.67±2.08 <sup>a</sup>	70.5	0.001
<i>Culex pipiens</i>	102.33±15.5 <sup>a</sup>	32.66±1.52 <sup>c</sup>	84.67±13.65 <sup>a,b</sup>	51.67±15.28 <sup>b,c</sup>	108.3±17.5 <sup>a</sup>	16.58	0.001
<i>Culex vagans</i>	16.66±1.52 <sup>b</sup>	5.33±1.52 <sup>d</sup>	11±2 <sup>c</sup>	10.33±2.08 <sup>c,d</sup>	23±3 <sup>a</sup>	31.19	0.001
<i>Anopheles subpictus</i>	120±18 <sup>a</sup>	18.67±2.08 <sup>b</sup>	101±16 <sup>a</sup>	0 <sup>b</sup>	107.67±16.04 <sup>a</sup>	55.31	0.001
<i>Anopheles stephensi</i>	33±3 <sup>a</sup>	8.67±2.08 <sup>b</sup>	30±1 <sup>a</sup>	0 <sup>c</sup>	34.67±2.08 <sup>a</sup>	201.8	0.001
<i>Anopheles culicifacies</i>	67.3±18 <sup>a</sup>	7±1 <sup>b</sup>	61.7±17.5 <sup>a</sup>	0 <sup>b</sup>	66.67±15.5 <sup>a</sup>	19.82	0.001
<i>Anopheles pulcherrimus</i>	19.33±1.52 <sup>a,b</sup>	6.33±1.52 <sup>c</sup>	17.33±2.52 <sup>b</sup>	0 <sup>d</sup>	23.67±2.08 <sup>a</sup>	94.29	0.001
<i>Anopheles nigerrimus</i>	11.66±1.52 <sup>a</sup>	2.67±2.08 <sup>b,c</sup>	6.33±1.52 <sup>b</sup>	0 <sup>c</sup>	11.33±2.08 <sup>a</sup>	30.09	0.001
<i>Aedes albopictus</i>	0	0	0	0	0	0	0
<i>Aedes aegypti</i>	0	0	0	0	0	0	0

**Table VI. Relationship of abundance of species of mosquitoes with seasons in Tehsil Yazman of District Bahawalpur.**

Species	Mean ± Standard deviation					F	P
	Autumn	Winter	Spring	Summer	Monsoon		
<i>Culex quinquefasciatus</i>	282±12.53 <sup>a</sup>	89.3±17.5 <sup>d</sup>	216.67±16.5 <sup>b</sup>	136.7±19 <sup>c</sup>	277.33±14.57 <sup>a</sup>	83.74	0.001
<i>Culex tritaenorrhynchus</i>	211.67±16.01 <sup>a</sup>	59±11.53 <sup>c</sup>	183.67±13.58 <sup>a</sup>	108.3±18 <sup>b</sup>	59±13.5 <sup>c</sup>	66.25	0.001
<i>Culex pseudovishnui</i>	29.33±2.52 <sup>a</sup>	10.33±1.52 <sup>c</sup>	21.67±2.08 <sup>b</sup>	15.67±2.52 <sup>c</sup>	27.33±2.08 <sup>a,b</sup>	39.95	0.001
<i>Culex pipiens</i>	87.67±11.02 <sup>a</sup>	25.33±2.08 <sup>c</sup>	77.7±17.5 <sup>a,b</sup>	50±1 <sup>b,c</sup>	85.67±13.05 <sup>a</sup>	18.02	0.001
<i>Culex vagans</i>	14.33±2.52 <sup>b</sup>	5.67±2.08 <sup>c</sup>	10.33±1.52 <sup>b,c</sup>	10±1 <sup>b,c</sup>	19.33±1.52 <sup>a</sup>	24.36	0.001
<i>Anopheles subpictus</i>	107±14.53 <sup>a</sup>	14±1 <sup>b</sup>	94.33±12.5 <sup>a</sup>	0 <sup>b</sup>	99±18.5 <sup>a</sup>	55.79	0.001
<i>Anopheles stephensi</i>	45.67±2.52 <sup>a</sup>	9.67±2.08 <sup>c</sup>	88.33±1.52 <sup>b</sup>	0 <sup>d</sup>	41.67±2.08 <sup>b</sup>	372.42	0.001
<i>Anopheles culicifacies</i>	74±19 <sup>a</sup>	7±1 <sup>b</sup>	69.67±16.62 <sup>a</sup>	0 <sup>b</sup>	83±11.53 <sup>a</sup>	30.86	0.001
<i>Anopheles pulcherrimus</i>	27±3 <sup>a</sup>	9±1 <sup>c</sup>	19.33±2.52 <sup>b</sup>	0 <sup>d</sup>	24.33±2.08 <sup>a,b</sup>	91.91	0.001
<i>Anopheles nigerrimus</i>	15±1 <sup>a</sup>	3±1 <sup>c</sup>	8±1 <sup>b</sup>	0 <sup>c</sup>	12.33±2.52 <sup>a</sup>	62.77	0.001
<i>Aedes albopictus</i>	0	0	0	0	0	0	0
<i>Aedes aegypti</i>	0	0	0	0	0	0	0

**Table VII. Relationship of abundance of species of mosquitoes with seasons in Tehsil Ahmadpur East of District Bahawalpur.**

Species	Mean ± Standard deviation					F	P
	Autumn	Winter	Spring	Summer	Monsoon		
<i>Culex quinquefasciatus</i>	229.33±12.01 <sup>a</sup>	80±15.1 <sup>b</sup>	89.67±17.01 <sup>b</sup>	80.67±11.02 <sup>b</sup>	230±20 <sup>a</sup>	81.52	0.001
<i>Culex tritaenorrhynchus</i>	227±15.52 <sup>a</sup>	77±18 <sup>b</sup>	99.67±11.02 <sup>b</sup>	93.33±17.04 <sup>b</sup>	220.33±14.57 <sup>a</sup>	69.04	0.001
<i>Culex pseudovishnui</i>	23.66±1.52 <sup>a</sup>	9.33±1.52 <sup>b</sup>	7±1 <sup>b</sup>	9.67±2.08 <sup>b</sup>	27±3 <sup>a</sup>	67.72	0.001
<i>Culex pipiens</i>	77.33±10.5 <sup>a</sup>	26.33±2.52 <sup>b</sup>	30.67±2.08 <sup>b</sup>	29.66±1.52 <sup>b</sup>	77.33±16.5 <sup>a</sup>	26.79	0.001
<i>Culex vagans</i>	9.67±2.08 <sup>a</sup>	2±1 <sup>b</sup>	6.67±2.08 <sup>a</sup>	8.33±1.52 <sup>a</sup>	10.33±1.52 <sup>a</sup>	11.59	0.001
<i>Anopheles subpictus</i>	86.67±16.56 <sup>a</sup>	13±1 <sup>b</sup>	86.67±15.28 <sup>a</sup>	0	85.3±18.5 <sup>a</sup>	3.99	0.001
<i>Anopheles stephensi</i>	116.67±2.08 <sup>b</sup>	4±1 <sup>c</sup>	18±3 <sup>b</sup>	0 <sup>c</sup>	23.66±1.52 <sup>a</sup>	90.18	0.001
<i>Anopheles culicifacies</i>	43.33±2.08 <sup>a,b</sup>	5.67±2.08 <sup>c</sup>	40±2 <sup>b</sup>	0 <sup>d</sup>	47.67±2.52 <sup>a</sup>	403.9	0.001
<i>Anopheles pulcherrimus</i>	10±1 <sup>b</sup>	4.67±2.08 <sup>c</sup>	15±2 <sup>a</sup>	0 <sup>d</sup>	10.66±1.52 <sup>b</sup>	43.46	0.001
<i>Anopheles nigerrimus</i>	5.33±1.52 <sup>a</sup>	2±1 <sup>b</sup>	7±1 <sup>a</sup>	0 <sup>b</sup>	7±1 <sup>a</sup>	27.72	0.001
<i>Aedes albopictus</i>	0	0	0	0	0	0	0
<i>Aedes aegypti</i>	0	0	0	0	0	0	0

**Table VIII. Relationship of abundance of species of mosquitoes with seasons in Tehsil Bahawalpur of District Bahawalpur.**

Species	Mean $\pm$ Standard deviation					F	P
	Autumn	Winter	Spring	Summer	Monsoon		
<i>Culex quinquefasciatus</i>	227 $\pm$ 15.72 <sup>a</sup>	74 $\pm$ 15.52 <sup>b</sup>	104 $\pm$ 17.09 <sup>b</sup>	98 $\pm$ 12 <sup>b</sup>	234.33 $\pm$ 12.06 <sup>a</sup>	82.78	0.001
<i>Culex tritaenorrhynchus</i>	251 $\pm$ 15 <sup>a</sup>	108 $\pm$ 14 <sup>b</sup>	108 $\pm$ 14.73 <sup>b</sup>	103.33 $\pm$ 11.02 <sup>b</sup>	251 $\pm$ 15.1 <sup>a</sup>	95.29	0.001
<i>Culex pseudovishnui</i>	31.33 $\pm$ 2.52 <sup>a</sup>	13.67 $\pm$ 2.08 <sup>b</sup>	10 $\pm$ 1 <sup>b</sup>	11 $\pm$ 2 <sup>b</sup>	27 $\pm$ 3 <sup>a</sup>	59.1	0.001
<i>Culex pipiens</i>	95.67 $\pm$ 16.5 <sup>a</sup>	34.33 $\pm$ 1.528 <sup>b</sup>	37.67 $\pm$ 2.08 <sup>b</sup>	38.67 $\pm$ 2.52 <sup>b</sup>	92 $\pm$ 14.73 <sup>a</sup>	29.18	0.001
<i>Culex vagans</i>	10 $\pm$ 1 <sup>a</sup>	2 $\pm$ 1 <sup>b</sup>	6.67 $\pm$ 2.08 <sup>a</sup>	8.33 $\pm$ 1.52 <sup>a</sup>	10.33 $\pm$ 1.52 <sup>a</sup>	15.65	0.001
<i>Anopheles subpictus</i>	96.67 $\pm$ 16.04 <sup>a</sup>	14 $\pm$ 3 <sup>b</sup>	104 $\pm$ 14.11 <sup>a</sup>	0	110 $\pm$ 13.11 <sup>a</sup>	66.93	0.001
<i>Anopheles stephensi</i>	19.33 $\pm$ 2.52 <sup>a</sup>	5 $\pm$ 1 <sup>b</sup>	19.67 $\pm$ 2.08 <sup>a</sup>	0 <sup>c</sup>	20 $\pm$ 2 <sup>a</sup>	87.69	0.001
<i>Anopheles culicifacies</i>	50.33 $\pm$ 1.52 <sup>a</sup>	6.67 $\pm$ 2.08 <sup>b</sup>	46.33 $\pm$ 1.52 <sup>a</sup>	0 <sup>b</sup>	55 $\pm$ 19 <sup>a</sup>	27.73	0.001
<i>Anopheles pulcherrimus</i>	11.33 $\pm$ 1.52 <sup>b</sup>	3 $\pm$ 1 <sup>c</sup>	16 $\pm$ 2 <sup>a</sup>	0 <sup>c</sup>	9.67 $\pm$ 2.08 <sup>b</sup>	53.64	0.001
<i>Anopheles nigerrimus</i>	6 $\pm$ 1 <sup>a</sup> <sup>b</sup>	2.67 $\pm$ 2.08 <sup>b</sup> <sup>c</sup>	6 $\pm$ 1 <sup>a</sup> <sup>b</sup>	0 <sup>c</sup>	8.67 $\pm$ 2.08 <sup>a</sup>	15.94	0.001
<i>Aedes albopictus</i>	64 $\pm$ 12.53 <sup>a</sup>	16.33 $\pm$ 2.08 <sup>b</sup>	24 $\pm$ 2 <sup>b</sup>	23.33 $\pm$ 2.08 <sup>b</sup>	83.33 $\pm$ 14.05 <sup>a</sup>	36	0.001
<i>Aedes aegypti</i>	28.33 $\pm$ 2.08 <sup>a</sup>	6.33 $\pm$ 1.52 <sup>c</sup>	9.67 $\pm$ 2.08 <sup>c</sup>	11 $\pm$ 2 <sup>c</sup>	21 $\pm$ 1 <sup>b</sup>	78.04	0.001

pattern of abundance as that of *Culex quinquefasciatus*. It was highest in the autumn season (Mean =119.33, SD=17.5), and it was lowest in the winter (Mean =0, SD = 0). It was highest during the monsoon season (Mean =119 SD=17.5) and spring (Mean =116, SD=16). This tehsil's *Aedes aegypti* population was found to be missing. Its existence in Tehsil Saddar (F = 31.15, df = 4, P = 0.001) demonstrated a consistent seasonal pattern, with monsoon season occurrence being highest (Mean =73, SD=13) and winter season occurrence being lowest (Mean=9, SD=1) (Table IV) The association between abundance and the season of the other species displayed the similar trend as that of *Culex* and *Anopheles* (Tables III-VIII).




Fig. 3. Significant results of difference of abundance of mosquito in rural and urban areas of Tehsil Hasilpur of District Bahawalpur.




Fig. 4. Non- significant results of difference of abundance of mosquito in rural and urban areas of Tehsil Saddar of District Bahawalpur.

#### Comparative analysis between rural and urban areas

The unpaired t-test was used to compare mosquito populations in different tehsils of the Bahawalpur district between rural and urban areas. According to the findings, the abundance of mosquitoes in the tehsils of Hasilpur, Ahmedpur East, Khairpur Tamewali, and Yazman were significantly high in rural as compared to urban areas (Fig. 3 and Table IX). Tehsil Yazman recorded a highly significant outcome ( $t = 2.64$ ,  $p = 0.028$ ,  $F = 3.530$ ). Results in the



tehsils of Bahawalpur ( $t= 1.617$ ,  $p= 0.144$ ) and Saddar ( $t= 2.059$ ,  $p= 0.073$ ) were not statistically significant (Fig. 4), indicating that these areas are heavily urbanized with little distinction between rural and urban areas.

**Table IX. Comparative analysis between rural and urban areas of different tehsils of district Bahawalpur.**

Tehsils	t	p	F	Difference between Means± Standard error
Hasilpur	2	0.0396	3.117	561.6±228.7
Ahmedpur east	2.38	0.044	3.163	410.4±171.9
Khairpur	2.50	0.0347	3.247	528±207.9
Yazman	2.664	0.0286	3.530	521.2±195.6
Saddar	2.059	0.0734	2.291	523.2±245
Bahawalpur	1.617	0.1446	1.86	326.4±201.9

## DISCUSSION

This is the first time in Pakistan that the spatiotemporal distribution of the mosquitoes in all the tehsils of Bahawalpur is studied along with the comparative analysis of diversity, composition, and abundance of all the genera of mosquitoes between rural and urban areas of different tehsils of Bahawalpur. This study also describes the effect of urbanization and environmental variables on the spatiotemporal distribution of mosquitoes in the Bahawalpur district.

Mashaal (1964) has reported only 2 species of *Anopheles*, *Anopheles stephensi* and *Anopheles culicifacies* as malaria vectors in South Punjab. Different species of *Anopheles* i.e., *Anopheles subpictus*, *Anopheles culicifacies*, *Anopheles stephensi*, *Anopheles pulcherrimus*, *Anopheles Peditaeniatus*, and *Anopheles nigerrimus* have also been reported from South Punjab (Herrel, 2001). All species except *Anopheles Peditaeniatus* were found in this current study. Similarly, *Anopheles subpictus*, *Anopheles stephensi*, *Anopheles culicifacies*, and *Anopheles pulcherrimus* have also been reported, previously. *Culex quinquefasciatus*, *Culex tritaeniorhynchus*, and *Aedes* mosquitoes were also found in wastewater in South Punjab (Mukhtar et al., 2003).

In the current study, the species of genus *Culex* (5 species) and genus *Anopheles* (5 species) were reported in all the rural and urban areas of Bahawalpur. Mosquito abundance and species richness were higher in rural areas than in urban areas. These results agreed with those from previous studies conducted in Europe (Hay et al., 1997;

Johnston et al., 2014) where anthropogenic habitats usually show the lowest abundance of mosquitoes. Rural areas, with freshwater, are more favorable breeding environments, despite those urban areas that may provide suitable habitats for some particular mosquito species (Cox et al., 2007).

*Aedes* mosquitoes were found absent from Tehsil Hasilpur, Khairpur Tamewali, Yazman, and Ahmedpur East. These are periurban areas that are in transition from rural to urban. zone. The species of *Aedes* were sparsely distributed in tehsils Bahawalpur and Saddar. The main reason for the occurrence of *Aedes* in these tehsils was heavy urbanization which was not found in other tehsils. With time, urbanization has increased in these areas due to the conversion of agricultural, forest, and fallow land into urban land (Hussain et al., 2020). With exceptional population growth and rapid urban development, the number of hospitals and clinics has increased several folds which has resulted in huge hospital waste and poor drainage providing favorable conditions for mosquito breeding, especially *Culex* spp. (Khan et al., 2020). The high number of animals in the Bahawalpur Zoo also benefits a variety of mosquito species (Derraik, 2004), especially the *Aedes* mosquitoes which require artificial containers for their breeding (Tuten et al., 2012). In highly urban areas, used tires are not disposed of properly. They effectively hold rainwater and rapidly warm in sunlight even during winter thus providing an ideal breeding habitat for *Aedes* (Khan et al., 2017). Similar results were found in a study conducted in North India where it was observed that tires provide favorable breeding sites for *Aedes aegypti* (Sikhon and Minhas, 2014).

Our study showed that the mosquito species occurred in all six tehsils of Punjab throughout the year in all seasons. The highest abundance was found in the season of monsoon that was during July and August. This has been proved in Pakistan (Resin and Milby, 1986; Mukhtar et al., 2003; Ashfaq et al., 2014; Akram et al., 2009; Fatima et al., 2016) and other countries (Roiz et al., 2014; Valentine et al., 2020). We found the positive effect of precipitation on the relative abundance of mosquito species. The precipitation increases the number of habitats of larvae (Evans et al., 2019). It also has a profound effect on the relative humidity which in turn increases the richness, life span, and host-seeking behavior of mosquitoes (Asigau and Parker, 2018). On the other hand, excessive rain may flush larvae from their habitats and decrease adult mosquito populations (Dieng, 2012).

The current study showed that a high number of species were also found during spring and autumn. During early spring and winter, few species were recorded. The number of species increased again in late April with an

especially high peak in May. In addition, the population density was higher in autumn, especially in September and October. The same results were observed in Pakistan, previously (Akram *et al.*, 2009) and around the globe (Alten *et al.*, 2000; Santos *et al.*, 2020).

It is concluded through this study that changes in the landscape due to urbanization strongly affect the abundance, community composition, and diversity of mosquitoes. Mosquito populations are also affected by changes in climate. This information in the future would be helpful for the public health to design surveillance programs to control the diseases caused by mosquitoes in the reported areas.

### ACKNOWLEDGEMENT

The author acknowledge the help of the Muhammad Asif (Entomologist), MEP Department, Institute of Public Health Lahore for the collection of mosquito.

#### Funding

The author received no financial support for research, authorship, and/or publication of this article

#### Statement of conflict of interest

The authors have declared no conflict of interest.

### REFERENCES

- Akram, W., Hafeez, F., Ullah, U.N., Kim, Y.K., Hussain, A. and Lee, J.J., 2009. Seasonal distribution and species composition of daytime biting mosquitoes. *Entomol. Res.*, **39**: 107-113. <https://doi.org/10.1111/j.1748-5967.2009.00204.x>
- Alten, B., Bellini, R., Caglar, S.S., Simsek, F.M. and Kaynas, S., 2000. Species composition and seasonal dynamics of mosquitoes in the Belek region of Turkey. *J. Vector Ecol.*, **25**: 146-154
- Ashfaq, M., Hebert, P.D., Mirza, J.H., Khan, A.M., Zafar, Y. and Mirza, M.S., 2014. Analyzing mosquito (Diptera: Culicidae) diversity in Pakistan by DNA barcoding. *PLoS One*, **9**: e97268. <https://doi.org/10.1371/journal.pone.0097268>
- Asigau, S. and Parker, P.G., 2018. The influence of ecological factors on mosquito abundance and occurrence in Galápagos. *J. Vector Ecol.*, **43**: 125–137. <https://doi.org/10.1111/jvec.12292>
- Attaullah, M., Gula, S., Bibia, D., Andaleeba, A., Ilahia, A., Sirajb, M., Ahmada, I., Ullaha, M. Alia, S. and Ullaha, Z., 2023. Diversity, distribution, and relative abundance of the mosquito fauna (Diptera: Culicidae) of Malakand and Dir Lower. *Braz. J. Biol.*, **83**: e247374. <https://doi.org/10.1590/1519-6984.247374>
- Barraud, P.J., 1934. *The fauna of British India, including Ceylon and Burma. Diptera: Family Culicidae; Tribes Megarhinini and Culicini*. Vol. 5, Taylor and Francis, London.
- Bashar, K. and Tuno, N., 2014. Seasonal abundance of *Anopheles* mosquitoes and their association with meteorological factors and malaria incidence in Bangladesh. *Parasit. Vectors*, **7**: 1-10. <https://doi.org/10.1186/1756-3305-7-442>
- Bayoh, M.N. and Lindsay, S.W., 2004. Temperature-related duration of aquatic stages of the Afrotropical malaria vector mosquito *Anopheles gambiae* in the laboratory. *Med. Vet. Ent.*, **18**: 174–179. <https://doi.org/10.1111/j.0269-283X.2004.00495.x>
- Byun, R. and Webb, C.E., 2012. *Guidelines for mosquito risk assessment and management in constructed wetlands. A risk assessment tool and management guidelines for councils in the Sydney West region of NSW*. Western Sydney
- Ceccato, P., Connor, S.J., Jeanne, I. and Thomson, M.C., 2005. Application of geographical information systems and remote sensing technologies for assessing and monitoring malaria risk. *Parassitologia*, **47**: 81-96.
- Christophers S.R., 1933. *The fauna of British India, including Ceylon and Burma. Diptera: Family Culicidae; Tribe Anophelini*. Vol. 4, Taylor and Francis, London.
- Cox, J., Grillet, M.E., Ramos, O.M., Amador, M. and Barrera, R., 2007. Habitat segregation of dengue vectors along an urban environmental gradient. *Am. J. trop. Med. Hyg.*, **76**: 820–826. <https://doi.org/10.4269/ajtmh.2007.76.820>
- de Almeida Costa, E.A.P., de Mendonça Santos, E.M., Correia, J.C. and de Albuquerque, C.M.R., 2010. Impact of small variations in temperature and humidity on the reproductive activity and survival of *Aedes aegypti* (Diptera, Culicidae). *Rev. Bras. Ent.*, **54**: 488-493. <https://doi.org/10.1590/S0085-56262010000300021>
- Derraik, G.B., 2004. Mosquitoes (Diptera: Culicidae) breeding in artificial habitats at the Wellington Zoo. *Weta*, **28**: 38-41.
- Dieng, H., Rahman, G.S., Hassan, A.A., Salmah, M.C., Satho, T., Miake, F. Boots, M. and Sazlay, A., 2012. The effects of simulated rainfall on immature population dynamics of *Aedes albopictus* and female oviposition. *Int. J. Biometeorol.*, **56**: 113–120. <https://doi.org/10.1007/s00484-011-0402-0>
- Evans, M.V., Hintz, C.W., Jones, L., Shiau, J., Solano,



- N., Drake, J.M. and Murdock, C.C., 2019. Microclimate and larval habitat density predict adult *Aedes albopictus* abundance in urban areas. *Am. J. trop. Med. Hyg.*, **101**: 362–370. <https://doi.org/10.4269/ajtmh.19-0220>
- Fatima, T., Rais, A., Khan, E., Hills, S.L., Chambers, T.V., Hotwani, A., Qureshi, S., Shafquat, S., Malik, S., Qamar, F., Mir, F., Marfin, A.A., Zaidi, A., Khowaja, A.R. and Shakoor, S., 2020. Investigation of Japanese encephalitis virus as a cause of acute encephalitis in Southern Pakistan, *PLoS One*, **15**: 1-9. <https://doi.org/10.1371/journal.pone.0234584>
- Ferraguti, M., Martínez-de la Puente, J., Roiz, D., Ruiz, S., Soriguer, R. and Figuerola, J., 2016. Effects of landscape anthropization on mosquito community composition and abundance. *Sci. Rep.*, **6**: 29002. <https://doi.org/10.1038/srep29002>
- Hay, S.I., Packer, M.J. and Rogers, D.J., 1997. The impact of remote sensing on the study and control of invertebrate intermediate hosts and vectors for disease. *Int. J. Remote Sens.*, **18**: 2899–2930. <https://doi.org/10.1080/014311697217125>
- Herrel, N., Amerasinghe, F.P., Ensink, J., Mukhtar, M., Van Der Hoek, W. and Konradsen, F., 2001. Breeding of *Anopheles* mosquitoes in irrigated areas of South Punjab, Pakistan. *Med. Vet. Ent.*, **15**: 236-248. <https://doi.org/10.1046/j.0269-283x.2001.00312.x>
- Higgs, S. and Vanlandingham, D., 2015. Chikungunya virus and its mosquito vectors. *Vector Borne Zoonot. Dis.*, **15**: 231-240. <https://doi.org/10.1089/vbz.2014.1745>
- Hussain, M., Khan, A.S., Faheem, M.Z. and Abuhala, M.M., 2020. Urban expansion and land use change in Bahawalpur city during 1998-2018. *Pak. Geogr. Rev.*, **75**: 54-70.
- Imran, M., Ye, J., Saleemi, M.K., Shaheen, I., Zohaib, A., Chen, Z. and Cao, S., 2022. Epidemiological trends of mosquito-borne viral diseases in Pakistan. *Anim. Dis.*, **2**: 1-10. <https://doi.org/10.1186/s44149-021-00034-4>
- Johnston, E., Weinstein, P., Slaney, D., Flies, A.S., Fricker, S. and Williams, C., 2014. Mosquito communities with trap height and urban-rural gradient in Adelaide, South Australia: Implications for disease vector surveillance. *J. Vector Ecol.*, **39**: 48-55. <https://doi.org/10.1111/j.1948-7134.2014.12069.x>
- Khan, A.A., Kanwal, N. and Khan, K., 2020. Medical solid waste generation and associated health problems in Bahawalpur city of southern Punjab, Pakistan. *Pak. Geogr. Rev.*, **75**: 115-137.
- Khan, I.H., Anwar, S. and Hashim, S., 2017. Youth knowledge, attitude and practices about malaria in district Layyah Punjab. *Rev. Econ. Dev. Stud.*, **3**: 125-134. <https://doi.org/10.26710/reads.v3i2.171>
- Kim, T.K., 2015. T test as a parametric statistic. *Korean J. Anesthesiol.*, **68**: 540-546. <https://doi.org/10.4097/kjae.2015.68.6.540>
- Kim, T.K., 2017. Understanding one-way ANOVA using conceptual figures. *Korean J. Anesthesiol.*, **70**: 22-26. <https://doi.org/10.4097/kjae.2017.70.1.22>
- Knowlton, K., Solomon, M.D., Miriam, M. and Rotkin, E., 2009. *Mosquito-borne dengue fever threat spreading in the Americas*. Natural Resources Defense Council, New York. pp. 1-3
- Lawler, S.P. and Lanzaro, G.C., 2005. *Managing mosquitoes on the farm*. UC ANR publication 8158 Department of Entomology, University of California, Davis, CA. <https://doi.org/10.3733/ucanr.8158>
- Mashaal, H.A.H., 1964. Pre and post transmission epidemiological investigations into malaria prevalence in Bahawalnagar district (West Pakistan). *Pak. J. Hlth.*, **14**: 50-82.
- Mohsin, M. and Chinyama, A., 2016. Impacts of solid waste management practices on environment and public health: A case of Bahawalpur city. *Pak. J. agric. Sci.*, **9**: 69-79.
- Mukhtar, M., Herrel, N., Amerasinghe, F.P., Ensink, J., Hoek, W. and Konradsen, F., 2003. Role of wastewater irrigation in mosquito breeding in south Punjab, Pakistan. *Southeast Asian J. trop. Med. Publ. Hlth.*, **34**: 72-80.
- Reisen, W.K. and Milby, M.M., 1986. Population dynamics of some Pakistan mosquitoes: Changes in adult relative abundance over time and space. *Annl Trop. Med. Parasitol.*, **80**: 53-68. <https://doi.org/10.1080/00034983.1986.11811984>
- Reiter, P., 2001. Climate change and mosquito-borne disease. *Environ. Hlth. Perspect.*, **109**: 141-161. <https://doi.org/10.1289/ehp.01109s1141>
- Roiz, D., Ruiz, S., Soriguer, R. and Figuerola, J., 2014. Climatic effects on mosquito abundance in Mediterranean wetlands. *Parasit Vector*, **7**: 1-13. <https://doi.org/10.1186/1756-3305-7-333>
- Rueda, L.M., 2004. Pictorial keys for the identification of mosquitoes (Diptera: Culicidae) associated with dengue virus transmission. *Zootaxa*, **14**: 1-60. <https://doi.org/10.11646/zootaxa.589.1.1>
- Santos, E.M., Favretto, M.A. and Müller, G.A., 2020. When and what time? On the seasonal and daily patterns of mosquitoes (Diptera: Culicidae) in an Atlantic Forest remnant from Southern Brazil.

- Austral. Entomol.*, **59**: 337-344. <https://doi.org/10.1111/aen.12454>
- Sikhon, H. and Minhas, S., 2014. A study of larval indices of *Aedes* and the risk for Dengue outbreak. *Sch. Acad. J. Biosci.*, **2**: 544-547.
- Strong, W.L., 2016. Biased richness and evenness relationships within Shannon–Wiener index values. *Ecol. Indicat.*, **67**: 703-713. <https://doi.org/10.1016/j.ecolind.2016.03.043>
- Styer, L.M., Lim, P., Louie, K.L., Albright, R.G., Kramer, L.D., Kristen A. and Bernard, K.A., 2011. Mosquito saliva causes enhancement of west Nile virus infection in mice. *J. Virol.*, **85**: 1-10. <https://doi.org/10.1128/JVI.01112-10>
- Tun-Lin, W., Burkot, T.R. and Kay, B.H., 2000. Effects of temperature and larval diet on development rates and survival of the dengue vector *Aedes aegypti* in north Queensland, Australia. *Med. Vet. Ent.*, **14**: 31–37. <https://doi.org/10.1046/j.1365-2915.2000.00207.x>
- Tuten, H.C., Bridge, S.W.C., Paul, K.S. and Adler, P.H., 2012. Blood-feeding ecology of mosquitoes in zoos. *Med. Vet. Ent.*, **26**: 407–416. <https://doi.org/10.1111/j.1365-2915.2012.01012.x>
- Valentine, M.J., Ciraola, B., Jacobs, G.R., Arnot, C., Kelly, P.J. and Murdock, C.C., 2020. Effects of seasonality and land use on the diversity, relative abundance, and distribution of mosquitoes on St. Kitts, West Indies. *Parasit. Vectors*, **13**: 1-14. <https://doi.org/10.1186/s13071-020-04421-7>
- Wongkoon, S., Jaroensutasinee, M. and Jaroensutasinee, K., 2013. Distribution, seasonal variation and dengue transmission prediction in Sisaket, Thailand. *Indian J. med. Res.*, **138**: 347-353.

Online First Article