



In-Vitro and *In-Vivo* Antibacterial Effects of Hydroxamic Acid in Broilers

Muhammad Adnan Shams¹, Sarzamin Khan¹, Muhammad Shuaib^{2*},
Muhammad Mushtaq¹, Najma Shams³, Muhammad Aamir⁴, Zahid Ullah⁵,
Shahrood Ahmed Siddiqui⁶ and Bilal Nadeem⁷

¹Department of Poultry Science, Faculty of Animal Husbandry and Veterinary Sciences, The University of Agriculture, Peshawar, Pakistan

²Arid Zone Small Ruminants Research Institute, Ghulam Banda, Kohat, Pakistan

³Higher Education Department Khyber Pakhtunkhwa, Peshawar

⁴Gomal University, Dera Ismail Khan, Pakistan

⁵Livestock Fisheries and Cooperative Department, Khyber Pakhtunkhwa, Pakistan

⁶Vaccine Production Unit Sindh Tandojam, Livestock and Fisheries Department Government of Sindh

⁷College of Veterinary Sciences, Faculty of Animal Husbandry and Veterinary Sciences, The University of Agriculture, Peshawar, Pakistan

ABSTRACT

This study investigated the antibacterial efficacy of hydroxamic acid extracted from Azam maize root powder. The methanolic extract was tested for the availability of hydroxamic acid as an antibacterial agent through *In-Vitro* and *In-Vivo* tests in locally isolated *E. coli* strain-infected broiler chicks. *In-Vitro* experiment, 280-day-old broiler (Cobb) chicks were distributed into 7 experimental groups each with 4 replicates (10 birds/replicate) e.g. CON (control), PC (positive control), Enro (enrofloxacin drug of choice against *E. coli* 5mg/kg body weight), SC (sweet corn dried roots not having any hydroxamic acid, 1000mg/kg feed), FD (freeze dried Azam maize roots, 1000mg/kg feed); SD (sun-dried Azam maize roots, 1000mg/kg feed) and OD (oven dried Azam maize roots, 1000mg/kg feed). Methanolic extract of Azam maize roots showed a significant reduction/inhibition in the growth of *E. coli* strain in which group FD in the range of 11.66 to 0.005mg and groups SD and OD in the range of 11.66 to 5.83 mg were effective. *In-vivo* tests, feed intake, body weight gain, and FCR were significantly improved in the FD group than in the PC, SC, SD, and OD groups during the post-infection phase. Similarly, the mortality index (%) showed a significant reduction in the FD group as compared to the PC, SC, SD, and OD treatment groups. Blood biochemistry parameters e.g., HB, PCV, RBCs, and WBCs were significantly affected in the FD group. Similarly, liver and kidney function tests e.g., AST, ALT, creatinine, and blood urea showed more potential antioxidant effects in the FD group as compared to the PC, SC, SD, and OD treatment groups. The histopathological examination of liver, kidney, and intestine tissues in infected birds showed significantly different architectural structures. The FD group showed normal sectional architectural study as compared to other treatment groups. Based on the above facts, it is concluded that among the different methods of methanolic extract of Azam maize roots, the FD group has a good antimicrobial effect as well as improved performance and serum biochemistry against locally isolated *E. coli* strain-infected broiler chicks.

Article Information

Received 09 October 2023

Revised 10 December 2023

Accepted 26 December 2023

Available online 01 May 2024

(early access)

Authors' Contribution

MAS, Animal trial, laboratory test, and manuscript writing. SK, Study design, feed formulation, and statistical analysis. MS, MS, Statistical analysis, data evaluation, manuscript writing, and review. NS, MA, ZU SAS, BN, Manuscript writing and review.

Key words

Medicinal plants, Broiler chicks, Hydroxamic acid, *E. coli*, Toxicity, Hematology

INTRODUCTION

Commercial poultry farming is gradually progressing and contributes to the provision of protein sources as well as employment either directly or indirectly (Hussain *et al.*, 2015) but still, this sector is facing many challenging diseases like colibacillosis which generates hurdles in the progress of this sector (Apostolakis *et al.*, 2021). Colibacillosis results in significant economic losses every year around the globe due to its high morbidity and mortality rates (Baranwal *et al.*, 2019). Systemic infection

* Corresponding author: shoaibwzr@gmail.com
0030-9923/2024/0001-0001 \$ 9.00/0



Copyright 2024 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

caused by *Escherichia coli* is the main reason for mortality (Guabiraba *et al.*, 2015). *E. coli* is a commensal facultative anaerobic microorganism that can grow in the lower intestinal tract of broiler birds shortly after birth and acts as a symbiont, helping in the synthesis of necessary vitamins for the host (Apostolakos *et al.*, 2021). Avian colibacillosis causes perihepatitis, airsacculitis, and pericarditis, although other syndromes such as coligranuloma, omphalitis, cellulitis, osteomyelitis/arthritis, and especially diarrhea may be noticed (Panth, 2019). Colibacillosis can occur as a secondary bacterial infection alongside other bacterial infections (Guabiraba and Schouler, 2015).

Pakistan produces about 7.9 million tonnes of maize annually and its production has increased 6% during 2019-2020. It contributes 0.6 % to the GDP (Khan *et al.*, 2021). Maize is being grown on an area of 1.016 million hectares with a typical grain yield of 2,864 kg/ha. Because of its protein composition, interest in the domesticated animals and poultry feed industry has increased. Azam maize is a medicinal plant so its chemical composition is the main concern for the research (Amanullah *et al.*, 2015). Hydroxamic acid is the obvious proof of its medicinal composition. Hydroxamic acid is prepared from a chemical compound an ester (via Lossen rearrangement reaction) or chlorides (by a reaction with hydroxylamine salts) as well as from aldehydes and sulfonyl hydroxylamine (via the Angeli-Rimini reaction). Hydroxamic acid and its chemically modified derivatives are efficient iron binders, bacterial protein synthesis inhibitors, and growth factor blockers and are good legends as well as metal chelators (Citarella *et al.*, 2021). They can remove heavy metals from the body easily through their excellent metal-binding ability (Bhawani *et al.*, 2014). Hydroxamic acid possesses *in-vitro* antibacterial properties (Pal and Saha, 2012). Many researchers have worked on the beneficial chemical effects of hydroxamic acids against microbial activities, and anti-cancer and anti-fungal properties as well (Citarella *et al.*, 2021). The medicinal properties of hydroxamic acid are solving many complicated drug resistance issues through its unique binding and chelating actions as well as anti-bacterial functions against different bacterial species and fungi (Griffith *et al.*, 2021). Syed *et al.* (2020) reported about the antibacterial activity of heterocyclic hydroxamic acids against some species of bacteria and fungi. Shekhawat and Rajput (2012) reviewed the same activity of hydroxamic acids against microbes and stated that they are efficient chelators, iron and lead binding agents. Therefore, this study was designed to determine the *in-vitro* and *in-vivo* antibacterial efficacy of hydroxamic acid extracted from the Azam maize plant in induced *E. coli* infection in broiler chicks.

MATERIAL AND METHODS

Study area

The experiments were performed for the investigation of *in-vitro* and *in-vivo* antibacterial effects of hydroxamic acids in the Azam variety of maize roots against the locally isolated *E. coli* strain-challenged broiler birds. The experiments were conducted at the poultry farm unit, Department of Poultry Science while the lab work was executed at the College of Veterinary Science, the University of Agriculture, Peshawar, Pakistan in the Pharmacology and Histopathology Labs.

Sample collection and maize seeds cultivation procedure

For the extraction of hydroxamic acid, local varieties of maize seeds i.e., Azam maize and sweet corn seeds were cultivated in a Petri plate using tissue paper as an anchor material. Ten seeds per petri plate were cultivated. For proper activation of the seeds, the Petri plate was kept completely dark with no light access. After a few days, the dormancy of the seeds was finished and became activated. On day 7 of the cultivation, the roots of these Azam maize plants and sweet corn plants were separated and collected. After collection, these roots were dried in equal amounts following three different drying procedures, namely freeze drying, oven drying, and sun drying procedure. After drying maize roots were crushed into powder through a grinding mill. Then using a sonicator machine methanolic extract was prepared from both the sweet corn and Azam maize roots powder for an *in vitro* test against *E. coli*. Two-fold dilutions were performed for 11.66 mg concentrations of methanolic extracts *in vitro* test against *E. coli*. There were 7 experimental groups e.g., control (CON), positive control (PC), enrofloxacin group (ENRO), sweet corn (SC) dried roots powder group, Azam maize freeze-dried (FD) roots powder group, Azam maize sun-dried (SD) roots powder group, Azam maize oven dried (OD) roots powder group with 4 replicates in each group.

Methodology for the collection and preparation of *E. coli* culture

Proper confirmation on API strip, locally isolated *E. coli* strain was collected from infected broiler visceral organs as per already established samples collection protocol (Mansour *et al.*, 2015). After the collection of *E. coli* bacteria, proper culturing of *E. coli* bacteria (10^7 CFU) in nutrient broth was made and the culture was kept overnight at 37 °C.

In-vitro antibacterial bioassay of the hydroxamic acids in broiler

The minimum inhibitory concentration test was

performed to explore the *in-vitro* antibacterial impacts of hydroxamic acid extracted from Azam maize roots and sweet corn roots against locally isolated strains of *E. coli*. The *in-vitro* antibacterial bioassay was performed using the broth dilution method as described by Neal *et al.* (2012) with some modifications: Nutrient broth 100 μ l was added to each/96 wells of microtitration plate. These wells were properly labeled and sterilized as well. In the 1st row, only broth and in the 2nd row only *E. coli* was added. In the 3rd row, the drug of choice, Enrofloxacin, and *E. coli* bacteria were added. In the 4th row, 11.66 μ l sweet corn root extract and locally isolated *E. coli* strain were added, and a 2-fold dilution up to well 12 was prepared. In the 5th row, 11.66 μ l Azam maize freeze-dried roots extract and *E. coli* bacteria were added and 2-fold dilution was prepared up to well number 12. In the 6th row, 11.66 μ l Azam maize oven-dried roots extract and *E. coli* bacteria were added and 2-fold dilution was prepared up to well number 12. In the 7th row, 11.66 μ l Azam maize sun-dried roots extract and *E. coli* bacteria were added and a 2-fold dilution was prepared up to well number 12. Then incubated in an oven for 1 to 2 days at 37°C. The bacterial growth was assessed by measuring optical density (OD) 600 nm with the help of a spectrophotometer at 24 h post-inoculation.

In-vivo antibacterial bioassay of the hydroxamic acids in broiler against E. coli

For *in-vivo* bioassays, from a commercial hatchery, 280 broiler chicks (Cobb) were purchased and distributed randomly into 7 experimental groups e.g., CON, PC, ENRO, SC dried roots powder group, Azam maize FD roots powder group, Azam maize SD roots powder group, Azam maize OD roots powder group with 4 replicates in each group having 10 broilers. The required data was collected on 2nd week of the trial and the total duration of the experimental trials was 35 days. After an adaptation period of 7 days, all the experimental groups were infected with locally isolated *E. coli* strain except the control group. After 2 to 3 days post-infection, the ENRO group was treated with the drug of choice enrofloxacin @ 5mg/kg body weight, and the SC group with sweet corn dried roots powder @1000mg/kg. FD, SD, and OD experimental groups were fed with Azam maize freeze, sun, and oven-dried roots powder @ 1000mg/kg feed accordingly for 7 days.

Production performance parameters

Feed intake (FI), body weight gain (BWG), feed conversion ratio (FCR), and mortality were recorded weekly as described by Shuaib *et al.* (2020). FI and BWG were calculated. The FCR was calculated by using the formula; total FI \div total BWG.

Hematology and serum biochemical analysis

At 2nd week of the trial, broiler chicks were slaughtered and both the blood and tissue samples were collected in sterile test tubes from 3 birds per replicate in each group. Whole blood samples were used for the determination of the hematological profile of the diseased broiler and to correlate the result with the good health status of the healthy birds. Total red blood cells (RBCs) and total white blood cells (WBCs) count, hemoglobin (HB) concentration, and packed cell volume (PCV) from the blood were determined using standard procedures (Sanderson and Philips 1981). Using 4000 rpm centrifugation for 10 min, serum was separated from the blood sample, which was used to foster biochemical analysis, and liver and kidney function tests according to the protocols adapted by IFCC. Commercially available kits (Sigma) were preferred for these tests according to the manufacturer's instructions.

Histopathology

After 1 week of the trial, three birds per replicate were slaughtered within each group. Postmortem was performed and gross lesions were observed and recorded in different locations especially visceral organs. The samples from the liver, kidney, and digestive tract/intestine were taken and fixed in 10% buffered formalin for histopathological examination as per standard protocol of Bancroft *et al.* (1994). Hematoxylin and eosin (H and E) were the basic stains preferred in this process. The slides were observed for any alterations in the morphology and also microscopic lesions on tissue. The digital camera was used for saving pictures.

Data analysis

Complete randomized design (CRD) was used for the statistical analysis of data. The least significant test at 5% probability was carried out to distinguish the mean differences. For analysis of the data, the statistical package statistics 8.1 was used.

RESULTS

The results regarding the minimum inhibitory concentration (MIC) for the growth of locally isolated *E. coli* strain against different dilutions are presented in Table I. Parent concentration of methanolic extract at 11.66 to 0.005 mg of Azam maize roots showed significant reduction/inhibition in growth of locally isolated *E. coli* strain. It is indicated from the MIC test outcomes that the root powder of Azam maize freeze-dried having hydroxamic acid was effective in the range of 11.66 to 0.005mg. The first 2 wells of Azam maize sun-dried root extract and Azam maize oven-dried root extract showed

Table I. Minimum inhibitory concentration of hydroxamic acid against *Escherichia coli*.

Group	Concentration (mg)											
	11.66	5.83	2.91	1.45	0.72	0.36	0.18	0.091	0.045	0.022	0.011	0.005
CON	0.042 ^b	0.048 ^c	0.039 ^c	0.042 ^c	0.032 ^c	0.048 ^d	0.030 ^c	0.038 ^c	0.020 ^c	0.054 ^c	0.037 ^c	0.041 ^c
PC	0.640 ^a	0.454 ^b	0.425 ^a	0.494 ^a	0.699 ^a	0.611 ^a	0.613 ^a	0.521 ^a	0.728 ^a	0.860 ^a	0.462 ^b	0.607 ^a
Enro	0.060 ^b	0.062 ^c	0.065 ^c	0.041 ^c	0.052 ^c	0.052 ^d	0.051 ^c	0.063 ^c	0.060 ^c	0.070 ^c	0.033 ^c	0.044 ^c
SC	0.640 ^a	0.631 ^a	0.425 ^a	0.494 ^a	0.699 ^a	0.611 ^a	0.613 ^a	0.521 ^a	0.728 ^a	0.864 ^a	0.462 ^b	0.607 ^a
FD	0.040 ^b	0.062 ^c	0.065 ^c	0.041 ^c	0.052 ^c	0.052 ^d	0.051 ^c	0.063 ^c	0.060 ^c	0.070 ^c	0.038 ^c	0.040 ^c
SD	0.080 ^b	0.096 ^c	0.282 ^b	0.217 ^b	0.300 ^b	0.230 ^c	0.266 ^b	0.425 ^b	0.635 ^b	0.607 ^b	0.521 ^a	0.635 ^a
OD	0.040 ^b	0.06 ^c	0.259 ^b	0.520 ^a	0.500 ^b	0.520 ^b	0.317 ^b	0.466 ^b	0.574 ^b	0.429 ^b	0.483 ^b	0.368 ^b
SEM	0.44	0.55	0.98	0.84	0.73	0.77	0.97	0.95	0.91	0.66	0.89	0.97
P. value	0.027	0.018	0.012	0.022	0.034	0.027	0.023	0.020	0.031	0.014	0.019	0.033

Means in the similar column with various superscripts are varies at ($p < 0.05$). CON, Control; PC, Positive control (bacteria effected); Enro, Enrofloxacin shows the drug of choice against *E. coli* bacteria (5mg/kg body weight); SC, Sweet corn dried roots not having any hydroxamic acid (1000mg/kg feed); FD, Freeze dried Azam maize roots having hydroxamic acid (1000mg/kg feed); SD, Sun dried Azam maize roots (1000mg/kg feed); OD, Oven dried Azam maize roots (1000mg/kg feed).

Table II. Antibacterial effects of hydroxamic acid on overall performance of broiler chicks during *E.coli* infection.

Groups	Performance parameters			
	FI (g)	BWG (g)	FCR	Mortality (%)
CON	550.1 ^a	399 ^a	1.37 ^c	0.00 ^c
PC	436.5 ^d	243 ^d	1.79 ^a	54.29 ^a
Enro	528.1 ^b	379 ^b	1.39 ^c	0.00 ^c
SC	430.12 ^d	244 ^d	1.76 ^a	53.30 ^a
FD	518.23 ^b	373 ^b	1.38 ^c	0.00 ^c
SD	475.15 ^c	303 ^c	1.50 ^b	20.00 ^b
OD	470.39 ^c	308 ^c	1.52 ^b	16.60 ^b
SEM	9.25	18.33	0.43	4.35
P value	0.001	0.001	0.004	0.001

Means in the similar column with various superscripts are varies at ($p < 0.05$). BWG, Body weight gain; FI, Feed intake; FCR, Feed conversion ratio. For groups description see [Table I](#).

antibacterial effects. [Table II](#) shows the results regarding the antibacterial activity of hydroxamic acid on the production performance parameters like FI, BWG, FCR, and mortality in broiler chicks. A significant improvement in FI, BWG, and FCR in the post-infection phase was recorded in the CON group as compared to all other groups. Among all other groups, ENRO and FD groups showed better results. Similarly, a significant reduction in mortality was recorded in the CON, ENRO, and FD groups than in the remaining groups. The results on hematological parameters like RBCs and WBCs, Hb, and PCV are presented in [Table III](#). The Hb was calculated significantly higher in the CON and FD

groups while PCV in the PC and SC groups but RBCs in the CON and WBCs in the FD diet groups as compared to all other groups. [Table IV](#) shows the results regarding the blood biochemical parameters e.g., AST, ALT creatinine, and blood urea level in broiler chicks. A significantly higher AST, ALT creatinine, and blood urea were calculated in the PC and SC diet groups as compared to all other groups. The results regarding the histopathological examination of the internal visceral organs like the liver, intestine, and kidney are described in [Figures 1, 2, and 3](#). The liver sample of the FD group demonstrated no histological reflection in contrast with a control group

Table III. Antibacterial effects of hydroxamic acid on overall blood profile of broiler chicks during *E.coli* infection.

Groups	Blood profile			
	Hb (g/dL)	PCV (%)	RBC ($10^{12}/\mu\text{L}$)	WBC ($10^9/\mu\text{L}$)
CON	9.62 ^a	30.22 ^c	3.56 ^a	3.40 ^b
PC	6.57 ^d	43.21 ^a	2.56 ^d	2.89 ^d
Enro	9.33 ^b	27.67 ^c	2.94 ^b	3.39 ^b
SC	6.60 ^d	42.22 ^a	2.57 ^d	2.92 ^d
FD	9.70 ^a	28.01 ^c	3.01 ^b	3.49 ^a
SD	8.70 ^c	36.60 ^b	2.77 ^c	3.17 ^c
OD	8.73 ^c	38.62 ^b	2.79 ^c	3.20 ^c
SEM	1.60	1.10	1.67	1.46
P value	0.812	0.030	0.010	0.080

Means in the similar column with various superscripts are varies at ($p < 0.05$). Hb, Hemoglobin; PCV, Pack cell volume; RBC, Red blood cells; WBC, White blood cells. For groups description see [Table I](#).

Table IV. Antibacterial effects of hydroxamic acid on blood biochemical parameters of broiler chicks during *E. coli* infection.

Groups	Blood biochemical parameters			
	AST (IU/ml)	ALT (IU/ml)	Creatinine (mg/dL)	Blood urea (mg/dL)
CON	18.1 ^c	27.5 ^c	0.55 ^c	13.9 ^c
PC	69.5 ^a	67.0 ^a	1.49 ^a	37.9 ^a
Enro	26.4 ^c	25.3 ^c	0.45 ^c	11.7 ^c
SC	64.5 ^a	66.0 ^a	1.35 ^a	32.9 ^a
FD	23.4 ^c	25.3 ^c	0.44 ^c	14.2 ^c
SD	54.0 ^b	50.9 ^b	1.01 ^b	23.6 ^b
OD	57.6 ^b	58.8 ^b	1.07 ^b	26.9 ^b
SEM	2.33	1.17	1.18	1.30
P value	0.851	0.151	0.225	0.001

Means in the similar column with various superscripts are varies at ($p < 0.05$). AST, Aspartate aminotransferase; ALT, alanine aminotransferase. For groups description see Table 1.




Fig. 1. A: NC (necrotic cell) hepatocyte pyknotic nuclei of *E. coli* affected broiler liver. B: NC (necrotic cell and hemorrhages) broiler liver treated with sweet corn roots powder. C: MNC (mild necrosis of cell) broiler liver treated with sun dried roots powder. D: MNC (mild necrosis of cell) broiler liver treated with oven dried roots powder. E: NCM (normal cell morphology) broiler liver treated with freeze dried roots powder. F: NCM (normal cell morphology) broiler liver treated with enrofloxacin. G: NCM (normal cell morphology) broiler liver of control group.




Fig. 2. A: NVM (necrosed villus morphology and effected crypt) *E. coli* affected broiler intestine. B: NVM (necrosed villus morphology and effected crypt) broiler intestine treated with sweet corn roots powder. C: MNVM (mild necrosed villus morphology and effected crypt) broiler intestine treated with sun dried roots powder. D: MNVM (mild necrosed villus morphology and effected crypt) broiler intestine treated with oven dried roots powder. E: NVLW (normal villus length and width) broiler intestine treated with freeze dried roots powder: intestine L 0.190mm, 0.006mm width. F: NVLW (normal villus length and width) broiler intestine treated with enrofloxacin: intestine L 0.200mm, 0.007mm width. G: NVLW (normal villus length and width) broiler intestine of control group: intestine L 0.194mm, 0.008mm width.

and the hepatocytes showed a typical appearance in every cell as compared to other treatment groups as presented in Figure 1. Similarly, an intestinal sample of the FD group demonstrated no histological reflection in contrast with a control group and the villi showed typical histology having pointed zenith and more extensive basal parts as compared to other treatment groups as shown in Figure 2. Similarly, the kidney sample of broiler chicks in the FD group demonstrated no histological reflection in contrast with the control group and all the cells showed normal glomerulus and distal convoluted tubules as compared to other treatment-provided groups as presented in Figure 3.




Fig. 3. A: HGDC (hemorrhagic glomerulus and distal convoluted tubules) *E. coli* effected broiler kidney. B: HGDC (hemorrhagic glomerulus and distal convoluted tubules) broiler kidney treated with sweet corn roots powder. C: MHGDC (mild hemorrhagic glomerulus and distal convoluted tubules) broiler kidney treated with sun dried roots powder. D: MHGDC (mild hemorrhagic glomerulus and distal convoluted tubules) broiler kidney treated with oven dried roots powder. E: NM (normal morphology) broiler kidney treated with freeze dried roots powder: F: NM (normal morphology) broiler kidney treated with enrofloxacin intestine. G: NM (normal morphology) broiler kidney of control group.

DISCUSSION

In vitro, antibacterial bioassay (MIC) of hydroxamic acid against *E. coli* shows that freeze-dried root powder of Azam maize was successful against the tried micro-organism (*E. coli*). The base inhibitory concentration of this concentrate against *E. coli* was 11.66 to 0.005 mg. The findings of the present study are in agreement with the results of [Zaltron *et al.* \(2020\)](#), [Ramisetty *et al.* \(2016\)](#), and [Jahangirian *et al.* \(2011\)](#). Usually, hydroxamic acid and their derivatives are prepared from a chemical compound an ester (via lossen rearrangement reaction), or chlorides by a reaction with hydroxylamine salts as well as from aldehydes and sulfonylhydroxylamine via the Angeli-Rimini reaction. Hydroxamic acid and its derivatives can

remove heavy metals from the body easily through their excellent metal binding ability ([Parveen *et al.*, \(2019\)](#)). The overall performance in broiler chicks at the post-infection phase was found significant in a group of birds that were treated with methanolic extract of freeze-dried Azam roots powder in induced locally isolated *E. coli* strain infection as compared to other treatment groups. Our findings agree with the results of [Ebeid *et al.* \(2021\)](#) who reported that overall performance was improved with the supplementation of organic acids in water or feed. Similarly, our results agree with the findings of [Adil *et al.* \(2010, 2011\)](#) who stated that FCR and weight gain were improved with supplementation of the organic acids in poultry feed. Similarly, our results agree with the findings of [Saaf *et al.* \(2008\)](#) who reported that weight gain increases with the supplementation of organic acids in broilers. The improvement in body weight gain in infected broiler chicks might be due to the anti-*E. coli* activity of hydroxamic acid. The decrease in *E. coli* activity may be due to the unavailability of growth-regulating factors (free iron) in the body. No free iron availability in the body may be due to the hydroxyamic acid conjugation with the free iron available in the body ([Weinberg, 1984](#)). Hydroxamic acid is denatured by heat treatment which is why the oven and sun-dried azam maize roots show low hydroxamic acid activates as compared to freeze-dried (freezing compacts the hydroxamic acid concentration). The decrease in *E. coli* infection may be due to its enzyme deactivation by the hydroxamic acid. The role of hydroxamic acid against *E. coli* may be due to its peptide deformylase enzyme-blocking activity ([Indu *et al.*, \(2010\)](#)). *In-vitro* exploration was the obvious proof that hydroxamic acids and their chemically modified derivatives are efficient iron binders, bacterial protein synthesis inhibitors, and growth factor blockers and are good legends as well as metal chelators.

Liver, kidney, and intestinal samples of broiler chicks offered with freeze-dried Azam maize roots showed no histological deviations in contrast with the control group. Our findings agree with the findings of [Rodríguez *et al.* \(2012\)](#) who stated that organic acid (1.0% sorbic acid and 0.2% citrus extract) supplementation fundamentally raised the villus width, height, and region of the duodenum, jejunum, and ileum of oven chicks at 14 days old enough. The findings of the current study agree with [García *et al.* \(2007\)](#) who announced that broilers took care of diets containing formic acid had the longest villi (1273 and 1250 μm for 0.5 and 1.0% formic acid, individually) contrasted with control (1088 μm). Our result agrees with the findings of [Saaf *et al.* \(2008\)](#) who concluded that the liver capabilities didn't antagonistically change in light of the increase in organic acidifiers. Similarly, our finding is in line with [Rizwan *et al.* \(2014\)](#) who reported that an

increase of organic acids in the drinking water did not show any effect on the strength of and kidney.

All these investigations and examinations revealed that freeze-dried Azam maize is a powerful medicinal plant with antibacterial characteristics both *in vitro* and *in vivo* research world. The Azam maize plant is extremely protected to utilize as it did not have any toxicopathological impacts in the broiler. Sun-dried roots and oven-dried roots of the Azam maize plant have a very small quantity of hydroxamic acid, which can be concluded that the decreased amount of hydroxamic acid in sun-dried and oven-dried roots of the Azam maize plant may be due to damage done by heat in both the groups during the drying procedure. Sweet corn dried roots did not show any antibacterial activity as it was double muted and did not possess any hydroxamic acid.

CONCLUSION

It is concluded that Azam maize roots powder dried through freeze method possess *In-vitro* antibacterial (*E. coli*) property in the range of 0.005 to 11.66 mg and has significantly improved the production performance, hematology and serum biochemistry parameters and intestinal histopathology and can be used at the range of 0.005 to 11.66 in the *E. coli* infected broiler chicks.

ACKNOWLEDGMENT

The authors would like to thank the staff of the Department of Poultry Science, The University of Agriculture Peshawar, for the laboratory and technical assistance.

Funding

Not applicable.

IRB approval

The experimental work was approved by the Advanced Studies and Research Board (No.219/PS/UAP) dated June 2022, The University of Agriculture Peshawar, KP, Pakistan.

Ethical statement

The study was approved by the ethical committee of the Faculty of Animal Husbandry and Veterinary Sciences, The University of Agriculture, Peshawar, before the practical execution of this experiment.

Statement of conflict of interest

The authors have declared no conflict of interest.

REFERENCES

- Adil, S., Bandy, M. T., Bhat, G. A., Khan, I. and Bhat, M.I., 2010. A study on the profitability of broiler chicken fed diets supplemented with organic acids. *Indian J. Poult. Sci.*, **45**: 220-223. <https://doi.org/10.4061/2010/479485>
- Adil, S., Bandy, M., Bhat, G., Qureshi, S. and Wani, S., 2011. Effect of supplemental organic acids on growth performance and gut microbial population of broiler chicken. *Livest. Res. Rural Dev.*, **3**: 203-214.
- Amanullah, K.I., Jan, A., Jan, M.T., Khalil, S.K., Shah, Z. and Afzal, M., 2015. Compost and nitrogen management influence productivity of spring maize (*Zea mays* L.) under deep and conventional tillage systems in semi-arid regions. *Commun. Soil Sci. Pl. Anal.*, **46**: 1566-1578. <https://doi.org/10.1080/00103624.2015.1043462>
- Apostolakos, I., Laconi, A., Mughini-Gras, L., Yapicier, Ö.Ş. and Piccirillo, A., 2021. Occurrence of colibacillosis in broilers and its relationship with avian pathogenic *Escherichia coli* (APEC) population structure and molecular characteristics. *Front. Vet. Sci.*, **8**: 737720. <https://doi.org/10.3389/fvets.2021.737720>
- Bancroft, J.D., Cook, H.C. and Stirling, R.W., 1994. Manual of histological techniques and their diagnostic application. In: *Manual of histological techniques and their diagnostic application*. pp. 457-457.
- Baranwal, A., Singh, N.P., Kumar, A., Kumar, A., Upadhyaya, P.K. and Srivastava, A.K., 2019. Colibacillosis in broilers: A case report. *J. Ent. Zool. Stud.*, **7**: 163-166.
- Citarella, A., Moi, D., Pinzi, L., Bonanni, D. and Rastelli, G., 2021. Hydroxamic acid derivatives: From synthetic strategies to medicinal chemistry applications. *ACS Omega*, **6**: 21843-21849. <https://doi.org/10.1021/acsomega.1c03628>
- Ebeid, T., Al-Homidan, I., Fathi, M., Al-Jamaan, R., Mostafa, M., Abou-Emera, O. and Alkhalaf, A., 2021. Impact of probiotics and/or organic acids supplementation on growth performance, microbiota, antioxidative status, and immune response of broilers. *Ital. J. Anim. Sci.*, **20**: 2263-2273. <https://doi.org/10.1080/1828051X.2021.2012092>
- Garcia, V., Catala-Gregori, P., Hernandez, F., Megias, M.D. and Madrid, J., 2007. Effect of formic acid and plant extracts on growth, nutrient digestibility, intestine mucosa morphology, and meat yield of

- broilers. *J. appl. Poult. Res.*, **16**: 555-562. <https://doi.org/10.3382/japr.2006-00116>
- Griffith, D.M., Li, H., Werrett, M.V., Andrews, P.C. and Sun, H., 2021. Medicinal chemistry and biomedical applications of bismuth-based compounds and nanoparticles. *Chem. Soc. Rev.*, **50**: 12037-12069. <https://doi.org/10.1039/D0CS00031K>
- Guabiraba, R. and Schouler, C., 2015. Avian colibacillosis: Still many black holes. *FEMS Microbiol. Lett.*, **362**: fnv118. <https://doi.org/10.1093/femsle/fnv118>
- Hussain, J., Rabbani, I., Aslam, S. and Ahmad, H.A., 2015. An overview of poultry industry in Pakistan. *World's Poult. Sci. J.*, **71**: 689-700. <https://doi.org/10.1017/S0043933915002366>
- Indu, S., Hemlata, K., Sunil, K., Arun, K., Suman, L. and Ashok, K., 2010. Synthesis of new coumarin derivatives as antibacterial agents. *Int. J. Chem. Tech. Res.*, **2**: 1745-1752.
- Jahangirian, H., Haron, J., Silong, S., Yusof, N., Shameli, K., Eissazadeh, S. and Jafarzade, M., 2011. Antibacterial effect of phenyl fatty hydroxamic acids synthesized from canola oil. *J. Med. Pl. Res.*, **5**: 4826-4831.
- Khan, A.G., Imran, M., Khan, A.U.H., Fares, A., Šimůnek, J., Ul-Haq, T. and Ali, S., 2021. Performance of spring and summer-sown maize under different irrigation strategies in Pakistan. *Sustainability*, **13**: 2757. <https://doi.org/10.3390/su13052757>
- Mansour, S.M., Ali, H., Chase, C.C. and Cepica, A., 2015. Loop-mediated isothermal amplification for diagnosis of 18 World Organization for Animal Health (OIE) notifiable viral diseases of ruminants, swine and poultry. *Anim. Hlth. Res. Rev.*, **16**: 89-106. <https://doi.org/10.1017/S1466252315000018>
- Neal, A.L., Ahmad, S., Gordon-Weeks, R. and Ton, J., 2012. Benzoxazinoids in root exudates of maize attract *Pseudomonas putida* to the rhizosphere. *PLoS One*, **7**: 35498. <https://doi.org/10.1371/journal.pone.0035498>
- Pal, D. and Saha, S., 2012. Hydroxamic acid- A novel molecule for anticancer therapy. *J. Adv. Pharm. Technol. Res.*, **3**: 92-99. <https://doi.org/10.4103/2231-4040.97281>
- Panth, Y., 2019. Colibacillosis in poultry: A review. *J. Agric. Natl. Resour.*, **2**: 301-311. <https://doi.org/10.3126/janr.v2i1.26094>
- Parveen, P., Brundavani, K.D., Mahathi, K., Bhavani, M.S. and Shaheda Sultana, S.K., 2019. Gene silencing and DNA methylation. *Am. J. Phytomed. clin. Therapeut.*, **2**: 1-12.
- Ramisetty, B.C.M., Raj, S. and Ghosh, D., 2016. *Escherichia coli* MazEF toxin-antitoxin system does not mediate programmed cell death. *J. Basic Microbiol.*, **56**: 1398-1402. <https://doi.org/10.1002/jobm.201600247>
- Rizwan, S., Naqshbandi, A., Farooqui, Z., Khan, A.A. and Khan, F., 2014. Protective effect of dietary flaxseed oil on arsenic-induced nephrotoxicity and oxidative damage in rat kidney. *Fd. Chem. Toxicol.*, **68**: 99-107. <https://doi.org/10.1016/j.fct.2014.03.011>
- Rodríguez-Lecompte, J.C., Yitbarek, A., Brady, J., Sharif, S., Cavanagh, M.D., Crow, G. and Camelo-Jaimes, G., 2012. The effect of microbial-nutrient interaction on the immune system of young chicks after early probiotic and organic acid administration. *J. Anim. Sci.*, **90**: 2246-2254. <https://doi.org/10.2527/jas.2011-4184>
- Saaf, E.S.M., El-Mednay, N. and Abdel-Azeem, F., 2008. Thyroid activity, some blood constituents, organs morphology and performance of broiler chicks fed supplemental organic acids. *Int. J. Poult. Sci.*, **7**: 215-222. <https://doi.org/10.3923/ijps.2008.215.222>
- Sanderson, J.H. and Phillips, C.E., 1981. *An atlas of laboratory animal haematology*. Clarendon Press.
- Shekhawat, S. and Rajput, S.K., 2012. Mechanistic investigation for the acidic hydrolysis of p-methyl benzyl-2-theno hydroxamic acid. *Chem. Sci. J.*, ink.gale.com/apps/doc/A335292529/AON-E?u=anon~3d1c3cb5&sid=googleScholar&id=ace41367.
- Shuaib, M., Ullah, N., Hafeez, A., Alhidary, I.A., Abdelrahman, M.M. and Khan, R.U., 2020. Effect of dietary supplementation of wild Cumin (*Bunium persicum*) seeds on performance, nutrient digestibility and circulating metabolites in broiler chicks during the finisher phase. *Anim. Biotechnol.*, **33**: 871-875. <https://doi.org/10.1080/10495398.20.1844222>
- Syed, Z., Sonu, K., Dongre, A., Sharma, G. and Sogani, M., 2020. A review on hydroxamic acids: Widespectrum chemotherapeutic agents. *Int. J. Biol. Biomed.*, **14**: 75-88. <https://doi.org/10.46300/91011.2020.14.12>
- Weinberg, E.D., 1984. Iron withholding: A defense against infection and neoplasia. *Physiol. Rev.*, **64**: 65-102. <https://doi.org/10.1152/physrev.1984.64.1.65>
- Zaltron, A., Merano, M., Mistura, G., Sada, C. and Seno, F., 2020. Optical tweezers in single-molecule experiments. *Eur. Phys. J. Plus*, **135**: 896. <https://doi.org/10.1140/epjp/s13360-020-00907-6>