



Assessing Wildlife and Ecology Amid Infrastructure Development in the Upper Indus Basin, Pakistan

Usman Ahmad¹, Sajid Rashid Ahmad^{1*}, Zulfiqar Ali^{2*} and Rida Ahmad^{2,3}

¹College of Earth and Environmental Sciences, University of the Punjab, Lahore

²Institute of Zoology, University of the Punjab, Lahore

³Department of Animal Sciences, University of the Punjab, Lahore, Pakistan

ABSTRACT

Understanding the baseline ecology of the Upper Indus Basin, Pakistan has become absolutely essential with ongoing infrastructure projects in the region as it would have various consequences on different habitats of the area. This study aims to provide a comprehensive understanding of the ecological dynamics, offering insights into the richness and diversity of fauna species of the area. The study area encompasses Upper Indus Basin, the Karakoram and Himalayan Mountain ranges, and was concentrated on the infrastructure developments in Khyber Pakhtunkhwa and Gilgit-Baltistan along the Indus River. Field surveys were conducted from May 2021 to December 2022 using the point count method for birds, pitfall method for reptiles and tracks and stools were utilized to identify mammals along with the direct observations, focusing on the nine identified habitats. Thorough search method was employed for all the taxa (birds, mammals, reptiles and amphibians). The surveyed species included birds, mammals, reptiles, and amphibians, and data were collected using GPS, binoculars, digital cameras, spotting scopes, and field guides. The study recorded a total of 322 species, including 242 birds, 47 mammals, 21 reptiles, and 12 amphibians. The values of Margalef, Menhinick's, Simpson, Shannon-Weiner and Evenness are as follows: 31.98, 2.13, 0.94, 3.65 and 0.63. Critical information on taxonomy, IUCN status, and trends for each group was analyzed along with hotspots. Overall, this study reveals that the area has high species diversity and provides habitat associations between nine habitats of the area. The infrastructure developments can disrupt these habitats, leading to biodiversity loss and altered species distribution. The findings contribute to a better understanding of the ecological dynamics in this region and can help in devising conservation and management strategies for the diverse habitats present.

Article Information

Received 23 February 2024

Revised 25 July 2024

Accepted 05 August 2024

Available online 26 September 2024 (early access)

Authors' Contribution

UA, SRA and ZA conceptualized the study. UA, ZA and RA conducted the field visits. UA, ZA and RA assembled and analyzed the dataset. UA and RA drafted the manuscript. SRA and ZA reviewed and improved the manuscript.

Key words

Infrastructure development, Habitat, Ecological baseline, Upper Indus Basin, Biodiversity hotspots, Conservation, Wildlife conservation strategies, Wildlife management strategies

INTRODUCTION

Situated in the mountainous expanse of the Hindu-Kush Karakoram Himalaya (HKH) region, the Upper Indus Basin (UIB) is characterized by a transnational river network that includes both the western rivers (Upper Indus, Kabul, Jhelum, and Chenab) and the eastern rivers (Ravi and Satluj) (Orr *et al.*, 2022). The water resources in the UIB exhibit pronounced seasonality, mainly dependent on the influx of water from snow and glacial melt during the spring and summer months, as well as the summer

monsoonal rainfall (Lutz *et al.*, 2014, 2016). In Pakistan, the majority of agricultural land relies on irrigation, and irrigation-based agriculture contributes approximately 90% of the total annual agriculture production (Hussain *et al.*, 2021), which accounts for about 25% of the country's GDP (Zahra *et al.*, 2023). The river holds immense importance as Afghanistan, Pakistan, India and China rely on river systems that originate from the mountainous region of the Upper Indus Basin (UIB) (Mukherji *et al.*, 2019; Wester *et al.*, 2019). The water resources within the UIB are vital for various purposes, including agriculture, power generation, domestic consumption, industrial processes, tourism, fishing, and religious ceremonies. Furthermore, they support a diverse range of terrestrial and aquatic ecosystems (Xu *et al.*, 2019). These rivers sustain the world's largest continuous irrigation system (Qureshi, 2011) and play a critical role in ensuring a reliable supply of electricity to downstream populations through numerous hydropower projects (Nie *et al.*, 2021; Hennig *et al.*, 2023).

Development of large transportation facilities, economic zones, and infrastructure are among the main measures for the development of the remote and

* Corresponding author: zali.zool@pu.edu.pk, sajidpu@yahoo.com
0030-9923/2024/0001-0001 \$ 9.00/0



Copyright 2024 by the authors. Licensee Zoological Society of Pakistan.

This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

disadvantaged areas of any country (Ramachandran and Linde, 2011). A well-built road network helps in accessibility to the basic facilities of life, such as health, education, employment, and economic opportunity (Sackey *et al.*, 2023). UIB is confronted with the challenge of massive and rapid infrastructure development in its entire history (Khan *et al.*, 2020). Mega infrastructure projects like China Pak Economic Corridor, Dams, High Voltage Transmission Lines, etc. are transforming natural landscape. It is expected to result in inducing very negative impacts on the environment, ecology and natural resources of the area (Huo *et al.*, 2021). Extensive anthropogenic stressed land use practices are already triggering climate change impacts (Xiaolong *et al.*, 2021). UIB is of great scientific importance, as it serves as a crucial source of freshwater for agricultural purposes and the generation of hydropower electricity (Yaseen *et al.*, 2020). Additionally, it plays a pivotal role in preserving biodiversity, owing to its high-altitude mountains, extensive glaciers, and unique ecological support systems (Zhang and Wang, 2021). The identification of ecologically vulnerable regions constitutes a fundamental aspect within the realm of ecological conservation and environmental management. This paper attempts to provide biodiversity hotspots and ecological baseline for UIB considering the major infrastructure developments in the area.

MATERIALS AND METHODS

Study area

The present study was conducted in the Upper Indus

Basin (72.20°–77.69° E; 32.92°–37.09° N), define as the Tarbela dam catchment area in Pakistan. The area lies at higher elevation ranging from 1000 to more than 5000 meters. It comprises of about 147,000 km², focused on the Karakorum and Himalaya Mountain ranges. Study includes primarily the areas of Haripur, Abbottabad, Mansehra, Battagram, Shangla and Kohistan districts of Khyber Pakhtunkhwa; and Diamer, Astore, Skardu and Ghanche districts of Gilgit-Baltistan, along river Indus.

The area was stratified into nine potential habitats: agriculture land, alpine grassland, bare rock/gravel, snow/glacier, sub-tropical broad leaf, sub-tropical chir pine, temperate forest, urban area/ settlements and waterbodies/swamp (Table I, Fig. 1).




Fig. 1. Map highlighting nine potential habitats in the Upper Indus Basin area studied for ecological assessment.

Table I. Brief description of habitats.

| S. | Habitat | Description |
|----|----------------------------|---|
| 1 | Agriculture land | Cultivated land used for growing crops, fertile soil, hot and dry climate in the plains, cooler climate in the hilly areas, mix of natural vegetation and cultivated crops. |
| 2 | Alpine grassland | Cold temperatures, low growing vegetation, grasses and herbs adapted to harsh environmental conditions. |
| 3 | Bare rock/ gravel | Rocky and gravelly terrain with little or no vegetation, thin and nutrient-poor soils, extreme temperatures |
| 4 | Snow/Glacier | Snow-capped peaks and glaciers, harsh environmental conditions, limited vegetation, wildlife adapted to cold conditions |
| 5 | Sub-tropical broad leaf | Dense forests with broadleaf trees adapted to warm, humid areas, diverse plant and animal life, mangrove forests in the Indus Delta region |
| 6 | Sub-tropical chir pine | Forests dominated by Chir Pine trees, wildlife includes the Himalayan black bear, leopard and several bird species |
| 7 | Temperate forest | Trees that shed leaves during winter months, diverse vegetation and wildlife, moderate temperatures |
| 8 | Urban area/ Settlements | Concentrated areas of buildings and infrastructure, high population density, limited natural vegetation |
| 9 | Waterbodies/ Swamp | Wetland areas with diverse plant and animal life, important breeding grounds for fish and bird species, provide important ecological services |

Field surveys

The monthly surveys were conducted between May 2021 and December 2022. Fifteen points were selected in each habitat for primary observations. In addition to spending 15 min at each point for direct observations of birds, other methods were employed for different habitats and species. In grassland and agricultural areas, the flushing method was used, while the sky view method was utilized for observing raptors. Points were also strategically selected to set up pitfalls for reptiles. For mammals, direct observations were complemented by identifying tracks and stool. Thorough search method was employed for all the taxa (birds, mammals, reptiles and amphibians). The equipment used for this study included GPS, binoculars (Bushnell power view, 60X 90 m), digital camera (Nikon p-900), spotting scope (Harrier 65mm ED Spotting Scope) and field guides Birds of Pakistan (Roberts, 1991, 1992; Mirza and Wasiq, 2007; Grimmitt *et al.*, 2008), Mammals of Pakistan (Roberts, 1997) and Amphibians and Reptiles of Pakistan (Khan, 2006). These guides were used for species identification.

Data analysis

The data were analyzed by using formulae of Simpsons index, Shannon-Wiener diversity index species evenness, Margalef's index, Menhinicks index, Relative abundance, Sorenson's similarity index, and Jaccard similarity index, Simpson's index $(D) = 1 - \sum (n_i) / N(N-1)$

Where, n is total number of individuals of a particular species, and N is total number of individuals of all species (Ahmad *et al.*, 2022).

Shannon-Wiener diversity index $(H') = -\sum_{i=1}^S P_i \ln P_i$

Where, H' Shannon-wiener diversity index, S is No. of Species, P_i is Proportion of a species relative to the total number of individuals (Ahmad *et al.*, 2022).

Species evenness: $(J') = H' / \ln(S)$

Where, J' is Pielou Index, H' is Shannon-Wiener Index, and S is No. of species (Kazam *et al.*, 2022)

Margalef's diversity index $(D_{Mg}) = S - 1 / \ln N$

Where, S is No. of species, and N is total no. of individuals in the sample (Gamito, 2010).

Menhinick's diversity index $(D_{Mn}) = S / \sqrt{N}$

Where, S is No. of species, and N is total no. of individuals in the sample (Mulya *et al.*, 2021).

Relative abundance $(RA) = n/N \times 100$

Where, n is number of individuals of the specie i in the specific area, N is total number of individuals of all species in a specific area (Kazam *et al.*, 2022).

Sorenson's similarity index $(SSI) = (2C/(a+b)) * 100$

Where, C is number of species in both habitats a and b , a is number of species in habitat a , b is number of species in habitat b . The similarity index was used to determine the

interspecific linkages between species in different habitats (Sørensen, 1948). SSI between two was calculated using the following formula (Nath *et al.*, 2005).

Jaccard similarity index $(C_j) = a / (a + b + c)$

Where, a is the number of common species in both habitats, b is the number of species in habitat a , not in habitat b while c is the number of species in habitat b , not present in habitat a . It determines and compares the similarities between various habitats.

It is important to note that neither coefficient (Sorenson's and Jaccard) takes into account the number of individuals present in each area (Barbour *et al.*, 1980; Magurran, 2004).

Identification of hotspots

Hotspots were identified based on the density and the highest number of infrastructure projects in an area. This method highlights regions with the most significant concentration of development activity.

RESULTS

The major infrastructure developments in the area include Dam/Hydropower, dry Ports/SEZ, roads, settlements and transmission lines as highlighted in Figure 2. A total of 322 species were recorded in different habitats of Upper Indus Basin. It included 242 birds, 47 mammals, 21 reptiles and 12 amphibians. Further details about order, family, IUCN status and trend are given in Supplementary Table I and a summary is given in Table II.

Table II. Taxonomic details and IUCN status.

| | Birds | Mam- mals | Rep- tiles | Amphib- ians |
|----------------------------|-------|--------------|---------------|-----------------|
| Order | 22 | 9 | 2 | 1 |
| Family | 63 | 23 | 9 | 2 |
| IUCN status | | | | |
| Least concern (LC) | 219 | 29 | 4 | 10 |
| Near threatened (NT) | 10 | 6 | | |
| Vulnerable (VU) | 8 | 6 | | |
| Endangered (EN) | 3 | 3 | 1 | |
| Critically endangered (CR) | 2 | | | |
| Data deficient (DD) | | | 1 | |
| Not available (NA) | | 3 | 15 | 2 |
| Trend | | | | |
| Increasing | 40 | 3 | | 1 |
| Decreasing | 77 | 18 | 2 | 1 |
| Stable | 100 | 9 | 1 | 7 |
| Unknown | 25 | 14 | 1 | 1 |
| Not available | | 3 | 17 | 2 |




Fig. 2. Hotspots identified by the density and number of infrastructure projects, indicating regions with high hydropower activity in Upper Indus Basin. (1) Tarbela Dam, (2) Allai Khwar, (3) Khan Khwar, (4) Thakot Dam, (5) Palas valley upper, (6) Palas valley lower, (7) Palas valley middle, (8) Pattan Dam site, (9) Keyal Khwar, (10) Dubair Khwar, (11). Spat Gah Lower, (12) Spat Gah Middle, (13) Spat Gah Upper, (14) Dasu Dam, (15) Karang, (16) Kaigah, (17) Tangir, (18) Basha Dam, (19) Phander, (20) Baru, (21) Kanch, (22) Sher Qila, (23) Naltar- 1, (24) Nasirabad, (25) Altit, (26) Bunji Dam Site, (27) Doyian, (28) Talu, (29) Yulbo Dam Site, (30) Tormic, (31) Harpo, (32) Tungas Dam Site, (33). Kachura, (34) Skardu, (35) Basha, (36) Satpara, (37) Shyok (Yugo) HPP.

Among birds, order Passeriformes was most abundant with 119 species followed by Charadriiformes, Accipitriformes with 25 and 17 species, respectively. The most abundant bird species were great cormorant (*Phalacrocorax carbo*), black-headed gull (*Larus ridibundus*), carrion crow (*Corvus corone*), large-billed crow (*Corvus macrorhynchos*) and house crow (*Corvus splendens*) with relative abundance 14.56, 10.48, 9.56, 7.46 and 5.00, respectively. Only two critically endangered species were recorded during surveys, sociable lapwing (*Vanellus gregarius*) and white-rumped vulture (*Gyps bengalensis*). Among mammals, order Carnivora exhibited the highest species richness with 15 species followed by Rodentia (14 species). Three endangered species were recorded *viz.*, Indian pangolin (*Manis crassicaudata*), Indian wild dog (*Cuon alpinus*) and Himalayan musk deer (*Moschus moschiferus*). In reptiles twenty species belonged to order Squamata and only one species belonged to order Testudines. All species belonged to order Anura in amphibians.

Three hotspots were identified based on the infrastructure developments in the study area. Hotspot A stretches from upstream of Battagram city of District Battagram up to District Kohistan in the North, all along the River Indus. The area inhabits 201 species (169 birds, 18

mammals, six reptiles and eight amphibians). Hotspot B spreads around Kachura town just downstream of Skardu, all along the River Indus. The area inhabits 79 species (62 birds, 11 mammals, five reptiles and one amphibian). Hotspot C lies upstream of Gilgit covering areas of Jutial and Naltar Valleys, along the Gilgit River. The area inhabits species (90 birds, 12 mammals and two reptiles). Forty-six species are common in these hotspots and can be considered as species of concern as their population may decline due to physical and environmental changes during construction and operation phases of the infrastructure developments (Supplementary Table II).

Table III. Species richness, abundance and diversity indices.

| | Rich-ness | Abun-dance | Indices | | | | |
|------------|-----------|------------|----------|-----------|----------|----------|-----------|
| | | | Margalef | Menhinick | Simp-son | Shan-non | Even-ness |
| Overall | 322 | 22850 | 31.98 | 2.13 | 0.94 | 3.65 | 0.63 |
| Birds | 242 | 22487 | 24.05 | 1.61 | 0.96 | 3.57 | 0.65 |
| Mammals | 47 | 148 | 9.21 | 3.86 | 0.96 | 3.49 | 0.9 |
| Reptiles | 21 | 194 | 3.80 | 1.51 | 0.89 | 2.43 | 0.79 |
| Amphibians | 12 | 21 | 3.61 | 2.62 | 0.94 | 2.37 | 0.95 |

Different indices were employed to understand the diversity and habitat associations of the area. The value of Margalef's index for birds was 31.98 referring to high richness while Menhinick's index value was 2.13. The minimum value for this index is one (01) and as the value increases, it corresponds to higher diversity. The indices details are given in Table III. The richness indices, such as Margalef and Menhinick's, offer insights into the diversity of each taxonomic group, indicating higher diversity among birds and mammals. The diversity indices, including the Simpson and Shannon-Weiner indices, suggest that overall species composition is relatively diverse and evenly distributed, with birds exhibiting slightly higher diversity compared to other groups. The Shannon-Weiner index value of 3.65 indicates a high level of diversity in the area.

Sørensen's and Jaccard's similarity coefficients are valuable tools for comparing species associations across different areas/habitats. Given that both Sørensen and Jaccard exhibited similar trends in the raw data (Fig. 2), the discussion will focus solely on Sørensen's coefficient. The habitat associations revealed that based on species composition, agriculture land has a close relationship with bare rock/gravel followed by urban area/settlements (Table IV). Moreover, Alpine grassland was closely related to snow/glaciers and Waterbodies/swamp. The urban area/settlements habitat, having a relatively lower similarity coefficient with other habitats, appears to be

distinct from natural ecosystems. Overall, these similarity coefficients offer insights into the ecological relationships

and associations among different habitats.

Table IV. Habitat associations based on sorenson index.

| | Alpine grassland | Bare rock/ Gravel | Snow/ Glacier | Sub-tropical broad leaf | Sub-tropical chir pine | Temperate forest | Urban area/ Settlements | Waterbodies/ swamp |
|-------------------------|---------------------|----------------------|------------------|----------------------------|---------------------------|---------------------|----------------------------|-----------------------|
| Agriculture land | 0.898 | 0.983 | 0.883 | 0.877 | 0.920 | 0.974 | 0.979 | 0.955 |
| Alpine grassland | | 0.895 | 0.984 | 0.817 | 0.875 | 0.915 | 0.884 | 0.929 |
| Bare rock / Gravel | | | 0.884 | 0.860 | 0.906 | 0.973 | 0.968 | 0.944 |
| Snow/Glacier | | | | 0.803 | 0.858 | 0.900 | 0.869 | 0.917 |
| Sub-tropical broad leaf | | | | | 0.902 | 0.883 | 0.891 | 0.851 |
| Sub-tropical chir pine | | | | | | 0.926 | 0.931 | 0.915 |
| Temperate forest | | | | | | | 0.969 | 0.941 |
| Urban area/ Settlements | | | | | | | | 0.939 |

DISCUSSION

Recent studies have improved our understanding of the global impacts of dams and hydropower, aided by emerging databases and maps. However, data limitations, particularly in rapidly developing regions like South Asia's Third Pole, hinder the quality of these analyses. Assessing biodiversity and ecology amid infrastructure development in the Upper Indus Basin (part of South Asia's Third Pole), Pakistan, holds paramount significance due to the region's unique ecological characteristics and the potential environmental impacts associated with large-scale development projects. A comprehensive assessment of biodiversity and ecology was essential to understand the baseline conditions and identify the key habitats. In the Upper Indus Basin. There are a total of around 567 species that can be found, comprising 400 birds, 103 mammals, 48 reptiles, and 16 amphibians according to the guides, Birds of Pakistan (Roberts, 1991, 1992; Mirza and Wasiq, 2007; Grimmett *et al.*, 2008), Mammals of Pakistan (Roberts, 1997) and Amphibians and Reptiles of Pakistan (Khan, 2006). In the course of the surveys conducted, a total of 322 species were observed, encompassing 242 birds, 47 mammals, 21 reptiles, and 12 amphibians. The most abundant bird species included great cormorant (14.56), black-headed gull (10.48), carrion crow (9.56), large-billed crow (7.46) and house crow (5.00) and Ahmad *et al.* (2022) also reported carrion crow (7.48) and larger-billed crow (6.24) as most abundant species of the area between Raikot and Thakot. While Ammanat *et al.* (2022) reported great cormorant (9.36) and carrion crow (5.73) as most abundant species in the area of Dasu Transmission Line (Raikot to Tarbela including the area of Islamabad West Grid Station). We identified three hotspots with 46 common species (forty birds and six mammals) based

on the infrastructure developments in the area. The birds mainly included waterbirds, raptors, pheasants and wheatears. May *et al.* (2021) also suggested that raptors followed by waterfowls were the most affected species specially in terms of collision and barrier during infrastructure developments in Norway. The mammals included common otter (*Lutra lutra*), Himalayan musk deer (*Moschus cupreus*), Himalayan pika (*Ochotona roylei*), Himalayan wood mouse (*Apodemus rusiges*), Turkestan rat (*Rattus turkestanicus*) and house mouse (*Mus musculus*). Common otter and Himalayan musk deer are endangered species and the infrastructure developments can cause further decline in their population. According to Khan *et al.* (2014), musk deer is at the brink of extinction due to exclusion from trophy hunting programs. The decline in musk deer population is not solely attributed to hunting but also to the destruction of their habitat, exacerbated by the growing human and livestock populations in mountainous regions (Nandy *et al.*, 2011, 2015; Navalgund *et al.*, 2019).

These species can be impacted by infrastructure developments either during construction or implementation phase. Ammanat *et al.* (2022) conducted a study on species vulnerable to collisions with the 765 kV Dasu Transmission Line, identifying 27 such species in the area. Of these, 23 species were also observed in the present study. Hydropower projects specially run off river, cause major variations in hydrological as well as ecological conditions (Ullah *et al.*, 2023). The Himalayan-Karakoram-Hindukush region harbors exceptionally diverse and globally significant biodiversity, but the implementation of large-scale hydropower development projects poses a significant threat to the habitats of numerous terrestrial plant and animal species (Sharma and Thakur, 2017). While hydropower has the potential to help address the challenges of climate change and enhance energy security

(Berga, 2016), it must not come at the cost of biodiversity loss (Carolli *et al.*, 2023). Research conducted in the HKH region has indicated that such projects can lead to deforestation, resulting not only in the loss of biodiversity within forest ecosystems but also adversely affecting communities reliant on forests for their livelihoods (Pandit and Grumbine, 2012; Batool and Abbas, 2017). Pandit and Grumbine (2012) predicted that the construction of dams in the Indian Himalayas could lead to the extinction of seven vertebrate taxa. These projects also have the potential to disrupt the habitats of aquatic species, further exacerbating the environmental consequences (Kumar and Katoch, 2014).

Jameel *et al.* (2022) also reported threats to the avian species because of illegal hunting, deforestation and major infrastructure development projects. Moreover, major infrastructure development projects pose multiple threats, encompassing issues such as sound and air pollution, resettlement and relocation of local communities residing near riverbeds (dam reservoirs), and the subsequent encroachment upon the pheasants habitats with higher conservation potential (Awan *et al.*, 2021; Bagaria *et al.*, 2021). The Western Tragopan and Koklass Pheasant, observed in the current study, may face similar risks due to ongoing infrastructure developments in the area. These projects also inflict severe damage upon both the avian population and their respective habitats (Hussain *et al.*, 2019). Moreover, hydropower plants have the adverse effect of interrupting and fragmenting the vast expanse of forest cover, which serves as a crucial predictor for species richness and functional biodiversity (Alho *et al.*, 2019). The construction activities associated with hydroelectric plants, regardless of their size, exert multifaceted impacts on both biodiversity and habitat (Alho, 2020). Habitat destruction and fragmentation can result in a loss of biodiversity as species struggle to find suitable places to live and reproduce, reducing the genetic variability (Valenzuela-Aguayo *et al.*, 2020). This can lead to declines in population sizes, and in some cases, even extinction of certain species (Palmeirim and Gibson, 2021). The forthcoming development of hydropower projects has the potential to impede the achievement of sustainable development goal 6 (clean water and sanitation) and sustainable development goal 15 (life on land) (Opoku, 2019) as it may engender local extinctions of various species, including fish, macroinvertebrates, as well as terrestrial flora and fauna (Hermoso *et al.*, 2019). Given that human well-being is intricately linked to biodiversity and the ecosystem services it provides, the preservation of biodiversity has emerged as a critical factor in attaining sustainable development objectives (Gillespie *et al.*, 2015).

Hydropower projects implemented within the Upper

Indus Basin of Pakistan have far-reaching implications for biodiversity. The construction of dams and reservoirs can alter and diminishes natural habitats, leading to the displacement or loss of local species. This disruption of habitats can disrupt migratory patterns and gene flow, ultimately affecting population dynamics (Alho, 2020). Furthermore, the regulation of water flow for hydropower purposes modifies the natural flow regimes of rivers, potentially causing significant impacts on aquatic ecosystems. Altered water flow patterns, including reduced downstream flow, can have detrimental effects on the availability of suitable habitats for various species (Bagaria *et al.*, 2021). Additionally, hydropower projects influence water quality through processes such as sedimentation, changes in temperature regimes, and alterations in nutrient levels. These combined factors contribute to the overall ecological impact of hydropower projects on the biodiversity of the region. This comprehensive understanding of the ecological dynamics in the region serves as a valuable resource for formulating conservation and management strategies tailored to the diverse habitats present. The research thus contributes significantly to the knowledge base necessary for effective conservation and sustainable development in this ecologically crucial area.

CONCLUSION

In conclusion, the research in the Upper Indus Basin has revealed rich biodiversity and highlighted the importance of balancing development with conservation efforts. By identifying key species and hotspots, the study offers valuable insights for informed decision-making. Prioritizing conservation measures alongside infrastructure development is crucial for maintaining ecosystem integrity and promoting sustainable growth in the region.

DECLARATIONS

Funding

The study did not receive any funding.

Ethical statement

The Advanced Study & Research Board at University of the Punjab, Lahore approved the study (Letter number: 6813/Acad, Dated: 12/07/2021).

Supplementary material

There is supplementary material associated with this article. Access the material online at: <https://dx.doi.org/10.17582/journal.pjz/20240223121505>

Statement of conflict of interest


The authors have declared no conflict of interest.

REFERENCES

- Ahmad, R., Ali, Z., Manzoor, F., Ahmad, U., Sidra, S., Zainab, I., Furqan, M., Batool, A. and Zona, Z., 2022. Avian diversity, abundance and habitat suitability index for threatened species in selected areas of Northern Pakistan. *Pakistan J. Zool.*, **55**: 2001-2500. <https://doi.org/10.17582/journal.pjz/20220224070218>
- Alho, C.J., 2020. Hydropower dams and reservoirs and their impacts on Brazil's biodiversity and natural habitats: A review. *World J. Adv. Res. Rev.*, **6**: 205-215. <https://doi.org/10.30574/wjarr.2020.6.3.0197>
- Alho, C.J., Mamede, S.B., Benites, M., Andrade, B.S. and Sepúlveda, J.J., 2019. Threats to the biodiversity of the Brazilian Pantanal due to land use and occupation. *Ambient. Soc.*, **22**. <https://doi.org/10.1590/1809-4422asoc201701891vu201913ao>
- Ammanat, M., Qadir, A., Ali, Z., Ahmad, R., Ahmad, U., Zainab, I. and Batool, A., 2022. Avian diversity around Indus River with collision prone species abundance at proposed 765 KV transmission line. *Pakistan J. Zool.*, **55**: 2385-2390. <https://doi.org/10.17582/journal.pjz/20220507120544>
- Awan, M.N., Saqib, Z., Buner, F., Lee, D.C. and Pervez, A., 2021. Using ensemble modeling to predict breeding habitat of the red-listed Western Tragopan (*Tragopan melanocephalus*) in the Western Himalayas of Pakistan. *Glob. Ecol. Conserv.*, **31**: e01864. <https://doi.org/10.1016/j.gecco.2021.e01864>
- Bagaria, P., Thapa, A., Sharma, L.K., Joshi, B.D., Singh, H., Sharma, C.M., Sarma, J., Thakur, M. and Chandra, K., 2021. Distribution modelling and climate change risk assessment strategy for rare Himalayan *Galliformes* species using archetypal data abundant cohorts for adaptation planning. *Clim. Risk Manage.*, **31**: 100264. <https://doi.org/10.1016/j.crm.2020.100264>
- Barbour, M.G., Burk, J.H. and Pitts, W.D., 1980. *Terrestrial plant ecology*. Benjamin/Cummings, Menlo Park. pp. 604.
- Batool, A. and Abbas, F., 2017. Reasons for delay in selected hydro-power projects in Khyber Pakhtunkhwa (KPK), Pakistan. *Renew. Sustain. Energy Rev.*, **73**: 196-204. <https://doi.org/10.1016/j.rser.2017.01.040>
- Berga, L., 2016. The role of hydropower in climate change mitigation and adaptation: A review. *Engineering*, **2**: 313-318. <https://doi.org/10.1016/J.ENG.2016.03.004>
- Carolli, M., de Leaniz, C.G., Jones, J., Belletti, B., Hudek, H., Pusch, M., Pandakov, P., Börger, L. and van de Bund, W., 2023. Impacts of existing and planned hydropower dams on river fragmentation in the Balkan region. *Sci. Total Environ.*, **871**: 161940. <https://doi.org/10.1016/j.scitotenv.2023.161940>
- Gamito, S., 2010. Caution is needed when applying Margalef diversity index. *Ecol. Ind.*, **10**: 550-551. <https://doi.org/10.1016/j.ecolind.2009.07.006>
- Gillespie, B.R., Desmet, S., Kay, P., Tillotson, M.R. and Brown, L.E., 2015. A critical analysis of regulated river ecosystem responses to managed environmental flows from reservoirs. *Freshw. Biol.*, **60**: 410-425. <https://doi.org/10.1111/fwb.12506>
- Grimmett, R., Roberts, T. and Inskipp, J., 2008. *Birds of Pakistan*. Christopher Helm, London Yale University Press, New Heaven. pp. 256.
- Hennig, T., Harlan, T., Tilt, B. and Magee, D., 2023. Hydropower development in South Asia: Data challenges, new approaches, and implications for decision-making. *WIREs Water*, **10**: p.e1654. <https://doi.org/10.1002/wat2.1654>
- Hermoso, V., Clavero, M. and Green, A.J., 2019. Don't let damage to wetlands cancel out the benefits of hydropower. *Nature*, **568**: 171-171. <https://doi.org/10.1038/d41586-019-01140-7>
- Huo, C., Hameed, J., Nawaz, A., Shah, S.A.R., Alqahtani, W., Maqsoom, A. and Anwar, M.K., 2021. Scientific risk performance analysis and development of disaster management framework: A case study of developing Asian countries *J. King Saud Univ. Sci.*, **33**: 101348. <https://doi.org/10.1016/j.jksus.2021.101348>
- Hussain, A., Qamar, F.M., Adhikari, L., Hunzai, A.I., Rehman, A.U. and Bano, K., 2021. Climate change, mountain food systems, and emerging opportunities: A study from the Hindu Kush Karakoram Pamir Landscape, Pakistan. *Sustainability*, **13**: 3057. <https://doi.org/10.3390/su13063057>
- Hussain, A., Sarangi, G.K., Pandit, A., Ishaq, S., Mamnun, N., Ahmad, B. and Jamil, M.K., 2019. Hydropower development in the Hindu Kush Himalayan region: Issues, policies and opportunities. *Renew. Sustain. Energy Rev.*, **107**: 446-461. <https://doi.org/10.1016/j.rser.2019.03.010>
- Jameel, M.A., Nadeem, M.S., Aslam, S., Ullah, W., Ahmad, D., Awan, M.N., Masroor, W., Mahmood, T., Ullah, R., Anjum, M.Z. and Ali, K., 2022. Impact of human imposed pressure on pheasants of western Himalayas, Pakistan: Implication for monitoring and conservation. *Diversity*, **14**: 752. <https://doi.org/10.3390/d14090752>
- Kazam, A., Sidra, S., Ali, Z., Ahmad, R., Bilal, A.

- and Batool, A., 2022. *Field validation of avian diversity at uchalli wetland complex: A Ramsar site in Khushab, Pakistan*. pp. 1-9. <https://doi.org/10.17582/journal.pjz/20220225100237>
- Khan, A.J., Koch, M. and Tahir, A.A., 2020. Impacts of climate change on the water availability, seasonality and extremes in the Upper Indus Basin (UIB). *Sustainability*, **12**: 1283. <https://doi.org/10.3390/su12041283>
- Khan, M.S., 2006. *Amphibians and reptiles of Pakistan*. Krieger Publishing Company, Malabar, Florida.
- Khan, M.Z., Khan, B., Khan, E., Garee, A., Khan, A., Rehmat, A., Abbas, A.S., Ali, M. and Hussain, E., 2014. Abundance distribution and conservation of key ungulate species in Hindu Kush Karakoram and Western Himalayan (HKH) mountain ranges of Pakistan. *Int. J. Agric. Biol.*, **16**: 1560-8530.
- Kumar, D. and Katoch, S.S., 2014. Harnessing water tower into power tower: A small hydropower development study from an Indian prefecture in western Himalayas. *Renew. Sustain. Energy Rev.*, **39**: 87-101. <https://doi.org/10.1016/j.rser.2014.07.052>
- Lutz, A.F., Immerzeel, W.W., Kraaijenbrink, P.D.A., Shrestha, A.B. and Bierkens, M.F.P., 2016. Climate change impacts on the upper Indus hydrology: Sources, shifts and extremes. *PLoS One*, **11**: e0165630. <https://doi.org/10.1371/journal.pone.0165630>
- Lutz, A.F., Immerzeel, W.W., Shrestha, A.B. and Bierkens, M.F.P., 2014. Consistent increase in high Asia's runoff due to increasing glacier melt and precipitation. *Nat. Clim. Change*, **4**: 587-592. <https://doi.org/10.1038/nclimate2237>
- Magurran, A.E., 2004. *Measuring biological diversity*. Blackwell Publishing, Oxford.
- May, R., Jackson, C.R., Middel, H., Stokke, B.G. and Veronesi, F., 2021. Life-cycle impacts of wind energy development on bird diversity in Norway. *Environ. Impact Assess. Rev.*, **90**: 106635. <https://doi.org/10.1016/j.eiar.2021.106635>
- Mirza, Z.B. and Wasiq, H., 2007. *A field guide to birds of Pakistan*. Published by WWF-Pakistan. Book land Lahore.
- Mukherji, A., Sinisalo, A., Nüsser, M., Garrard, R. and Eriksson, M., 2019. Contributions of the cryosphere to mountain communities in the Hindu Kush Himalaya: A review. *Region. Environ. Change*, **19**: 1311-1326. <https://doi.org/10.1007/s10113-019-01484-w>
- Mulya, H., Santosa, Y. and Hilwan, I., 2021. Comparison of four species diversity indices in mangrove community. *Biodiv. J. Biol. Divers.*, **22**: 3648-3655. <https://doi.org/10.13057/biodiv/d220906>
- Nandy, S., Kushwaha, S.P.S. and Dadhwal, V.K., 2011. Forest degradation assessment in the upper catchment of the river Tons using remote sensing and GIS. *Ecol. Indic.*, **11**: 509-513. <https://doi.org/10.1016/j.ecolind.2010.07.006>
- Nandy, S., Singh, C.D.K.K., Das, K.K., Kingma, N.C. and Kushwaha, S.P.S., 2015. Environmental vulnerability assessment of eco-development zone of Great Himalayan National Park, Himachal Pradesh, India. *Ecol. Indic.*, **57**: 182-195. <https://doi.org/10.1016/j.ecolind.2015.04.024>
- Nath, P.C., Arunachalam, A., Khan, M.L., Arunachalam, K. and Barbhuiya, A.R., 2005. Vegetation analysis and tree population structure of tropical wet evergreen forests in and around Namdapha National Park, northeast India. *Biodiv. Conserv.*, **14**: 2109-2135. <https://doi.org/10.1007/s10531-004-4361-1>
- Navalgund, R.R., Kumar, A.S. and Nandy, S., 2019. *Remote sensing of northwest Himalayan ecosystems* (Vol. 1). Singapore: Springer. <https://doi.org/10.1007/978-981-13-2128-3>
- Nie, Y., Pritchard, H.D., Liu, Q., Hennig, T., Wang, W., Wang, X., Liu, S., Nepal, S., Samyn, D., Hewitt, K. and Chen, X., 2021. Glacial change and hydrological implications in the Himalaya and Karakoram. *Nat. Rev. Earth Environ.*, **2**: 91-106. <https://doi.org/10.1038/s43017-020-00124-w>
- Opoku, A., 2019. Biodiversity and the built environment: Implications for the sustainable development goals (SDGs). *Resour. Conserv. Recycl.* **141**: 1-7. <https://doi.org/10.1016/j.resconrec.2018.10.011>
- Orr, A., Ahmad, B., Alam, U., Appadurai, A., Bharucha, Z.P., Biemans, H., Bolch, T., Chaulagain, N.P., Dhaubanjari, S., Dimri, A.P., Dixon, H., 2022. Knowledge priorities on climate change and water in the Upper Indus Basin: A horizon scanning exercise to identify the top 100 research questions in social and natural sciences. *Earth's Future*, **10**: e2021EF002619. <https://doi.org/10.1029/2021EF002619>
- Palmeirim, A.F. and Gibson, L., 2021. Impacts of hydropower on the habitat of jaguars and tigers. *Commun. Biol.*, **4**: 1358. <https://doi.org/10.1038/s42003-021-02878-5>
- Pandit, M.K. and Grumbine, R.E., 2012. Potential effects of ongoing and proposed hydropower development on terrestrial biological diversity in the Indian Himalaya. *Conserv. Biol.*, **26**: 1061-1071. <https://doi.org/10.1111/j.1523-1739.2012.01918.x>
- Qureshi, A.S., 2011. Water management in the Indus

- Basin in Pakistan: Challenges and opportunities. *Mt. Res. Dev.*, **31**: 252–260. <https://doi.org/10.1659/MRD-JOURNAL-D-11-00019.1>
- Ramachandran, P. and Linde, L., 2011. Integrating spatial support tools into strategic planning SEA of the GMS North–South economic corridor strategy and action plan. *Environ. Impact Assess. Rev.*, **31**: 602–611. <https://doi.org/10.1016/j.eiar.2010.04.002>
- Roberts, T.J., 1997. *The mammals of Pakistan*. Revised Edition. Karachi: Oxford University Press. pp. 525.
- Roberts, T.J., 1991. *The birds of Pakistan*. Volume 1. Oxford University Press, Karachi. pp. 598.
- Roberts, T.J., 1992. *The birds of Pakistan*. Volume 2. Oxford University Press, Karachi. pp. 617.
- Sackey, L.N., Quartey, J., Nortey, A.N., Obeng, A.T., Okyere, A.A. and Kayoung, P.Y., 2023. Road construction and its socio-economic and health impact: A case study of Atonsu lake road. *Discover Environment.*, **1**: 6. <https://doi.org/10.1007/s44274-023-00009-x>
- Sharma, A.K. and Thakur, N.S., 2017. Energy situation, current status and resource potential of run of the river (RoR) large hydro power projects in Jammu and Kashmir: India. *Renew. Sustain. Energy Rev.*, **78**: 233–251. <https://doi.org/10.1016/j.rser.2017.04.087>
- Sorensen, T., 1948. A Method of establishing groups of equal amplitude in plant sociology based on similarity of species content and its application to analyses of the vegetation on danish commons. *Biologiske Skrifter/Kongelige Danske Videnskabernes Selskab*, **5**: 1–34.
- Ullah, E.I., Ahmad, S., Khokhar, M.F., Azmat, M. and Khayyam, U., 2023. Hydrological and ecological impacts of run off river scheme; a case study of Ghazi Barotha hydropower project on Indus River, Pakistan. *Heliyon*, **9**: e12659. <https://doi.org/10.1016/j.heliyon.2022.e12659>
- Valenzuela-Aguayo, F., McCracken, G.R., Manosalva, A., Habit, E. and Ruzzante, D.E., 2020. Human-induced habitat fragmentation effects on connectivity, diversity, and population persistence of an endemic fish, *Percilia irwini*, in the Biobío River basin (Chile). *Evol. appl.*, **13**: 794–807. <https://doi.org/10.1111/eva.12901>
- Wester, P., Mishra, A., Mukherji, A. and Shrestha, A.B., 2019. The Hindu Kush Himalaya assessment: Mountains, climate change, sustainability and people. *Springer Nature*, pp. 627. <https://doi.org/10.1007/978-3-319-92288-1>
- Xiaolong, T., Gull, N., Iqbal, S., Asghar, M., Nawaz, A., Albasher, G., Hameed, J. and Maqsoom, A., 2021. Exploring and validating the effects of mega projects on infrastructure development influencing sustainable environment and project management. *Front. Psychol.*, **12**: 663199. <https://doi.org/10.3389/fpsyg.2021.663199>
- Xu, J., Badola, R., Chettri, N., Chaudhary, R.P., Zomer, R., Pokhrel, B., Hussain, S.A., Pradhan, S. and Pradhan, R., 2019. Sustaining biodiversity and ecosystem services in the Hindu Kush Himalaya. In: *The Hindu Kush Himalaya assessment* (eds. P. Wester, A. Mishra, A. Mukherji and A. Shrestha). Springer, Cham. https://doi.org/10.1007/978-3-319-92288-1_5
- Yaseen, M., Ahmad, I., Guo, J., Azam, M.I. and Latif, Y., 2020. Spatiotemporal variability in the hydrometeorological time-series over Upper Indus River Basin of Pakistan. *Adv. Meteorol.*, Article ID 5852760. <https://doi.org/10.1155/2020/5852760>
- Zahra, S.M., Shahid, M.A., Misal, M.A., Zaman, M., Imran, M., Azam, S. and Hussain, F., 2023. Sustainable water management in Indus Basin and vulnerability due to climate change. *Environ. Sci. Proc.*, **25**: 36. <https://doi.org/10.3390/ECWS-7-14203>
- Zhang, L. and Wang, J., 2023. Mountain biodiversity, species distribution and ecosystem functioning in a changing world. *Diversity*, **15**: 799. <https://doi.org/10.3390/d15070799>



Supplementary Material

Assessing Wildlife and Ecology Amid Infrastructure Development in the Upper Indus Basin, Pakistan

Usman Ahmad¹, Sajid Rashid Ahmad^{1*}, Zulfiqar Ali^{2*} and Rida Ahmad^{2,3}

¹College of Earth and Environmental Sciences, University of the Punjab, Lahore

²Institute of Zoology, University of the Punjab, Lahore

³Department of Animal Sciences, University of the Punjab, Lahore, Pakistan

Supplementary Table I. Species recorded at the study site.

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|------------------|-----------------|---------------------------------------|--------------------------------------|---------------|---------------|------------|-------|
| Amphibian | | | | | | | |
| Anura | Bufonidae | Green toad | <i>Bufo viridis</i> | LC | Decreasing | YRR | 1 |
| | | Hazara toad | <i>Bufo melanostictus hazarensis</i> | Not available | Not available | YRR | 2 |
| | | Himalayan toad | <i>Bufo himalayanus</i> | LC | Stable | YRR | 3 |
| | | Indus valley toad | <i>Bufo stomaticus</i> | LC | Stable | YRR | 3 |
| | | Ladakh toad | <i>Bufo latastii</i> | LC | Unknown | YRR | 2 |
| | | Siachine toad | <i>Bufo siachenensis</i> | Not available | Not available | YRR | 1 |
| | | Southeast Asian toad | <i>Bufo melanostictus</i> | LC | Increasing | YRR | 1 |
| | Swat green toad | <i>Bufo pseudoraddei pseudoraddei</i> | LC | Stable | YRR | 1 | |
| | Dicroglossidae | Alpine cricket frog | <i>Fejervarya limnocharis</i> | LC | Stable | YRR | 1 |
| | | Hazara fast-flowing stream frog | <i>Paa hazarensis</i> | LC | Stable | YRR | 3 |
| | | Hazara torrent frog | <i>Allopaa hazarensis</i> | LC | Stable | YRR | 1 |
| | | Skittering frog | <i>Euphlyctis cyanophlyctis</i> | LC | Stable | YRR | 2 |
| | Avifauna | | | | | | |
| Accipitri-formes | Accipitridae | Black kite | <i>Milvus migrans</i> | LC | Unknown | SB/YRR | 157 |
| | | Bonelli's eagle | <i>Hieraaetus fasciatus</i> | LC | Decreasing | WM/YRR | 1 |
| | | Booted eagle | <i>Hieraaetus pennatus</i> | LC | Unknown | WM/YRR | 19 |

Table continued on next page.....

* Corresponding author: zali.zool@pu.edu.pk, sajidpu@yahoo.com
0030-9923/2024/0001-0001 \$ 9.00/0



| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|----------------------|---------------|------------------------|---------------------------------|-------------|------------|---------------|-------|
| | | Egyptian vulture | <i>Neophron percnopterus</i> | EN | Decreasing | SB/YRR | 7 |
| | | Eurasian buzzard | <i>Buteo buteo</i> | LC | Stable | WM | 1 |
| | | Eurasian marsh harrier | <i>Circus aeruginosus</i> | LC | Increasing | WM/PM | 21 |
| | | Eurasian sparrowhawk | <i>Accipiter nisus</i> | LC | Stable | SB/WM | 4 |
| | | Hen harrier | <i>Circus cyaneus</i> | LC | Decreasing | WM | 7 |
| | | Himalayan griffon | <i>Gyps himalayensis</i> | NT | Stable | YRR | 8 |
| | | Indian spotted eagle | <i>Aquila hastata</i> | VU | Decreasing | Not available | 3 |
| | | Lammergeier | <i>Gypaetus barbatus</i> | NT | Decreasing | WM/YRR | 6 |
| | | Long-legged buzzard | <i>Buteo rufinus</i> | LC | Stable | WM | 3 |
| | | Pallas's fish eagle | <i>Haliaeetus leucoryphus</i> | EN | Decreasing | YRR | 9 |
| | | Pallid harrier | <i>Circus macrourus</i> | NT | Decreasing | WM/PM | 8 |
| | | Shikra | <i>Accipiter badius</i> | LC | Stable | SB/YRR | 29 |
| | | White-eyed buzzard | <i>Butastur teesa</i> | LC | Stable | YRR/SB | 2 |
| | | White-rumped vulture | <i>Gyps bengalensis</i> | CR | Decreasing | YRR | 4 |
| Anseri- formes | Anatidae | Bar-headed goose | <i>Anser indicus</i> | LC | Decreasing | PM/ IYRV | 371 |
| | | Common merganser | <i>Mergus merganser</i> | LC | Increasing | WM | 11 |
| | | Common teal | <i>Anas crecca</i> | LC | Unknown | WM | 357 |
| | | Eurasian wigeon | <i>Anas penelope</i> | LC | Decreasing | WM | 23 |
| | | Ferruginous pochard | <i>Aythya nyroca</i> | NT | Decreasing | PM/WM | 14 |
| | | Gadwall | <i>Anas strepera</i> | LC | Increasing | WM | 84 |
| | | Greylag goose | <i>Anser anser</i> | LC | Increasing | PM/ IYRV | 53 |
| | | Mallard | <i>Anas platyrhynchos</i> | LC | Increasing | WM | 42 |
| | | Northern pintail | <i>Anas acuta</i> | LC | Decreasing | WM | 216 |
| | | Northern shoveler | <i>Anas clypeata</i> | LC | Decreasing | WM | 94 |
| | | Ruddy shelduck | <i>Tadorna ferruginea</i> | LC | Unknown | PM/ IYRV/* | 37 |
| | | Tufted duck | <i>Aythya fuligula</i> | LC | Stable | PM/WM | 264 |
| Apodi- formes | Apodidae | Alpine swift | <i>Tachymartitis melba</i> | LC | Stable | SB | 26 |
| Buceroti- formes | Upupidae | Common hoopoe | <i>Upupa epops</i> | LC | Decreasing | SB/YRR/ WM | 53 |
| Caprimul- iformes | Apodidae | Common swift | <i>Apus apus</i> | LC | Stable | SB | 14 |
| | | House swift | <i>Apus affinis</i> | LC | Increasing | SB/YRR | 12 |
| | Caprimulgidae | Eurasian nightjar | <i>Caprimulgus europaeus</i> | LC | Decreasing | SB/PM | 4 |
| Charadrii- formes | Charadriidae | Little ringed plover | <i>Charadrius dubius</i> | LC | Stable | SB/WM | 5 |
| | | Northern lapwing | <i>Vanellus vanellus</i> | NT | Decreasing | WM | 41 |
| | | Red-wattled lapwing | <i>Vanellus indicus</i> | LC | Stable | SB/YRR | 22 |
| | | Sociable lapwing | <i>Vanellus gregarius</i> | CR | Decreasing | PM/WM | 3 |
| | Jacanidae | Pheasant-tailed jacana | <i>Hydrophasianus chirurgus</i> | LC | Decreasing | SB/YRR | 1 |
| | Laridae | Black-headed gull | <i>Larus ridibundus</i> | LC | Unknown | PM/WM | 2357 |
| | | Brown-headed gull | <i>Larus brunnicephalus</i> | LC | Stable | PM/WM | 242 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|---------------|------------------|---------------------------|----------------------------------|-------------|------------|------------|-------|
| | | Caspian tern | <i>Sterna caspia</i> | LC | Increasing | PM | 41 |
| | | Gull-billed tern | <i>Gelochelidon nilotica</i> | LC | Decreasing | PM/WM | 138 |
| | | Indian skimmer | <i>Rynchops albicollis</i> | VU | Decreasing | SB | 14 |
| | | Mew gull | <i>Larus canus</i> | LC | Unknown | WM | 38 |
| | Recurvirostridae | Pied avocet | <i>Recurvirostra avosetta</i> | LC | Unknown | WM | 3 |
| | Scolopacidae | Ruff | <i>Philomachus pugnax</i> | LC | Decreasing | PM | 6 |
| | | Common redshank | <i>Tringa totanus</i> | LC | Unknown | PM/WM | 1 |
| | | Common sandpiper | <i>Actitis hypoleucos</i> | LC | Decreasing | SB/WM | 22 |
| | | Common snipe | <i>Gallinago gallinago</i> | LC | Decreasing | PM/WM | 3 |
| | | Curlew sandpiper | <i>Calidris ferruginea</i> | NT | Decreasing | PM/WM | 14 |
| | | Eurasian curlew | <i>Numenius arquata</i> | NT | Decreasing | PM/WM | 1 |
| | | Eurasian woodcock | <i>Scolopax rusticola</i> | LC | Stable | SB/WM | 1 |
| | | Green sandpiper | <i>Tringa ochropus</i> | LC | Increasing | PM/WM | 9 |
| | | Jack snipe | <i>Lymnocyptes minimus</i> | LC | Stable | PM/WM | 5 |
| | | Little stint | <i>Calidris minuta</i> | LC | Increasing | PM/WM | 7 |
| | | Temminck's stint | <i>Calidris temminckii</i> | LC | Unknown | PM/WM | 3 |
| | | Wood sandpiper | <i>Tringa glareola</i> | LC | Stable | PM/WM | 17 |
| | Turnicidae | Small buttonquail | <i>Turnix sylvatica</i> | LC | Decreasing | SB | 18 |
| Ciconiiformes | Ciconiidae | Black stork | <i>Ciconia nigra</i> | LC | Unknown | PM | 1 |
| Columbiformes | Columbidae | European turtle dove | <i>Streptopelia turtur</i> | VU | Decreasing | PM | 14 |
| | | Hill pigeon | <i>Columba rupestris</i> | LC | Decreasing | SB/WM | 22 |
| | | Rock pigeon | <i>Columba livia</i> | LC | Decreasing | YRR | 63 |
| | Columbidae | Eurasian collared dove | <i>Streptopelia decaocto</i> | LC | Increasing | SB/YRR | 23 |
| | | Laughing dove | <i>Streptopelia senegalensis</i> | LC | Stable | YRR/SB | 27 |
| | | Oriental turtle dove | <i>Streptopelia orientalis</i> | LC | Stable | YRR/WM | 3 |
| | | Spotted dove | <i>Streptopelia chinensis</i> | LC | Increasing | SB/WM | 6 |
| Coraciiformes | Alcedinidae | Common kingfisher | <i>Alcedo atthis</i> | LC | Unknown | SB/YRR/WM | 49 |
| | | White-throated kingfisher | <i>Halcyon smyrnensis</i> | LC | Increasing | YRR | 19 |
| | Coraciidae | European roller | <i>Coracias garrulus</i> | LC | Decreasing | PM/SB | 47 |
| | | Indian roller | <i>Coracias benghalensis</i> | LC | Increasing | SB/YRR | 32 |
| | Meropidae | Blue-cheeked bee-eater | <i>Merops persicus</i> | LC | Stable | PM/SB | 13 |
| | | European bee-eater | <i>Merops apiaster</i> | LC | Stable | SB | 6 |
| | | Green bee-eater | <i>Merops orientalis</i> | LC | Increasing | SB/YRR | 24 |
| Cuculiformes | Cuculidae | Pied cuckoos | <i>Clamator jacobinus</i> | LC | Stable | SB | 4 |
| | | Eurasian cuckoo | <i>Cuculus canorus</i> | LC | Decreasing | SB | 1 |
| Falconiformes | Falconidae | Common kestrel | <i>Falco tinnunculus</i> | LC | Decreasing | WM/YRR | 18 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count | | |
|-------------------------|-------------------------------|---------------------------|----------------------------------|---------------------------------|-------------------------------|------------|------------|-------|---|
| Galliformes | Phasianidae | Eurasian hobby | <i>Falco subbuteo</i> | LC | Decreasing | SB/WM | 7 | | |
| | | Peregrine falcon | <i>Falco peregrinus</i> | LC | Stable | SB/WM | 1 | | |
| | | Saker falcon | <i>Falco cherrug</i> | EN | Decreasing | WM | 2 | | |
| | | Cheer pheasant | <i>Catreus wallichii</i> | VU | Decreasing | YRR | 1 | | |
| | | Chukar | <i>Alectoris chukar</i> | LC | Stable | YRR | 14 | | |
| | | Common quail | <i>Coturnix coturnix</i> | LC | Decreasing | PM/ IYRV | 12 | | |
| | | Grey francolin | <i>Francolinus pondicerianus</i> | LC | Stable | YRR | 4 | | |
| | | Himalayan monal | <i>Lophophorus impejanus</i> | LC | Decreasing | YRR | 4 | | |
| | | Himalayan snowcock | <i>Tetraogallus himalayensis</i> | LC | Stable | YRR | 1 | | |
| | | Kalij pheasant | <i>Lophura leucomelanos</i> | LC | Decreasing | YRR | 7 | | |
| | | Koklass pheasant | <i>Pucrasia macrolopha</i> | LC | Decreasing | YRR | 3 | | |
| | | See-see partridge | <i>Ammoperdix griseogularis</i> | LC | Stable | YRR | 6 | | |
| | | Snow partridge | <i>Lerwa lerwa</i> | LC | Decreasing | YRR | 4 | | |
| | | Western tragopan | <i>Tragopan melanocephalus</i> | VU | Decreasing | YRR | 2 | | |
| | | Gruiformes | Rallidae | Baillon's crane | <i>Porzana pusilla</i> | LC | Unknown | PM/WM | 1 |
| Little crane | <i>Porzana parva</i> | | | LC | Stable | PM/WM | 3 | | |
| White-breasted waterhen | <i>Amaurornis phoenicurus</i> | | | LC | Unknown | PM/WM | 59 | | |
| Common coot | <i>Fulica atra</i> | | | LC | Increasing | WM | 67 | | |
| Common moorhen | <i>Gallinula chloropus</i> | | | LC | Stable | YRR | 11 | | |
| Otidiformes | Otididae | Water rail | <i>Rallus aquaticus</i> | LC | Decreasing | PM/WM | 7 | | |
| | | Little bustard | <i>Tetrax tetrax</i> | NT | Decreasing | WM | 2 | | |
| | | Passeri- formes | Acrocephalidae | Blyth's reed warbler | <i>Acrocephalus dumetorum</i> | LC | Increasing | PM | 7 |
| | | | Aegithalidae | White-browed tit warbler | <i>Leptopoecile sophiae</i> | LC | Stable | SB/WM | 2 |
| | | Alaudidae | White-cheeked tit | <i>Aegithalos leucogenys</i> | LC | Stable | YRR/WM | 3 | |
| | | | Bimaculated lark | <i>Melanocorypha bimaculata</i> | LC | Stable | WM | 2 | |
| | | | Crested lark | <i>Galerida cristata</i> | LC | Decreasing | YRR | 29 | |
| | | | Desert lark | <i>Ammomanes deserti</i> | LC | Increasing | YRR | 8 | |
| | | | Eurasian skylark | <i>Alauda arvensis</i> | LC | Decreasing | WM | 6 | |
| | | | Oriental skylark | <i>Alauda gulgula</i> | LC | Decreasing | SB/YRR | 2 | |
| Campephagidae | Long-tailed minivet | | <i>Pericrocotus ethologus</i> | LC | Decreasing | SB/WM | 5 | | |
| | Rosy minivet | | <i>Pericrocotus roseus</i> | LC | Decreasing | YRR | 7 | | |
| Certhiidae | Eurasian treecreeper | <i>Certhia familiaris</i> | LC | Stable | YRR | 1 | | | |
| Cinclidae | Brown dipper | <i>Cinclus pallasii</i> | LC | Stable | YRR | 167 | | | |
| | White-throated dipper | <i>Cinclus cinclus</i> | LC | Decreasing | YRR | 49 | | | |
| Cisticolidae | Grey-breasted prinia | <i>Prinia hodgsonii</i> | LC | Stable | YRR | 1 | | | |
| | Striated prinia | <i>Prinia crinigera</i> | LC | Stable | YRR | 3 | | | |
| | Zitting cisticola | <i>Cisticola juncidis</i> | LC | Increasing | YRR | 14 | | | |
| Corvidae | Carrion crow | <i>Corvus corone</i> | LC | Increasing | WM/YRR | 2149 | | | |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|-------|-----------------|-----------------------------|-------------------------------|-------------|------------|---------------|-------|
| | | Common raven | <i>Corvus corax</i> | LC | Increasing | YRR | 274 |
| | | House crow | <i>Corvus splendens</i> | LC | Stable | YRR | 1124 |
| | | Large-billed crow | <i>Corvus macrorhynchos</i> | LC | Stable | WM/YRR | 1678 |
| | | Rook | <i>Corvus frugilegus</i> | LC | Decreasing | WM | 11 |
| | | Yellow-billed blue magpie | <i>Urocissa flavirostris</i> | LC | Stable | YRR | 18 |
| | | Yellow-billed chough | <i>Pyrrhonorax graculus</i> | LC | Stable | YRR | 4 |
| | Dicruridae | Ashy drongo | <i>Dicrurus leucophaeus</i> | LC | Unknown | SB | 37 |
| | | Black drongo | <i>Dicrurus macrocercus</i> | LC | Unknown | SB/PM/YRR | 138 |
| | Emberizidae | Chestnut bunting | <i>Emberiza rutila</i> | LC | Stable | Not available | 11 |
| | | Reed bunting | <i>Emberiza schoeniclus</i> | LC | Decreasing | WM | 4 |
| | | Rock bunting | <i>Emberiza cia</i> | LC | Increasing | SB/WM | 417 |
| | Estrildidae | Red avadavat | <i>Amandava amandava</i> | LC | Stable | SB/YRR | 1 |
| | Fringillidae | Chaffinch | <i>Fringilla coelebs</i> | LC | Increasing | WM | 1 |
| | | Common rosefinch | <i>Carpodacus erythrinus</i> | LC | Decreasing | SB/PM | 9 |
| | | Crested bunting | <i>Melophus lathami</i> | LC | Stable | SB | 7 |
| | | Eurasian goldfinch | <i>Carduelis carduelis</i> | LC | Decreasing | SB/WM | 1 |
| | | Orange bullfinch | <i>Pyrrhula aurantiaca</i> | LC | Stable | WM/YRR | 4 |
| | | Red-fronted rosefinch | <i>Carpodacus puniceus</i> | LC | Stable | YRR | 5 |
| | | Twite | <i>Carduelis flavirostris</i> | LC | Decreasing | SB/WM | 1 |
| | | White-winged grosbeak | <i>Mycerobas carnipes</i> | LC | Stable | YRR | 2 |
| | | Yellow-breasted green-finch | <i>Carduelis spinoides</i> | LC | Stable | SB/WM | 3 |
| | Hirundinidae | Asian house martin | <i>Delichon dasypus</i> | LC | Increasing | SB | 137 |
| | | Barn swallow | <i>Hirundo rustica</i> | LC | Decreasing | SB/WM | 44 |
| | | Eurasian crag martin | <i>Hirundo rupestris</i> | LC | Stable | SB/WM | 4 |
| | | Northern house martin | <i>Delichon urbicum</i> | LC | Decreasing | SB | 14 |
| | | Pale martin | <i>Riparia diluta</i> | LC | Unknown | SB/WM | 6 |
| | | Wire-tailed swallow | <i>Hirundo smithii</i> | LC | Increasing | SB/YRR | 49 |
| | Laniidae | Great grey shrike | <i>Lanius excubitor</i> | LC | Stable | Not available | 7 |
| | | Long-tailed shrike | <i>Lanius schach</i> | LC | Unknown | SB/YRR | 38 |
| | | Rufous-tailed shrike | <i>Lanius isabellinus</i> | LC | Stable | SB/WM | 11 |
| | | Southern grey shrike | <i>Lanius meridionalis</i> | VU | Decreasing | SB/YRR | 7 |
| | Leiotherichidae | Jungle babbler | <i>Turdoides striatus</i> | LC | Stable | YRR | 36 |
| | | Streaked laughingthrush | <i>Garrulax lineatus</i> | LC | Stable | YRR | 14 |
| | Locustellidae | Long-billed bush warbler | <i>Bradypterus major</i> | LC | Decreasing | SB | 6 |
| | Monarchidae | Asian paradise-fly-catcher | <i>Terpsiphone paradisi</i> | LC | Stable | SB/PM | 19 |
| | Motacillidae | Citrine wagtail | <i>Motacilla citreola</i> | LC | Increasing | SB/WM | 74 |
| | | Grey wagtail | <i>Motacilla cinerea</i> | LC | Stable | SB/WM | 698 |
| | | Long-billed pipit | <i>Anthus similis</i> | LC | Stable | WM/YRR | 19 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|-------|---------------|-----------------------------|---|-------------|------------|------------|-------|
| | | Paddyfield pipit | <i>Anthus rufulus</i> | LC | Stable | SB/YRR | 87 |
| | | Rosy pipit | <i>Anthus roseatus</i> | LC | Stable | SB/WM | 4 |
| | | Tawny pipit | <i>Anthus campestris</i> | LC | Stable | WM | 24 |
| | | Tree pipit | <i>Anthus trivialis</i> | LC | Decreasing | SB/PM | 8 |
| | | Upland pipit | <i>Anthus sylvanus</i> | LC | Stable | YRR | 3 |
| | | Water pipit | <i>Anthus spinoletta</i> | LC | Stable | WM | 37 |
| | | White wagtail | <i>Motacilla alba</i> | LC | Stable | SB/WM | 742 |
| | | Yellow wagtail | <i>Motacilla flava</i> | LC | Decreasing | PM | 287 |
| | Muscicapidae | Black redstart | <i>Phoenicurus ochruros</i> | LC | Increasing | SB/WM | 1 |
| | | Blue rock thrush | <i>Monticola solitarius</i> | LC | Stable | SB/WM | 32 |
| | | Blue whistling thrush | <i>Myophonus caeruleus</i> | LC | Unknown | SB/WM | 314 |
| | | Blue-capped redstart | <i>Phoenicurus coeruleocephalus</i> | LC | Stable | SB/WM | 23 |
| | | Blue-capped rock thrush | <i>Monticola cinclorhynchus</i> | LC | Stable | SB/PM | 14 |
| | | Blue-fronted redstart | <i>Phoenicurus frontalis</i> | LC | Stable | YRR | 1 |
| | | Bluethroat | <i>Luscinia svecica</i> | LC | Stable | SB/WM/PM | 4 |
| | | Common stonechat | <i>Saxicola torquata</i> | LC | Stable | SB/PM | 37 |
| | | Grey bushchat | <i>Saxicola ferrea</i> | LC | Stable | SB/WM | 12 |
| | | Hume's wheatear | <i>Oenanthe alboniger</i> | LC | Stable | YRR/WM | 182 |
| | | Indian blue robin | <i>Luscinia brunnea</i> | LC | Decreasing | SB/PM | 16 |
| | | Indian robin | <i>Saxicoloides fulicata</i> | LC | Stable | YRR | 38 |
| | | Isabelline wheatear | <i>Oenanthe isabellina</i> | LC | Stable | SB/WM/PM | 1 |
| | | Kashmir flycatcher | <i>Ficedula subrubra</i> | VU | Decreasing | SB | 6 |
| | | Little forktail | <i>Enicurus scouleri</i> | LC | Stable | YRR/WM | 1 |
| | | Pied buschhat | <i>Saxicola caprata</i> | LC | Stable | SB/YRR | 18 |
| | | Pied wheatear | <i>Oenanthe pleschanka</i> | LC | Stable | SB/PM | 47 |
| | | Plumbeous water redstart | <i>Rhyacornis fuliginosus</i> | LC | Stable | SB/WM | 924 |
| | | Red-throated flycatcher | <i>Ficedula parva</i> | LC | Increasing | WM/PM | 1 |
| | | Rufous-backed redstart | <i>Phoenicurus erythronota</i> | LC | Stable | WM | 4 |
| | | Rufous-tailed rock thrush | <i>Monticola saxatilis</i> | LC | Decreasing | PM | 19 |
| | | Slaty-blue flycatcher | <i>Ficedula tricolor</i> | LC | Stable | SB/WM | 1 |
| | | Ultramarine flycatcher | <i>Ficedula superciliaris</i> | LC | Stable | SB/WM | 3 |
| | | Variable wheatear | <i>Oenanthe picata</i> | LC | Stable | SB/WM | 358 |
| | | White-bellied redstart | <i>Hodgsonius phaenicuroides</i> | LC | Stable | YRR | 3 |
| | | White-capped water redstart | <i>Chaimarrornis leucocephalus</i> | LC | Stable | SB/WM | 892 |
| | | White-winged redstart | <i>Phoenicurus erythrogaster</i> | LC | Stable | SB/WM | 2 |
| | Nectariniidae | Purple sunbird | <i>Nectarinia asiatica</i> | LC | Stable | SB/YRR | 27 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|---------------------|-------------------------|------------------------------|----------------------------------|-------------|------------|---------------|-------|
| | Oriolidae | Eurasian golden oriole | <i>Oriolus oriolus</i> | LC | Stable | SB/PM | 9 |
| | Paridae | Great tit | <i>Parus major</i> | LC | Increasing | YRR/WM | 15 |
| | | Rufous-naped tit | <i>Parus rufonuchalis</i> | LC | Stable | YRR | 18 |
| | Passeridae | Eurasian tree sparrow | <i>Passer montanus</i> | LC | Decreasing | WM/YRR | 7 |
| | | House sparrow | <i>Passer domesticus</i> | LC | Decreasing | SB/YRR/PM | 247 |
| | | Russet sparrow | <i>Passer rutilans</i> | LC | Stable | WM/YRR | 14 |
| | | Spanish sparrow | <i>Passer hispaniolensis</i> | LC | Decreasing | WM/PM | 252 |
| | Phylloscopidae | Blyth's leaf warbler | <i>Phylloscopus reguloides</i> | LC | Stable | Not available | 3 |
| | | Brook's leaf warbler | <i>Phylloscopus subviridis</i> | LC | Stable | SB/WM | 3 |
| | | Common chiffchaff | <i>Phylloscopus collybita</i> | LC | Increasing | WM | 9 |
| | | Greenish warbler | <i>Phylloscopus trochiloides</i> | LC | Increasing | SB/WM/PM | 6 |
| | | Hume's warbler | <i>Phylloscopus humei</i> | LC | Stable | SB/WM | 5 |
| | | Mountain chiffchaff | <i>Phylloscopus sindianus</i> | LC | Stable | SB/WM | 5 |
| | | Tickell's leaf warbler | <i>Phylloscopus affinis</i> | LC | Stable | SB | 3 |
| | | Tytler's leaf warbler | <i>Phylloscopus tytleri</i> | NT | Decreasing | SB/PM | 1 |
| | Prunellidae | Alpine accentor | <i>Prunella collaris</i> | LC | Stable | SB/WM | 16 |
| | Pycnonotidae | Himalayan bulbul | <i>Pycnonotus leucogenys</i> | LC | Increasing | YRR | 214 |
| | | Red-vented bulbul | <i>Pycnonotus cafer</i> | LC | Increasing | YRR | 321 |
| | Rhipiduridae | White-throated fantail | <i>Rhipidura albicollis</i> | LC | Stable | YRR | 1 |
| | Sittidae | Wallcreeper | <i>Tichodroma muraria</i> | LC | Stable | YRR/WM | 9 |
| | Sturnidae | Brahminy starling | <i>Sturnus pagodarum</i> | LC | Unknown | YRR | 2 |
| | | Common myna | <i>Acridotheres tristis</i> | LC | Increasing | SB/YRR | 426 |
| | Sylviidae | Lesser whitethroat | <i>Sylvia curruca</i> | LC | Stable | SB/WM | 1 |
| | Troglodytidae | Winter wren | <i>Troglodytes troglodytes</i> | LC | Increasing | YRR | 17 |
| | Turdidae | Chestnut thrush | <i>Turdus rubrocanus</i> | LC | Unknown | WM/YRR | 3 |
| | | Eurasian blackbird | <i>Turdus merula</i> | LC | Increasing | YRR | 1 |
| | | Tickell's thrush | <i>Turdus unicolor</i> | LC | Unknown | SB/WM | 2 |
| | Zosteropidae | Oriental white-eye | <i>Zosterops palpebrosus</i> | LC | Decreasing | YRR | 7 |
| Pelecani- formes | Ardeidae | Cattle egret | <i>Bubulcus ibis</i> | LC | Increasing | YRR | 124 |
| | | Grey heron | <i>Ardea cinerea</i> | LC | Unknown | WM/PM/YRR | 74 |
| | | Indian pond-heron | <i>Ardeola grayii</i> | LC | Unknown | YRR | 138 |
| | | Little bittern | <i>Ixobrychus minutus</i> | LC | Decreasing | PM/WM | 8 |
| | | Little egret | <i>Egretta garzetta</i> | LC | Increasing | PM/YRR | 57 |
| | Threskiornithi- idae | Eurasian spoonbill | <i>Platalea leucorodia</i> | LC | Unknown | PM/WM | 2 |
| | | Glossy ibis | <i>Plegadis falcinellus</i> | LC | Decreasing | PM | 2 |
| Piciformes | Megalaimidae | Great barbet | <i>Megalaima virens</i> | LC | Decreasing | SB/YRR | 1 |
| | Picidae | Eurasian wryneck | <i>Jynx torquilla</i> | LC | Decreasing | SB/PM/WM | 2 |
| | | Grey-capped pygmy woodpecker | <i>Dendrocopos canicapillus</i> | LC | Stable | SB | 1 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|-----------------------|------------------------|-----------------------------|-----------------------------------|---------------|---------------|---------------|-------|
| | | Himalayan woodpecker | <i>Dendrocopos himalayensis</i> | LC | Stable | YRR | 3 |
| Podicipedi- formes | Podicipedidae | Great crested grebe | <i>Podiceps cristatus</i> | LC | Unknown | WM | 3 |
| | | Little grebe | <i>Tachybaptus ruficollis</i> | LC | Decreasing | SB/YRR | 34 |
| Psittaci- formes | Psittacidae | Alexandrine parakeet | <i>Psittacula eupatria</i> | NT | Decreasing | YRR | 4 |
| | | Rose-ringed parakeet | <i>Psittacula krameri</i> | LC | Increasing | YRR | 3 |
| | | Slaty-headed parakeet | <i>Psittacula himalayana</i> | LC | Stable | YRR | 1 |
| Pterocli- formes | Pteroclididae | Chestnut-bellied sandgrouse | <i>Pterocles exustus</i> | LC | Stable | YRR | 4 |
| Strigi- formes | Strigidae | Mountain scops owl | <i>Otus spilocephalus</i> | LC | Stable | YRR | 1 |
| | | Snowy owl | <i>Nyctea scandiaca</i> | VU | Decreasing | Not available | 1 |
| | Strigidae | Asian barred owl | <i>Glaucidium cuculoides</i> | LC | Increasing | YRR | 1 |
| | | Collared owl | <i>Glaucidium brodiei</i> | LC | Decreasing | SB/WM | 2 |
| | | Eurasian scops owl | <i>Otus scops</i> | LC | Decreasing | SB/WM | 1 |
| | | Pallid scops owl | <i>Otus brucei</i> | LC | Stable | SB/WM | 1 |
| | | Tawny owl | <i>Strix aluco</i> | LC | Stable | YRR/WM | 1 |
| Suliformes | Phalacrocoraci- dae | Great cormorant | <i>Phalacrocorax carbo</i> | LC | Increasing | PM/WM | 3274 |
| Mammals | | | | | | | |
| Artiodac- tyla | Bovidae | Punjab urial | <i>Ovis vignei punjabensis</i> | VU | Decreasing | Not available | 8 |
| | | Shapu | <i>Ovis vignei vignei</i> | VU | Decreasing | Not available | 2 |
| Carnivora | Canidae | Desert fox | <i>Vulpes vulpes pusilla</i> | Not available | Not available | Not available | 1 |
| | | Indian wild dog | <i>Cuon alpinus</i> | EN | Decreasing | Not available | 1 |
| | Felidae | Jungle cat | <i>Felis chaus</i> | LC | Decreasing | Not available | 1 |
| | | Leopard cat | <i>Prionailurus bengalensis</i> | LC | Stable | Not available | 1 |
| | | Pallas' cat | <i>Felis manul</i> | NT | Decreasing | Not available | 1 |
| | | Panther | <i>Panthera pardus</i> | VU | Decreasing | Not available | 4 |
| | | Snow leopard | <i>Panthera uncia</i> | VU | Decreasing | Not available | 2 |
| | Herpestidae | Common indian mongoose | <i>Herpestes edwardsi</i> | LC | Stable | Not available | 7 |
| | Hyaenidae | Striped hyaena | <i>Hyaena hyaena</i> | NT | Decreasing | Not available | 1 |
| | Mustelidae | Alpine weasel | <i>Mustela altaica</i> | NT | Decreasing | Not available | 1 |
| | | Common otter | <i>Lutra lutra</i> | NT | Decreasing | Not available | 3 |
| | | Indian otter | <i>Lutrogale perspicillata</i> | VU | Decreasing | Not available | 1 |
| | Ursidae | Asiatic black bear | <i>Ursus thibetanus</i> | VU | Decreasing | Not available | 4 |
| | Viverricula | Common palm civet | <i>Paradoxurus hermaphroditus</i> | LC | Decreasing | Not available | 1 |
| | | Himalayan palm civet | <i>Paguma larvata wroughtoni</i> | LC | Decreasing | Not available | 2 |
| Cetartio- dactyla | Bovidae | Astor markhor | <i>Capra falconeri</i> | NT | Increasing | Not available | 7 |
| | | Himalayan ibex | <i>Capra ibex siberica</i> | LC | Increasing | Not available | 1 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|-----------------|------------------|-------------------------------|---|---------------|---------------|---------------|-------|
| | | Pir panjal markhor | <i>Capra falconeri cashmiriensis</i> | Not Available | Not available | Not available | 4 |
| | Moschidae | Himalayan musk deer | <i>Moschus chrysogaster</i> | EN | Decreasing | Not available | 3 |
| | Suidae | Indian wild boar | <i>Sus scrofa</i> | LC | Unknown | Not available | 8 |
| Chiroptera | Pteropodidae | Indian flying fox | <i>Pteropus giganteus</i> | LC | Decreasing | Not available | 4 |
| | Rhinopomatidae | Big-eared horseshoe bat | <i>Rhinolophus macrotis</i> | LC | Stable | Not available | 2 |
| | Vespertilionidae | Asian barbastelle | <i>Barbastella leucomelas</i> | LC | Unknown | Not available | 2 |
| | | Common pipistrelle | <i>Pipistrellus pipistrellus</i> | LC | Stable | Not available | 1 |
| Lagomorpha | Leporidae | Indian hare | <i>Lepus nigricollis</i> | LC | Decreasing | Not available | 2 |
| | Ochotonidae | Himalayan pika | <i>Ochotona roylei</i> | LC | Stable | Not available | 3 |
| Pholidota | Manidae | Indian pangolin | <i>Manis crassicaudata</i> | EN | Decreasing | Not available | 7 |
| Primates | Cercopithecidae | Rhesus macaque | <i>Macaca mulatta</i> | LC | Unknown | Not available | 18 |
| Rodentia | Cricetidae | Burrowing vole | <i>Hyperacrius fertilis</i> | NT | Unknown | Not available | 1 |
| | | Murree vole | <i>Hyperacrius wynnei</i> | LC | Unknown | Not available | 1 |
| | Gliridae | Forest dormouse | <i>Dryomys nitedula</i> | LC | Unknown | Not available | 2 |
| | Hystriidae | Indian crested porcupine | <i>Hystrix indica</i> | LC | Stable | Not available | 6 |
| | Muridae | Himalayan rat | <i>Rattus nitidus</i> | LC | Stable | Not available | 4 |
| | | Himalayan wood mouse | <i>Apodemus rusiges</i> | LC | Unknown | Not available | 3 |
| | | House mouse | <i>Mus musculus</i> | LC | Stable | Not available | 7 |
| | | Indian mole rat | <i>Bandicota bengalensis</i> | LC | Increasing | Not available | 1 |
| | | Turkestan rat | <i>Rattus turkestanicus</i> | LC | Unknown | Not available | 4 |
| | Suiridae | Golden marmot | <i>Marmota caudata aurea</i> | Not available | Not available | Not available | 6 |
| | | Himalayan marmot | <i>Marmota himalayana</i> | LC | Unknown | Not available | 1 |
| | | Kashmir marmot | <i>Marmota caudata</i> | LC | Unknown | Not available | 2 |
| | | Northern palm squirrel | <i>Funambulus pennantii</i> | LC | Unknown | Not available | 1 |
| | | Small kashmir flying squirrel | <i>Hylopetes fimbriatus</i> | LC | Unknown | Not available | 1 |
| Soricidae | Eulipotyphla | Grey shrew | <i>Crocidura attenuata</i> | LC | Unknown | Not available | 1 |
| | | Musk shrew | <i>Suncus murinus</i> | LC | Stable | Not available | 3 |
| | | Pygmy shrew | <i>Suncus etruscus</i> | LC | Unknown | Not available | 1 |
| Reptiles | | | | | | | |
| Squamata | Agamidae | Agrore valley agama | <i>Laudakia agrorensis</i> | Not available | Not available | Not available | 9 |
| | | Black rock agama | <i>Laudakia melanura melanura</i> | Not available | Not available | Not available | 17 |
| | | Blue rock agama | <i>Laudakia tuberculata</i> | Not available | Not available | Not available | 12 |
| | | Himalayan agama | <i>Laudakia himalayana</i> | Not available | Not available | Not available | 19 |
| | | North-pakistan agama | <i>Laudakia pakistanica auffenbergi</i> | Not available | Not available | Not available | 32 |
| | | | <i>Laudakia pakistanica khani</i> | Not available | Not available | Not available | 24 |

Table continued on next page.....

| Order | Family | Common name | Scientific name | IUCN status | Trend | Occurrence | Count |
|------------|-------------|---------------------------|--|---------------|---------------|---------------|-------|
| | | | <i>Laudakia pakistanica pakistanica</i> | Not available | Not available | Not available | 37 |
| | Colubridae | Cliff racer | <i>Platyceps rhodorachis rhodorachis</i> | Not available | Not available | Not available | 1 |
| | Elapidae | Brown cobra | <i>Naja oxiana</i> | DD | Not available | Not available | 1 |
| | | Common krait | <i>Bungarus caeruleus caeruleus</i> | Not available | Not available | Not available | 3 |
| | Gekkonidae | Baltistan gecko | <i>Altigekko stoliczkai</i> | LC | Stable | Not available | 2 |
| | | Chitral gecko | <i>Mediodactylus walli</i> | LC | Unknown | Not available | 1 |
| | | Fat-tailed gecko | <i>Eublepharis macularius</i> | Not Available | Not available | Not available | 3 |
| | | Karakorum gecko | <i>Altigekko boehmei</i> | Not Available | Not available | Not available | 4 |
| | Scincidae | Himalayan skink | <i>Scincella himalayana</i> | Not Available | Not available | Not available | 3 |
| | | Orange-tailed skink | <i>Novoeumeces blythianus</i> | Not Available | Not available | Not available | 1 |
| | Typhlopidae | Kashmir blind snake | <i>Typhlops diardii platyventris</i> | Not Available | Not available | Not available | 1 |
| | Varanidae | Bengal monitor | <i>Varanus bengalensis</i> | LC | Decreasing | Not available | 19 |
| | | Yellow monitor | <i>Varanus flavescens</i> | LC | Not available | Not available | 1 |
| | Viperidae | Himalayan pit viper | <i>Gloydius himalayanus</i> | Not Available | Not available | Not available | 1 |
| Testudines | Geoemydidae | Yellow-spotted mud turtle | <i>Geoclemys hamiltonii</i> | EN | Decreasing | Not available | 3 |

Supplementary Table II. Species recorded in hotspot areas.

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|-------------------|---------------------------------|--------------------------------------|-----------|-----------|-----------|
| Amphibians | | | | | |
| 1 | Southeast asian toad | <i>Bufo melanostictus</i> | + | | |
| 2 | Hazara toad | <i>Bufo melanostictus hazarensis</i> | + | | |
| 3 | Siachine toad | <i>Bufo siachenensis</i> | | + | |
| 4 | Indus valley toad | <i>Bufo stomaticus</i> | + | | |
| 5 | Green toad | <i>Bufo viridis</i> | + | | |
| 6 | Skittering frog | <i>Euphlyctis cyanophlyctis</i> | + | | |
| 7 | Alpine cricket frog | <i>Fejervarya limnocharis</i> | + | | |
| 8 | Hazara fast-flowing stream frog | <i>Paa hazarensis</i> | + | | |
| 9 | Hazara torrent frog | <i>Allopaa hazarensis</i> | + | | |
| Birds | | | | | |
| 10 | Snow partridge | <i>Lerwa lerwa</i> | + | + | + |
| 11 | Chukar | <i>Alectoris chukar</i> | + | + | + |
| 12 | Common quail | <i>Coturnix coturnix</i> | + | | + |
| 13 | Himalayan snowcock | <i>Tetraogallus himalayensis</i> | | | + |
| 14 | Grey francolin | <i>Francolinus pondicerianus</i> | + | | |
| 15 | Western tragopan | <i>Tragopan melanocephalus</i> | + | | |
| 16 | Koklass pheasant | <i>Pucrasia macrolopha</i> | + | | |
| 17 | Himalayan monal | <i>Lophophorus impejanus</i> | + | | + |

Table continued on next page.....

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|--------|-----------------------------|----------------------------------|-----------|-----------|-----------|
| 18 | Cheer pheasant | <i>Catreus wallichii</i> | + | | |
| 19 | Greylag goose | <i>Anser anser</i> | + | | + |
| 20 | Bar-headed goose | <i>Anser indicus</i> | + | | + |
| 21 | Ruddy shelduck | <i>Tadorna ferruginea</i> | + | + | + |
| 22 | Gadwall | <i>Anas strepera</i> | + | | |
| 23 | Eurasian wigeon | <i>Anas penelope</i> | + | | |
| 24 | Mallard | <i>Anas platyrhynchos</i> | + | | |
| 25 | Common teal | <i>Anas crecca</i> | + | | |
| 26 | Northern pintail | <i>Anas acuta</i> | + | | |
| 27 | Northern shoveler | <i>Anas clypeata</i> | + | | |
| 28 | Ferruginous pochard | <i>Aythya nyroca</i> | + | | |
| 29 | Tufted duck | <i>Aythya fuligula</i> | + | | |
| 30 | Common merganser | <i>Mergus merganser</i> | + | | |
| 31 | Eurasian wryneck | <i>Jynx torquilla</i> | + | | + |
| 32 | Himalayan woodpecker | <i>Dendrocopos himalayensis</i> | + | | + |
| 33 | Common hoopoe | <i>Upupa epops</i> | + | + | + |
| 34 | European roller | <i>Coracias garrulus</i> | + | | + |
| 35 | Common kingfisher | <i>Alcedo atthis</i> | | + | |
| 36 | White-throated kingfisher | <i>Halcyon smyrnensis</i> | + | | |
| 37 | Blue-cheeked bee-eater | <i>Merops persicus</i> | + | | + |
| 38 | European bee-eater | <i>Merops apiaster</i> | + | | |
| 39 | Eurasian cuckoo | <i>Cuculus canorus</i> | | + | |
| 40 | Rose-ringed parakeet | <i>Psittacula krameri</i> | + | | |
| 41 | Alpine swift | <i>Tachymarptis melba</i> | + | | |
| 42 | Common swift | <i>Apus apus</i> | + | + | + |
| 43 | House swift | <i>Apus affinis</i> | + | | |
| 44 | Eurasian scops owl | <i>Otus scops</i> | | | + |
| 45 | Collared owl | <i>Glaucidium brodiei</i> | + | | |
| 46 | Tawny owl | <i>Strix aluco</i> | + | | |
| 47 | Eurasian nightjar | <i>Caprimulgus europaeus</i> | + | + | + |
| 48 | Rock pigeon | <i>Columba livia</i> | + | + | + |
| 49 | Hill pigeon | <i>Columba rupestris</i> | + | + | + |
| 50 | Laughing dove | <i>Streptopelia senegalensis</i> | + | | |
| 51 | Spotted dove | <i>Streptopelia chinensis</i> | + | | |
| 52 | Eurasian collared dove | <i>Streptopelia decaocto</i> | + | | |
| 53 | Little bustard | <i>Tetrax tetrax</i> | + | | |
| 54 | Water rail | <i>Rallus aquaticus</i> | + | | |
| 55 | Little crane | <i>Porzana parva</i> | + | | + |
| 56 | Baillon's crane | <i>Porzana pusilla</i> | + | | |
| 57 | Common moorhen | <i>Gallinula chloropus</i> | + | + | |
| 58 | Chestnut-bellied sandgrouse | <i>Pterocles exustus</i> | + | | |

Table continued on next page.....

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|--------|------------------------|---------------------------------|-----------|-----------|-----------|
| 59 | Eurasian woodcock | <i>Scolopax rusticola</i> | | | + |
| 60 | Common snipe | <i>Gallinago gallinago</i> | + | | |
| 61 | Eurasian curlew | <i>Numenius arquata</i> | + | | |
| 62 | Common redshank | <i>Tringa totanus</i> | | | + |
| 63 | Green sandpiper | <i>Tringa ochropus</i> | + | | + |
| 64 | Wood sandpiper | <i>Tringa glareola</i> | | | + |
| 65 | Common sandpiper | <i>Actitis hypoleucos</i> | + | + | + |
| 66 | Little stint | <i>Calidris minuta</i> | + | | |
| 67 | Temminck's stint | <i>Calidris temminckii</i> | + | | + |
| 68 | Curlew sandpiper | <i>Calidris ferruginea</i> | + | | |
| 69 | Ruff | <i>Philomachus pugnax</i> | + | | + |
| 70 | Pheasant-tailed jacana | <i>Hydrophasianus chirurgus</i> | + | | |
| 71 | Northern lapwing | <i>Vanellus vanellus</i> | + | | |
| 72 | Red-wattled lapwing | <i>Vanellus indicus</i> | + | | + |
| 73 | Sociable lapwing | <i>Vanellus gregarius</i> | + | | |
| 74 | Indian skimmer | <i>Rynchops albicollis</i> | + | | |
| 75 | Brown-headed gull | <i>Larus brunnecephalus</i> | + | + | |
| 76 | Gull-billed tern | <i>Gelochelidon nilotica</i> | | | + |
| 77 | Caspian tern | <i>Sterna caspia</i> | + | | |
| 78 | Black kite | <i>Milvus migrans</i> | + | + | |
| 79 | Pallas's fish eagle | <i>Haliaeetus leucoryphus</i> | + | + | |
| 80 | Lammergeier | <i>Gypaetus barbatus</i> | + | + | + |
| 81 | Egyptian vulture | <i>Neophron percnopterus</i> | + | | + |
| 82 | Himalayan griffon | <i>Gyps himalayensis</i> | + | + | + |
| 83 | Eurasian marsh harrier | <i>Circus aeruginosus</i> | + | | + |
| 84 | Hen harrier | <i>Circus cyaneus</i> | + | + | |
| 85 | Pallid harrier | <i>Circus macrourus</i> | + | | + |
| 86 | Shikra | <i>Accipiter badius</i> | + | | |
| 87 | Eurasian sparrowhawk | <i>Accipiter nisus</i> | + | + | + |
| 88 | White-eyed buzzard | <i>Butastur teesa</i> | + | | |
| 89 | Long-legged buzzard | <i>Buteo rufinus</i> | + | + | |
| 90 | Booted eagle | <i>Hieraaetus pennatus</i> | + | | |
| 91 | Common kestrel | <i>Falco tinnunculus</i> | + | + | + |
| 92 | Eurasian hobby | <i>Falco subbuteo</i> | + | + | |
| 93 | Saker falcon | <i>Falco cherrug</i> | + | | |
| 94 | Peregrine falcon | <i>Falco peregrinus</i> | + | | |
| 95 | Little grebe | <i>Tachybaptus ruficollis</i> | | + | |
| 96 | Little egret | <i>Egretta garzetta</i> | + | | |
| 97 | Grey heron | <i>Ardea cinerea</i> | + | | + |
| 98 | Indian pond-heron | <i>Ardeola grayii</i> | + | | |
| 99 | Little bittern | <i>Ixobrychus minutus</i> | + | | |

Table continued on next page.....

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|--------|-----------------------------|-------------------------------------|-----------|-----------|-----------|
| 100 | Glossy ibis | <i>Plegadis falcinellus</i> | | + | |
| 101 | Black stork | <i>Ciconia nigra</i> | + | | |
| 102 | Rufous-tailed shrike | <i>Lanius isabellinus</i> | + | | |
| 103 | Long-tailed shrike | <i>Lanius schach</i> | + | + | + |
| 104 | Southern grey shrike | <i>Lanius meridionalis</i> | + | | + |
| 105 | Yellow-billed chough | <i>Pyrrhocorax graculus</i> | + | + | + |
| 106 | Rook | <i>Corvus frugilegus</i> | | + | + |
| 107 | Carrion crow | <i>Corvus corone</i> | | + | + |
| 108 | Large-billed crow | <i>Corvus macrorhynchos</i> | + | + | + |
| 109 | Common raven | <i>Corvus corax</i> | | + | + |
| 110 | Eurasian golden oriole | <i>Oriolus oriolus</i> | + | + | + |
| 111 | Rosy minivet | <i>Pericrocotus roseus</i> | + | | |
| 112 | Long-tailed minivet | <i>Pericrocotus ethologus</i> | + | + | + |
| 113 | Black drongo | <i>Dicrurus macrocercus</i> | + | | |
| 114 | Ashy drongo | <i>Dicrurus leucophaeus</i> | + | | |
| 115 | Asian paradise-flycatcher | <i>Terpsiphone paradisi</i> | + | | |
| 116 | White-throated dipper | <i>Cinclus cinclus</i> | + | | |
| 117 | Brown dipper | <i>Cinclus pallasii</i> | + | + | + |
| 118 | Rufous-tailed rock thrush | <i>Monticola saxatilis</i> | + | | |
| 119 | Blue-capped rock thrush | <i>Monticola cinclorhynchus</i> | + | | |
| 120 | Blue rock thrush | <i>Monticola solitarius</i> | + | + | + |
| 121 | Blue whistling thrush | <i>Myophonus caeruleus</i> | + | + | |
| 122 | Tickell's thrush | <i>Turdus unicolor</i> | + | | |
| 123 | Eurasian blackbird | <i>Turdus merula</i> | + | | |
| 124 | Chestnut thrush | <i>Turdus rubrocanus</i> | + | | |
| 125 | Ultramarine flycatcher | <i>Ficedula superciliaris</i> | + | | |
| 126 | Slaty-blue flycatcher | <i>Ficedula tricolor</i> | + | | |
| 127 | Bluethroat | <i>Luscinia svecica</i> | + | + | + |
| 128 | Indian blue robin | <i>Luscinia brunnea</i> | + | | |
| 129 | Rufous-backed redstart | <i>Phoenicurus erythronota</i> | + | | + |
| 130 | Blue-capped redstart | <i>Phoenicurus coeruleocephalus</i> | | | + |
| 131 | Black redstart | <i>Phoenicurus ochruros</i> | + | | |
| 132 | White-winged redstart | <i>Phoenicurus erythrogaster</i> | + | | + |
| 133 | Blue-fronted redstart | <i>Phoenicurus frontalis</i> | + | | |
| 134 | White-capped water redstart | <i>Chaimarrornis leucocephalus</i> | + | | + |
| 135 | Plumbeous water redstart | <i>Rhyacornis fuliginosus</i> | + | | + |
| 136 | Little forktail | <i>Enicurus scouleri</i> | | | + |
| 137 | Common stonechat | <i>Saxicola torquata</i> | + | + | + |
| 138 | Pied bushchat | <i>Saxicola caprata</i> | + | | |
| 139 | Grey bushchat | <i>Saxicola ferrea</i> | + | | |
| 140 | Variable wheatear | <i>Oenanthe picata</i> | + | + | + |
| 141 | Pied wheatear | <i>Oenanthe pleschanka</i> | + | + | + |

Table continued on next page.....

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|--------|--------------------------|----------------------------------|-----------|-----------|-----------|
| 142 | Isabelline wheatear | <i>Oenanthe isabellina</i> | + | | |
| 143 | Brahminy starling | <i>Sturnus pagodarum</i> | + | | |
| 144 | Common myna | <i>Acridotheres tristis</i> | + | | |
| 145 | Wallcreeper | <i>Tichodroma muraria</i> | + | + | + |
| 146 | Winter wren | <i>Troglodytes troglodytes</i> | + | + | + |
| 147 | Rufous-naped tit | <i>Parus rufonuchalis</i> | + | | + |
| 148 | Great tit | <i>Parus major</i> | + | + | + |
| 149 | White-cheeked tit | <i>Aegithalos leucogenys</i> | + | | |
| 150 | Pale martin | <i>Riparia diluta</i> | + | | |
| 151 | Eurasian crag martin | <i>Hirundo rupestris</i> | + | + | + |
| 152 | Barn swallow | <i>Hirundo rustica</i> | + | | + |
| 153 | Northern house martin | <i>Delichon urbicum</i> | | + | + |
| 154 | Asian house martin | <i>Delichon dasypus</i> | + | + | + |
| 155 | Himalayan bulbul | <i>Pycnonotus leucogenys</i> | + | | |
| 156 | Red-vented bulbul | <i>Pycnonotus cafer</i> | + | | |
| 157 | Grey-breasted prinia | <i>Prinia hodgsonii</i> | + | | |
| 158 | Oriental white-eye | <i>Zosterops palpebrosus</i> | + | | |
| 159 | White-browed tit warbler | <i>Leptopoeile sophiae</i> | + | | + |
| 160 | Lesser whitethroat | <i>Sylvia curruca</i> | | + | |
| 161 | Long-billed bush warbler | <i>Bradypterus major</i> | | | + |
| 162 | Blyth's reed warbler | <i>Acrocephalus dumetorum</i> | + | | |
| 163 | Common chiffchaff | <i>Phylloscopus collybita</i> | + | | + |
| 164 | Mountain chiffchaff | <i>Phylloscopus sindianus</i> | | + | + |
| 165 | Tickell's leaf warbler | <i>Phylloscopus affinis</i> | + | | + |
| 166 | Brook's leaf warbler | <i>Phylloscopus subviridis</i> | + | | + |
| 167 | Hume's warbler | <i>Phylloscopus humei</i> | + | | + |
| 168 | Greenish warbler | <i>Phylloscopus trochiloides</i> | + | | + |
| 169 | Tytler's leaf warbler | <i>Phylloscopus tytleri</i> | + | | |
| 170 | Streaked laughingthrush | <i>Garrulax lineatus</i> | + | + | + |
| 171 | Bimaculated lark | <i>Melanocorypha bimaculata</i> | | | + |
| 172 | Crested lark | <i>Galerida cristata</i> | + | | + |
| 173 | Eurasian skylark | <i>Alauda arvensis</i> | + | + | + |
| 174 | Oriental skylark | <i>Alauda gulgula</i> | + | | + |
| 175 | House sparrow | <i>Passer domesticus</i> | + | + | + |
| 176 | Spanish sparrow | <i>Passer hispaniolensis</i> | + | | + |
| 177 | Russet sparrow | <i>Passer rutilans</i> | + | | |
| 178 | Eurasian tree sparrow | <i>Passer montanus</i> | + | | |
| 179 | White wagtail | <i>Motacilla alba</i> | + | + | + |
| 180 | Citrine wagtail | <i>Motacilla citreola</i> | + | + | + |
| 181 | Yellow wagtail | <i>Motacilla flava</i> | + | | + |
| 182 | Grey wagtail | <i>Motacilla cinerea</i> | + | + | + |

Table continued on next page.....

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|----------------|-------------------------------|--------------------------------------|-----------|-----------|-----------|
| 183 | Tawny pipit | <i>Anthus campestris</i> | + | | |
| 184 | Long-billed pipit | <i>Anthus similis</i> | + | | |
| 185 | Upland pipit | <i>Anthus sylvanus</i> | + | | |
| 186 | Tree pipit | <i>Anthus trivialis</i> | + | + | + |
| 187 | Rosy pipit | <i>Anthus roseatus</i> | + | + | + |
| 188 | Water pipit | <i>Anthus spinoletta</i> | + | + | + |
| 189 | Alpine accentor | <i>Prunella collaris</i> | + | + | + |
| 190 | Yellow-breasted greenfinch | <i>Carduelis spinoides</i> | + | | |
| 191 | Eurasian goldfinch | <i>Carduelis carduelis</i> | | + | |
| 192 | Twite | <i>Carduelis flavirostris</i> | | + | |
| 193 | Common rosefinch | <i>Carpodacus erythrinus</i> | + | + | |
| 194 | Red-fronted rosefinch | <i>Carpodacus puniceus</i> | + | | + |
| 195 | Orange bullfinch | <i>Pyrrhula aurantiaca</i> | + | + | |
| 196 | White-winged grosbeak | <i>Mycerobas carnipes</i> | + | | |
| 197 | Crested bunting | <i>Melophus lathamii</i> | + | | |
| 198 | Rock bunting | <i>Emberiza cia</i> | + | + | + |
| 199 | Reed bunting | <i>Emberiza schoeniclus</i> | + | | |
| 200 | Snowy owl | <i>Nyctea scandiaca</i> | + | | |
| Mammals | | | | | |
| 201 | Musk shrew | <i>Suncus murinus</i> | + | | |
| 202 | Common pipistrelle | <i>Pipistrellus pipistrellus</i> | | | + |
| 203 | Asian barbastelle | <i>Barbastella leucomelas</i> | | + | |
| 204 | Rhesus macaque | <i>Macaca mulatta</i> | + | | |
| 205 | Indian wild dog | <i>Cuon alpinus</i> | | + | |
| 206 | Asiatic black bear | <i>Ursus thibetanus</i> | + | | |
| 207 | Alpine weasel | <i>Mustela altaica</i> | | | + |
| 208 | Common otter | <i>Lutra lutra</i> | + | + | + |
| 209 | Himalayan palm civet | <i>Paguma larvata wroughtoni</i> | + | | |
| 210 | Leopard cat | <i>Prionailurus bengalensis</i> | | | + |
| 211 | Panther | <i>Panthera pardus</i> | + | | |
| 212 | Snow leopard | <i>Panthera uncia</i> | | | + |
| 213 | Himalayan musk deer | <i>Moschus chrysogaster</i> | + | + | + |
| 214 | Himalayan ibex | <i>Capra ibex siberica</i> | | + | |
| 215 | Astor markhor | <i>Capra falconeri</i> | | + | |
| 216 | Pir panjal markhor | <i>Capra falconeri cashmiriensis</i> | + | | |
| 217 | Himalayan pika | <i>Ochotona roylei</i> | + | + | + |
| 218 | Small Kashmir flying squirrel | <i>Hylopetes fimbriatus</i> | | + | |
| 219 | Kashmir marmot | <i>Marmota caudata</i> | + | | + |
| 220 | Indian crested porcupine | <i>Hystrix indica</i> | + | | |
| 221 | Forest dormouse | <i>Dryomys nitidula</i> | + | | |
| 222 | Himalayan wood mouse | <i>Apodemus rusiges</i> | + | + | + |

Table continued on next page.....

| S. No. | Common name | Scientific name | Hotspot A | Hotspot B | Hotspot C |
|-----------------|----------------------|---|-----------|-----------|-----------|
| 223 | Turkestan rat | <i>Rattus turkestanicus</i> | + | + | + |
| 224 | Himalayan rat | <i>Rattus nitidus</i> | + | | |
| 225 | House mouse | <i>Mus musculus</i> | + | + | + |
| 226 | Murree vole | <i>Hyperacrius wynnei</i> | + | | |
| 227 | Burrowing vole | <i>Hyperacrius fertilis</i> | + | | |
| 228 | Shapu | <i>Ovis vignei vignei</i> | | | + |
| 229 | Himalayan agama | <i>Laudakia himalayana</i> | | + | + |
| Reptiles | | | | | |
| 230 | North-Pakistan agama | <i>Laudakia pakistanica pakistanica</i> | | + | |
| 231 | North-Pakistan agama | <i>Laudakia pakistanica auffenbergi</i> | | + | |
| 232 | Blue rock agama | <i>Laudakia tuberculata</i> | + | | + |
| 233 | Baltistan gecko | <i>Altigekko stoliczkai</i> | | + | |
| 234 | Chitral gecko | <i>Mediodactylus walli</i> | + | | |
| 235 | Himalayan skink | <i>Scincella himalayana</i> | + | + | |
| 236 | Bengal monitor | <i>Varanus bengalensis</i> | + | | |
| 237 | Common krait | <i>Bungarus caeruleus caeruleus</i> | + | | |
| 238 | Brown cobra | <i>Naja oxiana</i> | + | | |